

Original Research Article

Prevalence and determinants of antimalarial self-medication in southern Benin

ABSTRACT

Aims: To determine the prevalence of antimalarial self-medication and identify its determinants in Comé, Southern Benin households

Study design: Cross-sectional community based study.

Place and Duration of Study: Benin southern setting, from April to July 2015.

Methodology: It was a cross-sectional study conducted from 1st to 14th July 2015 which involved 480 households randomly selected. Data on socioeconomic and demographic factors, the use of antimalarial drugs, health system and knowledge regarding malaria and dangers of self-medication were collected by questionnaire. Multivariate logistic regression was used to identify determinants of antimalarial self-medication in households.

Results: The prevalence antimalarial self-medication in households was 69.77% [CI95: (65.64; 73.89)]. The use of antimalarial drug from illicit market was reported in 83.75%. In 28.73% of cases, the main anti-malarial drug used in self-medication was artemisinin-based combination. Determinants of antimalarial self-medication were level of economic status: quintiles poor (1 and 2) versus the richest quintile : OR = 6.50 [95%IC (3.43 – 10.68)] ; quintiles less poor (3 et 4) versus the richest quintile : OR = 3,05 [95%IC (1,49 – 6,25)]; knowledge of the dangers of antimalarial self-medication, knowledge versus ignorance, OR = 0.26 [95%IC (0.14 – 0.48)], knowledge of dangers associated with street drugs: knowledge versus ignorance; OR = 0,40 [95% (0,19 – 0,86)], knowledge of consequences of poorly treated malaria, knowledge versus ignorance: OR = 0,36 [95%IC (0,16 – 0,8)].

Conclusion: The results point out the high prevalence of antimalarial self-medication in households in Comé, Southern Benin. Strengthening knowledge on consequences of antimalarial self-medication, street drugs consumption and poorly treated malaria is needed to avert antimalarial self-medication practices in Comé in Southern Benin

Keywords: antimalarial self-medication, prevalence, determinants, households, Benin

INTRODUCTION

Malaria is the commonest cause of fever and morbidity in the tropics and a significant source of mortality, especially among infants and young children in Sub-Saharan Africa [1-3]. To treat this disease, some individuals practice antimalarial self-medication [4-8]. According to the World Health Organization, self-medication refers to the use of a drug, on its own initiative or on relatives' advices, in order to treat a disease or symptom without recourse to a health worker [9]. Self-medication includes both modern medicines as traditional medicines [10, 11]. The irrational use of antimalarial by self-medication exposes to drug poisoning and adverse drug interactions. The practice of self-medication and some unnecessary presumptive treatment prescribed by health professionals contribute to the emergence of drug resistance [3, 4].

With the wide drug resistance to chloroquine, several sub-Saharan countries including Benin have adopted the combination therapy base on artemisinin (ACTs) as recommended by WHO for the effective treatment of malaria and for preventing the development of drug resistance [12]. The fight against the practice of antimalarial self-medication may help preventing the early onset of resistance to ACTs in parasites. In Benin, studies reported that antimalarial self-medication was widespread in urban area [13] but there is a dearth of information in semi-urban and rural settings.

27 **2. OBJECTIVE**

28 The objective of the study was to determine the prevalence of antimalarial self-medication
29 and examine its determinants in household in a semi-urban area and its rural outskirts in
30 southern Benin.

31

32 **3. SUBJECTS AND METHODS**

33

34 **3.1. Setting**

35 The study was carried out in the small size city of Com  (80 km from Cotonou, the economic capital of
36 Benin) and its rural outskirts. Com  includes 38 villages and city neighborhoods. According to census
37 in 2013, the population of Com  was estimated at 79 665 inhabitants. The epidemiological profile is
38 characterized by the predominance of communicable diseases, mainly malaria and emergence of
39 some non-communicable diseases (hypertension, diabetes and obesity).

40 To improve timely access to malaria treatment, the National Program against Malaria has
41 strengthened the laboratory capacity for malaria biologic diagnosis at public, private and community
42 levels by increasing availability of the rapid diagnostic test for malaria at health centers and in the
43 community (for community health workers). With the updated management guidance of malaria cases
44 in February 2011, laboratory confirmation of malaria before treatment is mandatory in health facilities
45 and in the community [14].

46

47 **3.2. Study design and subjects**

48 This was a cross-sectional study conducted from April to July 2015. The study population consisted of
49 the heads of households in Com .

50

51 **3.3. Sample size and subjects selection**

52 **3.3.1. Sample size**

53 The minimum sample size estimated was 480 heads of households using Schwartz formula
54 and assuming α error of 5%, an accuracy of 4%, the cluster sampling effect of 3 and the
55 prevalence of antimalarial self-medication reported in in urban areas (93.1%) [13].

56 **3.3.2. Selection of participants**

57 Randomised multistage cluster sampling technique was used to select participants: the villages or city
58 neighborhoods, concessions and finally, heads of households.

- 59 - First stage: a total of 30 villages or city neighborhoods were selected by simple random
60 sampling among the 38 villages and city neighborhoods.
- 61 - Second stage: From the center of each selected village or city neighborhood, the investigator
62 has chosen the direction indicated by the tip of launched pen. In this direction, concessions
63 were numbered and the first to visit was selected by simple random. From this first house
64 visited, the investigator went from concession to concession until the expected number of
65 house in each village.
- 66 - Third stage: in the selected concession, the head of household or his representative over 18
67 years was identified. If the concession had two or more eligible household chiefs, one was
68 randomly selected. Clusters including 16 (480/30) participants were formed in each selected
69 village or city neighbour.

70

71 **3.4. The study variables**

72 **3.4.1. The dependent variable**

73 The dependent variable was the practice of antimalarial self-medication. In the study, all personal
74 initiative of consuming one or more antimalarial drugs without prior medical assessment or health
75 worker advices was considered as self-medication. This could concern CTA, quinine, sulfadoxin-
76 pyrimethamin, chloroquine and other preparations such as herbal beverages and traditional products
77 intended for treating malaria. The period considered covered the 30 days past the day before the
78 survey.

79

80

3.4.2. The independent variables

Independent variables included:

- Socio-economic and demographic factors: age, sex, religion, educational level, occupation, socioeconomic status and household size,
- Behavioural factors: health services utilization, preference for self-treatment, opinion on health care fees;
- Factors related to households head's knowledge: knowledge of the dangers of self-medication, knowledge of symptoms of malaria, knowledge of the consequences of poorly treated malaria and knowledge of adverse effects associated with street drugs consumption.
- Environmental factors: Distance from home to the illicit drug market and exposure to advertising of antimalarial traditional medicine products.
- Factors related to the health system: the distance from home to health facilities and satisfaction regarding quality of health care.

3.5. Data collection procedures

Questionnaires were used for data collection in study participants. The questionnaires were pre-tested in other health district by checking the answerability and the appropriateness of questions. The result of the pre-test informed the investigator on the required adjustments in the questionnaire used.

3.6. Data analysis

Data were analyzed by the software STATA11.0. A weighted variable was created to take into account the design of the survey. Proportions were estimated and odds ratio (OR) were calculated with confidence interval of 95% to measure the strength of the association between antimalarial self-medication and independent variables. Univariate analyzes were used initially to identified factors associated with the practice of antimalarial self-medication. To identify determinants of the practice of antimalarial self-treatment, multivariate analysis was performed taking into account only the factors associated with the practice of self-medication in the univariate analysis was. The significance level was set at $p < 0.05$.

3.7. Ethical considerations

The objectives of the study were explained to participants. Voluntary free and informed oral consent was obtained before starting the interview. This consent claims that participants are not at risk by refusing to participate in the survey or stopping their collaboration during the study. The confidentiality and anonymity of information collected were ensured.

4. results

4.1. Socioeconomic and demographic characteristics of the participants

Table I summarizes socioeconomic and demographic characteristics of the participants. Among the participants, 46.5% were women and 39.14% of the respondents aged 31 - 46 years. Participants with primary education accounted for 31.8% of respondents. At least one respondent in five, (20.0%) belonged to the second quintile of economic well-being. Of the respondents, 25.9% were traders.

4.2. Behavioral and environmental and knowledge of participants

Table II shows that among the participants, 54.7% preferred to consume antimalarial drug that was not prescribed by a health worker and 79.4% of the respondents showed a poor knowledge of dangers associated with antimalarial self-medication. Among the 480 participants, 49.3% failed to identify symptoms of malaria, 79.2% had poor knowledge of the consequences of poorly treated malaria and 68.1% were unaware of the risks associated with street drugs consumption.

Among the participants, 55.2% declared to reside close to illicit drug mark and 41.6% were exposed to advertisements on traditional medicine products used as antimalarial during the month preceding the survey. Furthermore, 76.2% of respondents lived within five kilometers of a health center and 61.6% were not satisfied with care provided in health centers (table II)

140 Table I: Socioeconomic and demographic characteristics of participants, Comé, Southern Benin,
141 2015, (n=480)
142

Characteristics	Absolute frequency	% weighted
Age		
15-30 years	187	38,63
31-46 years	178	39,14
47-61 years	73	13,66
62-77 years	38	7,12
78-93 years	4	1,45
Sex		
Women	227	46,50
Men	253	53,50
Education		
No schooling	139	30,58
Primary	138	31,83
Secondary	167	30,56
University	36	7,03
Religion		
Catholicism	145	32,81
Protestant	20	3,91
Celestial Christianity	64	10,67
Muslim	18	3,07
Animism	142	31,62
Atheist	91	17,92
Occupation		
Official	46	8,08
Trader	121	25,97
Farmer	45	11,12
Student	78	14,12
Motor cycle taxi driver	46	9,48
House wife	73	17,33
Unemployed	19	4,00
Other	52	9,90
Economic status		
Quintile 1	96	20,02
Quintile 2	72	20,05
Quintile 3	101	19,95
Quintile 4	93	20,29
Quintile 5	118	19,69

143

144

145

146

147

148

149

150

151

152

153

154

155 Table II: Behavioral and environmental characteristics and knowledge of participants, Comé,
156 Southern Benin, 2015, (n=480)
157

Characteristics of participants	Absolute frequency	% weighted
Preference for non-prescribed by a health worker medicines		
Yes	273	54,76
No	207	45,24
Utilization of health facilities		
Yes	259	48,75
No	221	51,25
Will to not pay medical examination fees		
Yes	201	48,73
No	279	51,27
Knowledge of danger of antimalarial self-medication		
Yes	109	20,58
No	371	79,42
Knowledge of malaria symptoms		
Yes	260	50,67
No	220	49,33
Knowledge of consequences of poorly treated malaria		
Yes	107	20,76
No	373	79,24
Knowledge of dangers of street drugs consumption		
Yes	165	31,84
No	315	68,16
Distance from home to health facilities		
Long (>5Km)	114	23,26
Short (<5Km)	366	76,74
The proximity to the illicit market		
Yes	245	55,20
No	235	44,80
Advertising for traditional medicine products		
Yes	216	41,66
No	264	58,34
Satisfaction with health care		
Yes	267	61,63
No	213	38,37
Trust in pharmacists		
Yes	172	35,90
No	308	64,10

158

159 4.3. Antimalarial self-medication during the last month

160 Among participants, 69.7% [95% CI (65.64 - 73.89)] reported that at least one member of their
161 household has practiced antimalarial self-medication during the thirty days preceding the survey.
162 Among households who practiced self-medication, 83.7% have bought drug at illicit market.
163 Antimalarial drug that have been used for self-medication were: CTA (28.7%), quinine (26.5%),

chloroquine (23.3%), sulfadoxin-pyrimethamin (8.38%) and other preparations such as herbal beverages and other traditional products intended to treat malaria (13.0%).

4.4. Factors associated with antimalarial self-medication

Univariate analyzes identified twelve variables that were associated with the practice of antimalarial self-medication as shown in table III.

Table IV presents the determinants of self-medication antimalarial according to the multivariate model. Determinants of antimalarial self-medication were: the level of socioeconomic welfare quintiles poor (1 and 2) versus the richest quintile OR = 6.50 [95% CI (3.43 to 10.68)], less poor quintiles (3 and 4) versus the richest quintile OR = 3.05 [95% CI (1.49 to 6.25)]; knowledge of the dangers of self-medication antimalarial: knowledge versus ignorance: OR = 0.26 [95% CI (0.14 to 0.48)]; knowledge of risks associated with street drugs consumption, knowledge versus ignorance: OR = 0.40 [95% (0.19 to 0.86)] and knowledge of the consequences of a poorly treated malaria, knowledge versus ignorance: OR = 0.36 [95% CI (0.16 to 0.8)].

Variables such as age, sex, preference for drugs non-prescribed by a health worker, knowledge of symptoms of malaria, satisfaction with care provided were not associated with the practice of self-medication antimalarial.

Tableau III: Factors associated with antimalarial self-medication in households in Comé, Southern Benin, 2015, univariate analysis (n=480)

Independent variables	Antimalarial self-medication				P-value
	Yes	No	OR	IC ₉₅ %	
Education					
No schooling	101	38	1		
Primary	99	39	1,08	[0,63 – 1,85]	0,752
Secondary	94	73	0,68	[0,34 – 1,38]	0,281
University	9	27	0,083	[0,02 – 0,28]	0,000
Religion					
Catholicism	81	64	1		
Protestant	15	5	3,59	[0,96 – 13,43]	0,056
Celestial Christianity	38	26	0,80	[0,40 – 1,57]	0,509
Muslim	10	8	1,58	[0,40 – 6,21]	0,500
Animist	109	33	1,94	[1,05 – 3,58]	0,035
Atheist	50	41	1,11	[0,52 – 2,37]	0,771
Occupation					
Official	11	35	1		
Trader	86	35	5,96	[2,17 – 16,3]	0,001
Farmer	33	12	6,31	[2,64 – 15,06]	0,000
Student	40	38	1,57	[0,30 – 8,12]	0,578
Motor cycle taxi driver	37	9	4,5	[0,69 – 29,09]	0,110
House wife	52	21	6,22	[2,03 – 19,07]	0,002
Unemployed	13	6	4,51	[1,30 – 15,65]	0,019
Others	31	21	2,69	[0,54 – 13,34]	0,214
Socioeconomic status					
Quintile 5	43	75	1		
Quintile 1	72	24	5,58	[2,49 – 12,49]	
Quintile 2	59	13	8,05	[3,46 – 18,75]	0,000
Quintile 3	70	31	5,00	[2,33 – 10,73]	0,000
Quintile 4	59	34	3,23	[1,64 – 6,35]	0,000
Utilization of health facilities					
No	170	51	1		
Yes	133	126	0,50	[0,27 – 0,93]	0,03
Will to not to pay medical examinations fees					
No	161	118	1		
Yes	142	59	1,95	[1,33 – 2,85]	0,001
Knowledge of danger of antimalarial self-medication					
No	263	108	1		
Yes	40	69	0,23	[0,13 – 0,41]	0,000
Knowledge of consequences of poorly treated malaria					
No	253	120	1		
Yes	50	57	0,35	[0,19 – 0,64]	0,001

Independent variables	Antimalarial self-medication				
	Yes	No	OR	IC ₉₅ %	P-value
Knowledge of dangers of street drugs consumption					
Yes	182	133	1		
No	121	44	2,43	[1,20 – 4,85]	0,015
Distance from home to health facilities					
Long (>5Km)	90	24	1		
Short (<5Km)	213	153	0,32	[0,15 – 0,69]	0,005
Proximity of illicit drug market					
No	111	124	1		
Yes	192	53	3,59	[1,70 – 7,18]	0,001
Advertising for traditional medicine products					
No	188	76	1		
Yes	115	101	0,53	[0,31 – 0,91]	0,025

Table IV: Determinants of antimalarial self-medication in households in Comé, Southern Benin, multivariate analysis, 2015 (n=480)

Independent variables	OR	IC ₉₅ %	P
Economic status			
Quintile 5	1		
Quintile 1 et 2	6,05	[3,43 – 10,68]	<0,000
Quintile 3 et 4	3,05	[1,49 – 6,25]	0,003
Knowledge of consequences of poorly treated malaria			
Non	1		
Oui	0,36	[0,16 – 0,8]	0,014
Knowledge of danger of antimalarial self-medication			
Non	1		
Oui	0,26	[0,14 – 0,48]	<0,000
Knowledge of dangers of street drugs consumption			
Non	1		
Oui	0,40	[0,19 – 0,86]	0,021

5. DISCUSSION

The study examined the prevalence and determinants of antimalarial self-medication in a semi-urban city and its rural outskirts in southern Benin. The prevalence of antimalarial self-medication among households was 69.7% and CTA (28.7%), quinine (26.5%) and chloroquine (23.3%) were the most concerned drugs. Determinants of antimalarial self-medication were socioeconomic status, knowledge of the dangers of antimalarial self-medication, knowledge of dangers associated with street drugs consumption and knowledge of the consequences of poorly treated malaria.

5.1. Prevalence of antimalarial self-medication in households

The prevalence of antimalarial self-medication reported in this study is lower than 93.1% previously reported in Benin in children under five years [14] and (89.6%) in in Kisangani [15] and (99%) in Lubumbashi [5] in Democratic Republic of Congo. The reported prevalence in this study is similar to 72.7% observed in a survey conducted in Burkina Faso in 2014 [16]. Lower prevalence were observed among children under five in Ghana (16.8%) [17] and in Togo (42%) [18]. Inconsistences in results could be explained by the fact that studies had different targets, geographic scope and participants' ages. Indeed, the study previously carried out in Benin involved children under 5 years [13] as well as the Ghana study [17] while the present study concerned adults. The study in Burkina Faso was carried out in hospital [16].

5.2. Factors associated with self-medication antimalarial

5.2.1. Socioeconomic and demographic factors

The low economic status was a determinant of the practice of antimalarial self-medication. Tshomba et al., had observed that higher monthly income to 60,000 FCFA protected against the practice of self-medication [15]. Also Dargicho et al. reported in Ethiopia that the monthly income was negatively and significantly associated with antimalarial self-medication [19]. Indeed, the person with low economic status prefers to buy drugs at illicit market to avoid paying medical examination fees in health centers. In contrast with results of Tshomba et al., [15] and Almeida et al., [19], this study confirms that the level of education were not directly associated with the practice self-medication.

Participants' age was not associated with antimalarial self-medication according to studies conducted by Tshomba et al. Congo [15], Yameogo et al. Burkina [20] and Deressa et al. Ethiopia [21]. However some authors had identified age as a factor associated with antimalarial self-medication. In Ghana, authors reported that older than five subjects were more likely to experience antimalarial self-medicate than those less than five years [17]. In Senegal, in a study conducted in hospitals, authors observed that self-medication for malaria fever was significantly more frequent among patients under fifteen. [22] Among student in Nigeria, authors found that the prevalence of self-medication was higher in the age group 25 to 44 years and lower in age from 15 to 24 years. [4]

The present study found that both men and women equally practiced antimalarial self-medication but studies conducted in Saudi Arabia and Tanzania reported that men were most concerned [23-25] by this practice. Variations in gender considerations according to different societies may explain divergences in results.

5.2.2. Behavioral factors

The preference for drug consumption without health workers' advices and the non-utilization, health care facilities were not associated with antimalarial self-medication. This finding may be explained by the fact that these behaviors are partly related to the economic level with was associated with the practice of self-medication. The effect of these behavioral factors would therefore be mediated by the economic level. The results of this study are not in contradiction with those observed in Tanzania Chipwaza et al. [24] Kenya by Watsierah et al. [26] and Afolabi Nigeria [27] who reported that some behaviour such as the preference for drug non-recommended by health workers, the non-utilization health care facilities were associated with antimalarial self-medication.

5.2.3. Factors related to patient knowledge

A significant relationship was observed between antimalarial self-medication and ignorance of its dangers, knowledge of the consequences of poorly treated malaria and dangers associated with street drugs consumption.

Dangers of antimalarial self-medication with street drugs are relative to the inadequate dosages of active chemicals [28]. Inadequate contents in active chemicals contribute to emergence of drug resistance, but overdoses can also be dangerous for health. This shows the importance for the population to be informed not only on the dangers of antimalarial self-medication but also the consequences of poorly treated malaria. This study showed that the participants were sensitized on the poor quality of street drugs. Nearly 32% of respondents declared that drugs sold in illicit market had no consequence on health. The poor storage of street antimalarial drugs could enhance their toxicity and the suboptimal content in active chemicals was factors could contribute to drug-resistance of Plasmodium falciparum [29].

5.2.4. Environmental factors

Exposure to advertising on traditional medicine products used as malaria and living close to illicit drug market were not associated with antimalarial self-medication. These results are not consistent with other study. In Lomé in Togo, D'almeida observed that drug advertising and living near illicit drug market were factors promoting self-medication [18]. Study participants declared that vehicles were used to advertise supposed effective products against malaria. This encouraged antimalarial self-medication

5.2.5. Factors related to the health system

The distance to health facilities was not significantly associated with antimalarial self-medication. This result is not consistent with those observed in a study conducted in the Middle East where the authors had shown that geographic access to health centers contributed to self-medication practices [25]. In Comé, dwellers use motorcycle for transportation for low fee, For this, it is possible that the distance to the health center does not influence the behavior of antimalarial self-medication practices.

Furthermore, non-satisfaction with the care provided in health centers was not associated with the practice of self-medication in contrast to results of Alghanim et al., who reported that health care quality strongly influenced self-medication [25].

The limitations of the present study are relative to the fact that the results can be only extrapolated to the city and its rural outskirts but not to all the South Benin. However, the results reported in this study are very useful for the design and implementation of interventions against the antimalarial self-medication

6. CONCLUSION

Antimalarial self-medication was common in households in Comé. Interventions to increase knowledge on consequences of antimalarial self-medication, dangers of poorly treated malaria and adverse effects of street drugs consumption are timely for rational use of antimalarial drugs and prevention of drug resistance in household in Comé, southern Benin.

REFERENCES

1. World Health Organisation. Division of control of tropical diseases. Several and complicated malaria. 2nd ed. Trans R Soc Trop Med Hyg 1990 ; 84 : 21-65.
2. Institut Pasteur. Paludisme. Aide-mémoire, [en ligne], Mars 2013 [Consulté le 15/02/2015]: «<http://www.pasteur.fr/fr/institut-pasteur/presse/fiches-info/paludisme>»
3. Olliaro PL, Taylor RWJ. Developing artemisin based drug combinations for the treatment of resistant falciparum malaria: A review. J Postgrad Med 2004; 50:40-4.
4. Osemene KP, Lamikanra A. Study of the prevalence of self-medication practice among university students in southwestern Nigeria. Tropical Journal of Pharmaceutical Research 2012; 11 (4): 683-689
5. Chiribagula, VB. Mboni H.M, Amuri S.B, kamulete G.S, Byanga J.K., Duez P, Simbi JB. L. Prévalence et caractéristiques de l'automédication chez les étudiants de 18 à 35 ans résidant au Campus de la Kasapa de l'Université de Lubumbashi. Pan African Medical Journal. 2015; 21:107 doi:10.11604/pamj.2015.21.107.5651
6. Akangan AS. L'automédication en milieu urbain béninois à Cotonou, à propos d'une enquête auprès de 275 ménages. Thèse de doctorat n° 0265.1986 ; Faculté de Médecine, Université Nationale du Bénin.
7. D'Almeida AGAA. Problématique de l'automédication dans la commune urbaine de Lomé (TOGO). Thèse de doctorat n° 52. Faculté de médecine, de pharmacie et d'odontologie ; Université Cheikh Anta Diop de Dakar ; 2003 :59
8. Donkor S, Tetteh-Quarcoop P B, Nartey P, Agyeman OI. Self-Medication Practices with Antibiotics among Tertiary Level Students in Accra, Ghana: a cross-sectional study. Int J Environ Res Public Health. 2012; 9(10): 3519-3529
9. WHO. Guidelines for the regulatory assessment of medicinal products for use in self-medication. WHO/EDM/QSM/002000.Disponible sur <http://apps.who.int/medicinedocs/pdf/s2218e/s2218e.pdf>, consulté le 29 /04/2016
10. Sanfo L. L'automédication dans la ville de Ouagadougou : une enquête réalisée auprès des officines pharmaceutiques : Thèse de doctorat n 25, 1999. Faculté des sciences de la santé, Université de Ouagadougou, Ouagadougou.77p.
11. Konate L. Etude de l'automédication dans les officines de la ville de Sikasso. Thèse de doctorat, 2005. Faculté de Médecine, de Pharmacieet d'Odontostomatologie,Université de Bamako. 78 p.

12. World Health Organization. Antimalarial Drug Combination Therapy: Report of a WHO Technical Consultation. Geneva, Switzerland: 2001.
13. Ahounou D. Prise en charge à domicile du paludisme simple chez les enfants d'âge inférieur à 5 ans en milieu côtier lagunaire du Bénin. Les cas de Ladji et Toweta 2 (Cotonou) – résultats préliminaires. Bull Soc Pathol Exot, 2006, 99, 5, 301-340
14. Ministère de la Santé. Plan stratégique national de lutte contre le paludisme 2011-2015, Cotonou, 2011
15. Tshomba OA, Atoba BC, Batina AS, Kazadi MD, Kayembe TC, Lutumba P. Determinants socio-economiques de l'automédication aux antimalariens à Kisangani, RD Congo. Rev. méd. Gd. Lacs 2013, 2(1): 37-51
16. Ouédraogo LT, Somé IT, Diarra M, Guissou I.P. Automédication dans le traitement de l'accès palustre : étude auprès de clients d'officines pharmaceutiques privées de la ville de Ouagadougou, Burkina Faso. Bull Soc Pathol Exot, 2008, 101, 2, 124-127
17. Aborah S, Akweongo P, Adjuik M, Atinga R, Welaga P, Adongo P. The use of non-prescribed anti-malarial drugs for the treatment of malaria in the Bolgatanga municipality, Ghana. Malaria Journal 2013, 12:266
18. D'almeida A. Problématique de l'automédication dans la commune urbaine de Lome ; Thèse de doctorat, Université du Bénin, Togo 2003.
19. Dargicho A, Nasir Tajure W, Mulugeta T. Self-Medication with Antibiotics and Antimalarials in the Community of Silte Zone, South Ethiopia. Prev Med Bull 2012; 11(5): 529-536
20. Yaméogo T M, et al. Chemin de soin des cas présomptifs de paludisme admis dans un hôpital de district au Burkina Faso. Med et Santé Trop 2014 ; 24 : 301-306
21. Deressa W, Ali A, Enqusellassie F. Self-treatment of malaria in rural communities, Butajira, southern Ethiopia. Bull of the World Health Organization 2003, 81 (4)
22. Ndiaye P, Tal-dia A, Diedhiou A, Juergens-Behr A, Lemort JP. L'automédication de la fièvre dans le district nord du Sénégal. Med Trop 2006; 66 : 74-78
23. Kigodi NK, Magreth K. Malaria and anti-malarial drugs utilisation among adults in a rural coastal community of Tanzania: knowledge, attitude and practice study, Dar es Salaam. Medical Students Journal 2006 ; 14 (7) : 1-10
24. Chipwaza B, Mugasa JP, Mayumana I, Amuri M, Makungu C, Gwakisa PS. Self-medication with anti-malarials is a common practice in rural communities of Kilosa district in Tanzania despite the reported decline of malaria, Arusha. Malaria Journal 2014, 13:252
25. Alghanim S A. Self-medication practice among patients in a public health care system, Arabie Saoudite. Eastern Mediterranean Health Journal 2011, 17 (5)
26. Watsierah C, Jura W, Oyugi H, Abong B, Ouma C. Factors determining anti-malarial drug use in a peri-urban population from malaria holoendemic region of western kenya. Malaria Journal 2010, 9:295
27. Afolabi A O. Factors Influencing the Pattern of Self-Medication in an Adult Nigerian Population, Nigeria. Annals of African Med 2008 ; 7(3) : 120-127
28. Hamani Abdou I. Les médicaments de la rue à Niamey: Modalités de vente et contrôle de qualité de quelques médicaments anti-infectieux. Thèse de doctorat en pharmacie, Université de Bamako, 2005. 140p
29. Baba-Moussa F., Bonou J., Ahouandjinou H., Tamègnon V., Dougnon V., Kpavode L., Raimi N. et al. Control of Selected Antimalarials Sold in the Illicit Market: An Investigation Conducted

391 in Porto-Novo City (Republic of Benin). Advances in Bioscience and Biotechnology, 2015, 6,
392 637-644