

Parasitological Assessment and Public Health Implications of Water used for Domestic Purposes in Ebonyi Local Government Area of Ebonyi State, Nigeria.

ABSTRACT

Aims: Water borne diseases have been a major cause of morbidity in most developing countries including Nigeria maybe due to poor quality of water used for domestic and industrial purposes. This study was on the parasitological assessment and public health implications of water used for domestic purposes in Ebonyi Local Government Area, Ebonyi State, Nigeria.

Study design: This was an observational study.

Place and Duration of Study: Ebonyi Local Government Area of Ebonyi State, Nigeria between September, 2014 and October, 2014.

Methodology: Samples from different water bodies in each of four communities in Ebonyi Local Government Area were collected and analyzed using standard parasitological methods.

Results: The results revealed the presence of seven (7) different parasites including *Giardia lamblia*, *Schistosoma haematobium*, *Ascaris lumbricoides*, *Entamoeba histolytica*, *Strongyloides stercoralis*, *Diphylobotrium latum* and *Taenia spp.* The result showed that *Giardia lamblia* has the highest occurrence of 23.57% while *Entamoeba histolytica* recorded the least parasite present with 7.14%. The pond water showed the highest occurrence of parasites present with 44.29% while the least occurrence was borehole with 0.71%. Only borehole from Nkaleke community showed the presence of a parasite (*Entamoeba histolytica*).

Conclusion: This study result is of alarming concern as the water bodies sampled are the major water sources available in those areas and by this study are unsafe for domestic use because of their high parasite presence. It is therefore imperative to suggest that the government of Ebonyi state and Nigeria at large should make available infrastructures that would make the water bodies safe for domestic use. Meanwhile, the inhabitants of at risk areas and the general populace are advised to boil or treat their drinking water with water guard before use. Also, indiscriminate dumping of refuse or faecal materials should become everyone's concern and greatly discouraged.

Keywords: Water, *Giardia lamblia*, *Schistosoma haematobium*, *Ascaris lumbricoides*, *Entamoeba histolytica*, *Strongyloides stercoralis*, *Diphylobotrium latum*, *Taenia spp.*, Faecal contamination

1. INTRODUCTION

The survival of man in any given geographical area depends on the availability and quality of water in the area. Indeed, virtually all human activities involve water usage ranging from drinking, washing, cooking, irrigation and for industrial activities [1]. However, the human race is under tremendous threat due to undesired changes in the physical, chemical and biological characteristics of air, water and soil [2]. Due to increased human population, industrialization, use of fertilizers and man-made activities, water is highly polluted with different harmful contaminants [3]. Although the water content of the biosphere is enormous, only a small portion is available for human domestic use due to contamination [3]. Faecal matter from patients and healthy carriers of pathogenic microorganism poses the most threatening form of water pollution [4]. The most predominant waterborne disease, diarrhea, has an estimated annual incidence of 4.6 billion episodes and causes 2.2 million deaths every year [5]. Children are the main victims of diarrhea and other faecal-oral disease, and also the most likely source of infection [6]. Since parasites live in the intestine of an infected host and the cyst of the parasite can be passed out as part of the faecal matters of infected person or animal, indiscriminate defecation is seen as the culprit to the high prevalence of parasites in water bodies, which consequently cause waterborne diseases [7]. Biological contaminants such as bacteria,

viruses, fungi, protozoa and helminth constitute the major cause of foodborne and water-borne diseases with varying degrees of severity, ranging from mild indisposition to chronic or life-threatening illness, or both ([7]. Water borne diseases of parasitic origin are of major concern to public health globally. Majorly in developing countries, water borne diseases have been frequently associated with high mortality and morbidity especially in children [8]. The WHO has estimated that food- and water-borne infectious diseases currently infect 3.5 billion people in developing countries and cause about 160,000 deaths per year and 80% of these occur in children less than 5 years of age [9]. This study was conducted to estimate the parasitological quality of domestic water sources in Ebonyi Local Government Area of Ebonyi State, Nigeria.

2. MATERIALS AND METHODS

2.1 Study Area

This study was carried out in Ebonyi Local Government Area, one of the 13 Local Government Areas in Ebonyi State, Nigeria. Ebonyi Local Government Area is an agrarian council area. It is located between longitude $7^{\circ}59'E$ and $8^{\circ}20'E$ and latitude $6^{\circ}22'N$ and $6^{\circ}43'N$. The area falls within the climatic region of South Eastern Nigeria where the rainy season spans from April to October and dry season from October to April [10]. The average annual rainfall of the study area is about 1500mm with actual surface temperature (seasonal temperature) of between $24-36^{\circ}C$ during dry season and about $18^{\circ}C$ during the rainy season [10]. In the local government area, four randomly selected communities, namely: Ndiabor, Mbeke, Ndiagu and Nkaleke were used for the study.

2.2 Sample Size and Source

Samples were collected from 320 sample points (10 points for each water body in all communities) in two containers in each point. Well, stream, borehole and pond water bodies were used for the study. Samples were collected separately in clean sterilized plastic bottles in the months of September and October, 2014 from Ndiabor, Mbeke, Ndiagu and Nkaleke communities of Ebonyi Local Government Area of Ebonyi State, Nigeria.

2.3 Sample Collection

Samples were collected in one liter containers already pre-sterilized at $121^{\circ}C$ for 15 minutes and preserved with 10% formalin as described by Roohul-Amin *et al.* [11]. While sampling in the streams, the collection bottle was lowered in water at a depth of about 15cm to 30cm. The bottle was held at the base and placed against the direction of the water flow. Immediately the water was collected, the bottle was covered up to avoid, carefully labeled and transported to the Department of Biological Science Laboratory, Ebonyi State University, Abakaliki for immediate processing.

2.4 Parasitological Analysis of Water Samples

The preserved one liter sample was allowed to settle for 24 hours. The supernatant was carefully removed and discarded without disturbing the sediment using a siphon. The sediment was then transferred to centrifuge tubes. The walls of the sedimentation container was washed thoroughly using a spray bottle with detergent solution and the rinsing was added to the sediments in the centrifuged tubes. All the recovered materials were centrifuged at 1000g for 15 minutes. The supernatant was removed and discarded and the sediments transferred to one tube. Flotation technique which is ideal for the recovery of parasite eggs/larva/cyst was used by suspending the pellet in an equal volume of aceto-acetic buffer (pH 4.5) [12,13]. The mixture was re-centrifuged at 1000g for 15 minutes and the pellets were transferred to microscope counting cell for final examination using objectives 10x/ 40x magnification. The parasites were identified with a practical guide of Ukaga *et al.* [14]. The number of parasites per liter of water was calculated following the equation as used by Kpoda *et al.* [15].

$$N = AX/PV$$

Where:

N = no of eggs/cyst/larva/ova of parasite

A = no of parasites counted on the Mac master slide

X = volume of the final product (ml)

82 P = Mac master cell capacity (0.3ml)
83 V = Pellet volume

84 2.5 Statistical Analysis

85 Data were analyzed using Statistical Package for Social Sciences (SPSS) version 19.0.

86

87 3. RESULTS

88 The parasitological examination of domestic water samples from four communities in Ebonyi Local
89 Government Area of Ebonyi State, Nigeria revealed the presence of seven (7) different parasites
90 including: *Giardia lamblia*, *Schistosoma haematobium*, *Ascaris lumbricoides*, *Entamoeba histolytica*,
91 *Strongyloides stercoralis*, *Diphylobotrium latum* and *Taenia spp.* (Plates 1-6). In this study, *Giardia*
92 *lamblia* had the highest prevalence rate of 8(25.81%), followed by *Schistosoma haematobium* and
93 *Taenia spp.*, both having the prevalence rate of 6(19.35%); however *Diphylobotrium latum* was not
94 seen in the water samples from Ndiabor. Out of the 31 parasites seen in water samples from Ndiabor,
95 8(25.80%) was from well, 9(29.03%) from stream and 14(45.16%) was from pond (NP), whereas no
96 parasite was seen in borehole (Table 1).

97 The results of the analysis of water samples from Mbeke community show that both *Giardia lamblia*
98 and *Taenia spp.* had the highest prevalence rate of 10(26.32%), followed by *Ascaris lumbricoides*
99 having 8(21.05%) prevalence rate; *Entamoeba histolytica* was observed least, having the occurrence
100 rate of 1(2.263%). Out of the 38 parasites seen in water samples from Mbeke community, 20(52.63%)
101 was from pond, 11(28.95%) well and 7(18.42%) from stream, whereas no parasite was seen in
102 borehole water sampled (Table 2). The analysis of water samples from Ndiagu community of Ebonyi
103 Local Government Area showed that *Giardia lamblia* and *Taenia spp.* occurred highest at the rate of
104 8(25.00%) each, followed by *Schistosoma haematobium* having 6(18.75), while *Entamoeba histolytica*
105 and *Strongyloides stercoralis* had the lowest occurrence of 2(6.25%) each. Out of the 32 parasites
106 seen in water samples from Ndiagu community, 13(40.62%) was from well, 12(37.50%) was from
107 pond and 7(21.88%) was from stream; no parasite was seen in borehole water samples (Table 3).

108 In Nkaleke community, the results showed that *Ascaris lumbricoides* had the highest occurrence of
109 8(20.00%) followed by *Giardia lamblia* and *Taenia spp.* having the occurrence of 7(17.50%) each;
110 *Strongyloides stercoralis* had the least occurrence of 2(5.00%). Out of the 40 parasites seen in
111 samples from Nkaleke community, 16(40.00%) was from pond, 13(32.50) from stream and
112 10(25.00%) was from well; only 1(2.50%) was from the borehole water samples in Nkaleke
113 community. Table 5 summarizes the mean percentages by both parasites and water sources. Pond
114 water recorded the highest (44.3%) parasites presence while borehole water recorded the least
115 percentage (0.71%) parasite presence. Meanwhile, *Giardia lamblia* was leading with 23.6% presence
116 while *Entamoeba histolytica* had the least percentage presence of 7.14% presence.

117

118



Plate 1: Cyst of *Giardia lamblia*




Plate 2: Egg of *Schistosoma haematobium*




Plate 3: Egg of *Ascaris lumbricoides*




Plate 4: Cyst of *Entamoeba histolytica*



Plate 5: Larvae of *Strongyloides stercoralis*




Plate 6: Egg of *Diphylobotrium latum*

119 120 Plates 1-6: Eggs/cysts/larvae of parasites observed in the study

121 **Table 1. Parasitological Examination of water Samples from Ndiabor**

S/N	Parasite	NW*	NS*	NB*	NP*	Total
1	<i>Giardia lamblia</i>	3	1	0	4	8
2	<i>Schistosoma haematobium</i>	2	4	0	0	6
3	<i>Ascaris lumbricoides</i>	2	0	0	3	5
4	<i>Entamoeba histolytica</i>	0	0	0	1	1
5	<i>Strongyloides stercoralis</i>	0	2	0	3	5
6	<i>Diphylobotrium latum</i>	0	0	0	0	0
7	<i>Taenia spp</i>	1	2	0	3	6
Total		8	9	0	14	31

122 NW = Ndiabor well, NS = Ndiabor stream, NB= Ndiabor bore hole

123 NP = Ndiabor pond, * = Number of egg/cyst/larva

124

125 **Table 2: Parasitological Examination of Water Samples from Mbeke Community**

S/N	Parasite	MW*	MS*	MW*	MP*	Total
1	<i>Giardia lamblia</i>	5	2	0	3	10
2	<i>Schistosoma haematobium</i>	2	1	0	2	5
3	<i>Ascaris lumbricoides</i>	1	1	0	6	8
4	<i>Entamoeba histolytica</i>	0	0	0	1	1
5	<i>Strongyloides stercoralis</i>	0	0	0	2	2
6	<i>Diphylobotrium latum</i>	0	0	0	2	2
7	<i>Taenia spp</i>	3	3	0	4	10
Total		11	7	0	20	38

126 MW = Mbeke well, MS = Mbeke stream, MB = Mbeke bore hole

127 MP = Mbeke pond, * = Number of egg/cyst/larva

128

129 **Table 3: Parasitological Examination of Water Samples from Ndiagu Community**

s/n	Parasites	NDW*	NDS*	NDB*	NDP*	Total
1	<i>Giardia lamblia</i>	3	1	0	4	8
2	<i>Schistosoma haematobium</i>	2	2	0	2	6
3	<i>Ascaris lumbricoides</i>	1	1	0	1	3
4	<i>Entamoeba histolytica</i>	1	0	0	1	2
5	<i>Strongyloides stercoralis</i>	2	0	0	0	2
6	<i>Diphylobotrium latum</i>	1	2	0	0	3
7	<i>Taenia spp</i>	3	1	0	4	8
Total		13	7	0	12	32

130 NDW = Ndiagu well, NDS = Ndiagu stream, NDB = Ndiagu bore hole

131 NDP = Ndiagu pond, * = Number of egg/cyst/larva

132

133 **Table 4: Parasitological Examination of Water Samples from Nkaleke Community**

s/n	Parasites	NKW	NKS	NKB	NKP	Total
1	<i>Giardia lamblia</i>	3	0	0	4	7
2	<i>Schistosoma haematobium</i>	2	2	0	0	4
3	<i>Ascaris lumbricoides</i>	1	6	0	1	8
4	<i>Entamoeba histolytica</i>	1	2	1	2	6
5	<i>Strongyloides stercoralis</i>	1	1	0	0	2
6	<i>Diphylobotrium latum</i>	0	1	0	5	6
7	<i>Taenia spp</i>	2	1	0	4	7
Total		10	13	1	16	40

134 NKW = Nkaleke well, NKS = Nkaleke stream, NKB = Nkaleke bore hole

135 NKP = Nkaleke pond, * = Number of egg/cyst/larva

136

137

138

139

140

141 **Table 5: Percentage Parasites Presence in all Sampled Domestic Water**

S/N	Parasites	Well	Stream	Borehole	Pond	Total (%)
1	<i>G. lamblia</i>	14	4	0	15	23.6
2	<i>S. haematobium</i>	8	9	0	4	15.0
3	<i>A.. lumbricoides</i>	5	8	0	11	17.1
4	<i>E. histolytica</i>	2	2	1	5	7.1
5	<i>S. stercoralis</i>	3	3	0	5	7.9
6	<i>D. latum</i>	2	3	0	7	8.6
7	<i>Taenia spp</i>	7	7	0	15	20.7
Total (%)		29.3	25.7	0.7	44.3	100

142 143 144 **4. DISCUSSION** 145

146 The result of the parasitological survey of the four domestic water sources from Ebonyi Local
147 Government Area showed that the water samples contained varying degrees of presence of seven (7)
148 intestinal parasites including *Giardia lamblia*, *Ascaris lumbricoides*, *Taenia spp*, *Strongyloides*
149 *stercoralis*, *Entamoeba histolytica*, *Schistosoma haematobium* and *Diphylobotrium latum*. The most
150 occurring parasite observed in this study was *Giardia lamblia* (23.6%), followed by *Taenia spp*.
151 (20.7%). The high prevalence might be due to high contamination of water bodies with faecal
152 materials as inhabitants of the study area indiscriminately defecate in their environment. This result
153 corroborate with other previous studies of Stott [16], Coelho *et al.* [17], Asaolu *et al.* [18], Valbuena
154 [19] and Iyayi *et al.* [20] who showed the frequent presence and high concentration of *Giardia lamblia*
155 and *Ascaris lumbricoides* in water and raw sewage. Similarly, Kpoda *et al.* [15] assessed the quality
156 of vegetable irrigation water in Ougadougou city, Burkina-Faso Parasitologically. The results showed
157 high presence of *Ancylostoma duodenalis*, *Giardia lamblia*, *Hymenolepis nana*, *Taenia spp*, *Acaris*
158 *lumbricoides*, *Strongyloides Stercoralis*, *Entamoeba histolytica*. However, this present study did
159 record any occurrences of *Ancylostoma duodenalis* and *Hymenolepis nana*.

160 Among the water sources, pond water recorded the highest prevalence of parasite (44.29%). This
161 could be attributed to the fact that during the rains, ponds serves as reservoirs that collect run-off
162 water from different routes. This conforms to previous studies of Chollom *et al.* [9], and Ani and Itiba
163 [21]. The well water also recorded a high prevalence of parasites (29.29%), which agrees with CDC
164 [22], that wells may be more vulnerable to contamination after flooding particularly if the wells are
165 shallow, have been dug or bored or have been submerged by flooding water for long period of time;
166 this may also be due to lack of toilet facilities in these communities and the fact that wells in these
167 communities are always open.

168 Water samples from borehole had low rate of parasite infestation (0.7%); this could be explained by
169 the fact that boreholes operate a closed system and this restricts faecal contaminations and contact
170 with parasite vectors. Chollom *et al.* [9] reported that infestation of water is contaminative in nature as
171 human parasites do not directly use water bodies for life cycle development; rather their vectors
172 inhabit water bodies thereby associating their transmission to water. However, the *E. histolytica*
173 observed in a sample of borehole from Nkaleke community is in disparity with the report of Chollom *et*
174 *al.* [9] that water samples from boreholes were found to be free from parasites. This variation can be
175 as a result of closeness to septic tanks, soakaway pit, shallowness of the borehole, as well as
176 possible contamination during sample collection.

177 178 **5. CONCLUSION**

179 Since this study showed less than one parasite per liter of water in three of the four domestic water
180 samples examined, it however means that with the exception of borehole water samples, the rest of
181 the water sources are not good for drinking and other domestic purposes. This finding is in
182 accordance with the earlier work of Nwele *et al.* [12] and Hajjami *et al.* [23] which recorded mean
183 concentration of 2.4 eggs/l and 8.98 eggs/l respectively. The cause of this is relatively connected to
184 poor hygiene of individuals, low sanitary facilities and different human activities in and around the
185 water bodies [12]. This study result is of alarming concern as the water bodies sampled are the major
186 water sources available in those areas and by this study are unsafe for domestic use because of their
187 high parasite presence. It is therefore imperative to suggest that the government of Ebonyi state and

Nigeria at large should make available infrastructures that would make the water bodies safe for domestic use. Meanwhile, the inhabitants of **at** risk areas and the general populace are advised to boil or treat their drinking water with water guard before use. Also, indiscriminate dumping of refuse or faecal materials should become everyone's concern and greatly discouraged.

REFERENCES

- [1] Oko OJ, Aremu MO, Odoh R, Yebpella G, Shenge GA. Assessment of Water Quality Index of Borehole and Well Water in Wukari Town, Taraba State, Nigeria. *Journal of Environment and Earth Science*. 2014;4(5):1-9.
- [2] Patil P N, Sawant DV, Deshmukh RN. Physicochemical parameters for testing of water – A review. *International Journal of Environmental Sciences*. 2013;3(3): 1194-1207.
- [3] Agah MV, Onochie CC, Nworie O, Oji AE, Nnachi AU, Chukwu OS, Udu-Ibiam OE, Ukaegbu CO, Ani C. Physicochemical and Bacteriological impact of Effluents from Ebonyi Fertilizer and Chemical Plant on Azuiyokwu Creek in Abakaliki, Ebonyi State, Nigeria. *Journal of Applied and Environmental Microbiology*. 2014; 2(4): 135-142.
- [4] Oyedele O, Olutiola PO, Mononuola MA. Microbiological quality of packaged drinking water brands marketed in Ibadan metropolis and Ile-Ile city in South Western Nigeria. *Journal of Microbiology Research*. 2010 4(1): 96 - 102.
- [5] WHO. Water for health, WHO Guidelines for Drinking-water Quality. WHO Geneva, Switzerland; 2010.
- [6] Tsega N, Sahile S, Kibret M, Abera B. Bacteriological and physico- chemical quality of drinking water sources in a rural community of Ethiopia. *African Health Sciences*. 2013; 13(4): 1156- 1161.
- [7] Okonko IO, Ogun AA, Adejoye OD, Ogunjobi AA, Nkang AO, Adebayo-Tayo BC. Hazards analysis critical control points (HACCP) and Microbiology qualities of Sea-foods as affected by Handler's Hygiene in Ibadan and Lagos, Nigeria. *African Journal of Food Sciences*. 2009; 3(1):035-050.
- [8] O'Ryan M, Prado V, Pickering LK. A millennium update on pediatric diarrheal illness in the developing world. *Seminars in Pediatric Infectious Diseases*. 2005;16(2):125-136.
- [9] Chollom, SC, Iduh MU, Gyang BJ, Idoko MA, Ujah A, Agada GO, Peter J, Akele YR, Okwori JA. Parasitological Evaluation of Domestic Water Sources in a Rural Community in Nigeria. *British Microbiology Research Journal*. 2013;3(3): 393-399.
- [10] Ezeh H, Ugwu S. Geo-electric investigation of sulphide mineralization at Egu-Echaraunuhu, Abakaliki, South-East, Nigeria. *Journal of Applied Natural Sciences*. 2007; 1(1):59-63.
- [11] Roohul-Anim SSA, Zubair A, Jabar ZKK. Microbial analysis of drinking water and water distribution system in new urban Peshawar. *Current Research Journal of Biological Science*. 2012; 4(6):731-737.
- [12] Nwele DE, Uhuo AC, Okonkwo EC, Ibiam GA, Onwe CS, Ugwu JI, Onwe JC, Nwuzo AC. Parasitological examination of Ava stream used in irrigation in Enugu state, south-eastern Nigeria: An implication for helminth transmission. *Journal of Parasitology and Vector Biology*. 2013; 5(8): 112-115.
- [13] Grantzer C, Gaspard P, Galnez L, Huyard A, Dumouthier N, Schwartzbrod J. Monitoring of bacterial and parasitological contamination during various treatment of sludge. *Water Research*. 2001;35(16):3763-3770.
- [14] Ukaga CN, Onyeka PIK, Nwoke BEB. *Practical Medical Parasitology*. Aran Global Publication, Nigeria; 2002.
- [15] Kpoda NW, Oueda A, Some YSC, Cisse G, Maiga AH, Kabre GB. Physicochemical and parasitological quality of vegetables irrigation water in Ouagadougou city, Burkina-Faso. *Africa Journal of Microbiology Research*. 2015;9(5):307-317.
- [16] Stott R. A survey of the microbial quality of waste waters in Ismailia, Egypt and the implications of waste water reuse. *Water science Technology*. 1997;35(11-12): 211-217.
- [17] Coelho LM, Oliveira SM, Milman MH, Karasawa KA, Santos RD. Detection of transmissible forms of ectoparasites in water and vegetables consumed at school in Sorocaba, Sao Paulo state, Brazil. *Revised Brazilian Medical Journal*. 2001;34: 479-48.
- [18] Asaolu SO, Ofoezie IE, Odumuyiwa PA, Sowemimo OA, Ogunniyi, TAB. Effect of water Supply and Sanitation on the prevalence and intensity of *Ascaris lumbricoides* among pre-school-age children in Ajebandele and Ifewara, Osun State, Nigeria. *Transaction of the Royal Society of Tropical Medicine and Hygiene*. 2002;96(6):600-604.

- 247 [19]Valbuena D. Detention of intestinal helminthes and contamination- indicator bacteria in
 248 treated and untreated waste water. Integrated Science. 2002; 27(12): 710-714.
- 249 [20]Oyibo IF, Ahmadu L, Stephen OO, Ameh YC. Evaluation of parasites of medical importance
 250 in drinking water sources in Okura district, Dekina Local Government, Kogi State, Nigeria.
 251 International Journal of Development Research. 2015; 6(4):7290-7294.
- 252 [21]Ani OC, Itiba OL. Evaluation of the Parasitic Contamination of Local Sources of Drinking
 253 Water in Abakaliki, Ebonyi State, Nigeria. IOSR Journal of Dental and Medical Sciences
 254 (IOSR-JDMS). 2015;14(9 Ver. I):30-34.
- 255 [22]CDC. Diseases and contaminants. Centers for Disease Control and Prevention; 2015.
- 256 [23]Hajjami K, Ennaji MM, Fouad S, Qubrim N, Cohen N. Waste water reuse for irrigation in
 257 Morocco. Journal of Parasitology. 2013;4:163-162.