

Original Research Article

Superficial Fungal Skin Infections in Patients Attending Zliten Teaching Hospital(North west of Libya)

Abstract:

Superficial fungal infection(SFI) of skin and it's appendages is common health problem worldwide. Dermatophytosis and Pityriasis versicolor are the main SFI in humans. The risk factors are hot humid environment leading to excessive sweating, and contact with animals. over a period of 9 Months, 3303 patients of various skin complaints were examined at outpatient Department of Zliten Teaching Hospital.

A Retrospective study was carried out to analyze the data to find out the distribution pattern of different clinical types of SFI in relation to age and sex. 161(4.9%) patient were found to be affected with SFI. 22.9% of infected patients were less than 7 years old, followed by age group from 21-27 years(14.9%). Male to female ratio was 1.2:1. The most frequent clinical pattern were, Tinea capitis (25.5%), followed by Tinea pedis (22.4%), Tinea corporis (21.7%) and Pityriasis versicolor(18.6%). Tinea capitis where the most frequent superficial fungal infection before 14 years of age, and Pityriasis versicolor was the most frequent fungal infection in the age group 13-20 years.

Key words: Superficial fungal infection, Dermatophytosis, Pityriasis versicolor, tinea capitis.

1- Introduction:

The Fungi are living organisms present in our environment and there are fungi considered to be normal flora such as *Malassezia* species. However, the organisms affect humans and cause disease.¹ Furthermore, the non-dermatophyte molds could be a cause of fungal infection of nails (Onychomycosis).² Fungal infections of skin, nails and hairs are one of most frequent diseases and in the last decades, the prevalence of fungal infections of skin and appendages has increased to affect around 25% of population of different world societies.³ Marques *et al* (2000) found that Dermatophyte infections of skin affect one fifth of world's population⁴.

The SFI are mainly confined to stratum corneum(upper layer of Epidermis) and cornified appendages (hairs and nails).² Dermatophytosis, is a group of important superficial fungal

infections of skin and it's appendages and are caused by three etiologic species of dermatophytes, which are microsporum, trichophyton and epidermophyton, and each type of these three is characterized by specific morphologic features of cultured colonies and of Macroconidia under Microscope. Dermatophyte infections of skin also known as ringworm infections and named after affected part of the body, e.g. Tinea capitis and Tinea pedis.⁵ Symptoms and signs of Dermatophyte infections of skin and it's appendages differ according to the affected site of the body and the source of fungi. Generally, skin lesions are scaly well defined erythematous plaques with active margin showing more scales with or without vesicles and pustules. The lesions tend to extend peripherally and clear centrally. Zoophilic fungi exhibit more inflammation compared to anthropophilic fungi.² Another common SFI is Pityriasis versicolor which is mainly caused by Malassezia species which is a yeast, and manifested as round or oval patches in different colours (white, red and brown) with fine scales.^{2,6}

The risk factors for SFI are hot, humid climate, which encourages sweating and appearance of fungal infections of skin; furthermore contact with infected animals, like cats could lead to skin infection of humans with zoophilic fungi.⁶

In Libya, there are few studies were conducted about SFI, furthermore, no previous studies about this issue were performed in Zliten. To know about the Epidemiology of dermatophytosis and Pityriasis versicolor in this area we have conducted this study.

2- Methods and subjects:

2.1. Area of Study:

Zliten is a Libyan city with more than 263,000 people and located on Mediterranean Sea about 150 kilometers east of Tripoli, with hot weather and high humidity in summer months.⁷

2.2 Subjectives:

This study carried out to discover the Epidemiology of dermatophytosis and Pityriasis versicolor in Zliten (North west of Libya).

2.3. Methods:

The data for all patients were used for retrospective study, the patients who have seek dermatological advice at outpatient Department of Zliten teaching Hospital. The study was

60 carried out in the period of time from 01.10.2014 until 30.06.2015. All patients were from
61 area of study. Patients were examined with wood's lamp, scraping of skin, nails and hairs
62 were examined under microscope with use of KOH 20% solution. The culture of scrapings
63 were used to reach definitive diagnosis. Scrapings were taken with sterile scalpel into a glass
64 slide and 20% potassium hydroxide (KOH) as aqueous solution were applied and the slide
65 was covered with cover slide and the scraping examined under microscope. For culture
66 purposes, the scraped site was cleaned aseptically with 70% ethanol and specimens was
67 directly collected into petri dishes. The culture was performed on sabroud dextrose agar
68 media with typical formula containing mycological peptone 10.0: glucose 40.0: Agar 15.0;
69 petri dishes were incubated at 26C° and the growth of cultures was observed twice weekly
70 and petri dishes were discarded only after 3 weeks in the absence of growth.

71 2.4. Statistical analysis:

72 The data were analysed with use of SPSS 20 to study the distribution and frequency of
73 superficial fungal infections of skin, hairs and nails.

74

75 **Results:**

76 A total number of 3303 Patients sought medical advice because of skin problems at
77 outpatient Department of Zliten Teaching Hospital. 161/3303(4.9%) were found to have SFI,
78 88/161 (54.7%) were males and 73/161(45.3%) were females. The ratio of male to female
79 was 1.2:1. The age of patients ranged from 1 to 80 years (mean, 25,3 years). Of 161 patients,
80 37(22.9%) of patients were located in the range between 0-6 years old, followed by age group
81 from 21-27 years (14.9%). We observed that 36% of patients with SFI were less than 14
82 years old(childhood age group). 34.2% were found to be in age groups from 14 to 34 years
83 which represent adolescence and adulthood. The age groups from 35 to 55 years were found
84 to represent 24.2% of total. From above mentioned results, we find out that about 94.4% of
85 patients with superficial fungal infection are less than 56 years old. The frequency of
86 occurrence of superficial fungal infections in different age groups and sex are shown in
87 (Figure 1) and (Table 1).

88 In relation of clinical type of fungal infection to age group, in children less than 7 years,
89 tinea capitis constituted 67.6% of different superficial fungal infections, while Pityriasis
90 versicolor represented 46.2% in age group from 14-20 years.

91

92

93

94

95 Table 1. The frequency of occurrence of different superficial fungal infections in 161 patients and their relation to gender and age.
 96 gender, had no statistically significant influence ($P \leq 0.05$) on the identification of the frequency of occurrence of superficial fungal infections.
 97

Diagnosis	Number (%)													
	Different age groups (years)											gender		Total Numder(%) of each clinical type
	0-6	7-13	14-20	21-27	28-34	35-41	42-48	49-55	56-62	63-69	>=70	Male	female	
Tinea capitis	25 (67.6)	10 (47.6)	4 (30.8)	1 (4.2)	1 (5.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	23 (26.1)	18 (24.7)	41 (25.5)
Tinea faciei	3 (8.1)	0 (0.0)	0 (0.0)	1 (4.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.1)	3 (4.1)	4 (2.5)
Tinea corporis	5 (13.5)	3 (14.3)	3 (23.1)	8 (33.3)	4 (22.2)	4 (33.3)	3 (20)	4 (33.3)	0 (0.0)	0 (0.0)	1 (25)	19 (21.6)	16 (21.9)	35 (21.7)
Tinea manuum	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (5.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.4)	1 (0.6)
Tinea cruris	1 (2.7)	0 (0.0)	0 (0.0)	2 (8.3)	0 (0.0)	1 (8.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (2.3)	2 (2.7)	4 (2.5)
Tinea pedis	3 (8.1)	2 (9.5)	0 (0.0)	2 (8.3)	4 (22.2)	5 (41.7)	8 (53.3)	6 (50)	3 (75)	1 (100)	2 (50)	22 (13.7)	14 (19.2)	36 (22.4)
Tinea unguium	0 (0.0)	4 (19.1)	0 (0.0)	1 (4.2)	2 (11.1)	0 (0.0)	3 (20)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (3.4)	7 (9.6)	10 (6.2)
Pityriasis versicolor	0 (0.0)	2 (9.5)	6 (46.2)	9 (37.5)	6 (33.3)	2 (16.7)	1 (6.7)	2 (16.7)	1 (25)	0 (0.0)	1 (25)	18 (20.5)	12 (16.4)	30 (18.6)
Total Number(%) in each age group	37 (22.9)	21 (13)	13 (8.1)	24 (14.9)	18 (11.2)	12 (7.5)	15 (9.3)	12 (7.5)	4 (2.5)	1 (0.6)	4 (2.5)	88 (54.7)	73 (45.3)	161 (100)




Figure 1. Graph, representing the frequency of occurrence of superficial fungal infections in different age groups




Figure 2. frequency of each type of superficial fungal infections.

The most common fungal infection was Tinea capitis (25.5%), followed by Tinea pedis, which represented 22.4% of total cases of superficial fungal infections, followed by Tinea corporis (21.7%) and Pityriasis versicolor (18.6%). However, Tinea unguium represented only 6.2% of total superficial fungal infections. The lowest frequent superficial fungal infection was Tinea manuum (Figure2).

The pattern of specific types in relation to different age groups were characteristic. Regarding distribution of Tinea capitis in different age groups, children before 13 years old were found to be more susceptible. Adults were rarely be affected with Tinea capitis. changed pattern of frequency distribution of Tinea capitis in relation to early age groups (childhood and young adults) is given in figure 3. Conversely, Pityriasis versicolor were found to occur more frequently between 13 years and 35 years old (Figure 4). Tinea pedis were more frequent in the age groups from 28 to 55 years (Table1).




Figure 3. Change pattern of frequency distribution of tinea capitis in relation to early age groups (childhood and young adults)




Figure 4. Frequency distribution of Pityriasis versicolor in different age groups.

Discussion:

The real prevalence of superficial mycoses of skin in Libya is not known because of SFI are not noticeable diseases.⁸

In this study the mean age of patients with SFI was 25.3 years which is relatively similar to results of studies conducted in Kosova (30.8)⁹ and in Monastir region in Tunisia (33 years).¹⁰

The frequency of fungal infections was dramatically reduced after 55 years of age and this corroborate the finding in one Egyptian study¹¹ and opposite the finding of study conducted in Malaysia, in which the most of patients were older than 50 years.¹² This variation may be related to variations in demography, geography and exposure to risk factors.

Male to female ratio in our study was 1.2, identical to result in study conducted in Malaysia, and opposite what was found in Tunisia with male to female sex-ratio (0.82).^{10,12}

In this study, the most susceptible age group was children less than 7 years, similar to what was found in Tripoli (Libya)⁸, in which the majority of patients (85%) were children less than 15 years old, and also similar distribution was found in patients attending to some hospitals in Egypt and the susceptibility was high to tinea capitis, same to our finding.¹¹ The second

more affected age group in our study was from 21 years up to 27 years, showing more frequent Pityriasis versicolor. While in study conducted by (Kaur *et al*, 2015) the age group 21–30 years was the most affected followed by age group 31–40 years.⁶

In study about childhood dermatomycoses in Sfax Hospital in Tunisia, They have found that Tinea capitis was the most frequent before 13 years of age, and Pityriasis versicolor was the most frequent after 13 years of age. This finding corroborated with our results.¹³

In our study tinea capitis represented 25.5% of all superficial mycosis, taking in consideration that candidiasis were not involved in our study, this distribution of Tinea capitis is similar to that found in patients attended to three hospitals in Egypt and Sfax in Tunisia, which found to be 69.4% in Tunisia and 28.6% in Egypt and followed with Tinea pedis in our study and in Egypt, but followed with Tinea corporis in Sfax (Tunisia). Tinea corporis in our study has represented 21.7% of total while has only formed about 15% in some hospitals in Egypt.^{11,13}

Conclusion:

Fungal infection is worldwide health problem and this study has improved our understanding of epidemiology of such common SFI in Zliten area. About 22.9% of infected patients were less than 7 years old , followed by age group from 21-27 years(14.9%). In contrast, Infection was less frequent after age of 55 years (5.6%). The most frequent clinical type was Tinea capitis (25.5%), followed by Tinea pedis, Tinea corporis and Pityriasis versicolor in descending order (22.4%,21.7%,18.6%).

References:

- 1- Hay R.J. and Ashbee H.R. Mycology. Burns T., Breathnach S., Cox N., Griffiths C. *Rook's Textbook of clinical dermatology*. (8. th ed.) 2010: Wiley-Blackwell; 36.1-36.92.
- 2- El-Ghameriny M.S. Cutaneous fungal infections. *Ghameriny's textbook of clinical dermatology* 2014; 26: 903-986.
- 3- Havlickova B, Czaika V, Friedric M. Epidemiological trends in skin mycoses worldwide. journal compilation.2008; Blackwell publishing Ltd. Mycoses, 51(suppl. 4): 2-15.
- 4- Marques SA, Robles AM, Tortorano AM, Tuculet MA, Negroni R, Mendes RP. Mycoses associated with AIDS in the Third World. *Med Mycol*. 2000;38 Suppl 1:269–279.
- 5- Andrews MD, Burns M. Common tinea infections in children. *American family physician* 2008; 77:1415-1420.
- 6- Kaur R, Panda P.S , Sardana K, Khan S. Mycological Pattern of Dermatomycoses in a Tertiary Care Hospital. *Journal of Tropical Medicine*. 2015; 1: 1-5.
- 7- Monthly Report on population number and statistics of Zliten Office for Civil Registration. Dec. 2015.
- 8- Ellabib M.S, Khalifa Z. M. DERMATOPHYTES AND OTHER FUNGI ASSOCIATED WITH SKIN MYCOSES IN TRIPOLI, LIBYA , *Ann Saudi Med* 2001;21(3-4):193-195.

- 175 9- Kocinaj A. F, Kotori M. G, Koraqi A, Fida M. Dermatophytes frequency and localization sites. *Med*
176 *Arh.* 2015 Feb; 69(1):58-59.
- 177 10- Khorchani H, Haouet H, Amri M, Zanned I, Babba H, Azaiz R. Epidemiological and clinical profile of
178 superficial mycoses in Monastir region (Tunisia). *Arch Inst Pasteur Tunis.* 1996 jul-Oct ;73(3-4):179-
179 84.
- 180 11- Abd Elmegeed Al Sh. M. , Ouf S.A., Moussa T. A.A, Eltahlawi S.M.R. Dermatophytes and other
181 associated fungi in patients attending to some hospitals in Egypt. *Brazilian journal of Microbiology*
182 *4(2015); 3:799-805*
- 183 12- Murtaza M, Rajainthran S, George B. A Mycological Study of Superficial Mycoses at the Skin Clinic
184 in Sabah, Malaysia. *International Journal of Pharmaceutical Science Invention* ISSN (Online): 2319 –
185 6718, ISSN (Print): 2319 – 670X www.ijpsi.org Volume 2 Issue 3 || March 2013 || PP.45-4
- 186 13- Sellami A, Sellami H, Makni F, Mezghani S, Cheikh-Rouhou F, Marrekchi S, Turki H, Ayadi A.
187 Childhood dermatomycoses study in Sfax Hospital, Tunisia. *Mycoses.* 2008 Sep; 11-51(5):451-4.