

Short Research Article

Title:

Prevalence of soil-transmitted helminths and intestinal protozoa in shanty towns of Libreville, Gabon

ABSTRACT

Background: Intestinal parasitic infections (IPIs) are significant public health problems in sub-Saharan Africa. Establishment of effective control strategies such as mass drug administration of anthelmintic are implementing while recent local data in urban areas are lacking.

Aim: to assess the frequency of IPIs and coparasitism in patients from shanty towns of Libreville, the capital city of Gabon.

Study Design: Cross-sectional and observational study.

Place and Duration of Study: Department of Parasitology Mycology at the Université des Sciences de la Santé, Libreville, from February to April 2014.

Methods: Stool samples collected from 101 individuals, aged between 13 and 74 years old, were analysed by direct microscopic examination, merthiolate-iodine-formol concentration and coproculture. Age, education level, type of house and existence of latrine were also recorded.

Results: Among analysed stools, 75.2% (n=76/101) were infected with at least one parasite. Protozoa represented 94.3% of the detected parasites with *Blastocystis hominis* being the predominant species (41.6%; n=42/101). *Entamoeba histolytica/dispar* (6.9%; n=7/101) and *Giardia intestinalis* (2.0%; n=2/101) were the less common protozoa. Soil transmitted helminths (STH) were found in 7.9% (n=8/101) of stool examined, *Trichuris trichiura* (5.9%) was the most common helminth detected, followed by *Ascaris lumbricoides* (2.0%), hookworm (1.0%) and *Strongyloides stercoralis* (1.0%). Coparasitism was present in 35.6% (n=36/101) of positive cases. Dual and triple infections accounted for 23.8% and 6.9%, respectively. Having no education and living in a wooden

house were associated with IPIs ($p<0.01$), particularly STH which were more frequent among women and children ($p<0.01$).

Conclusion: This study reports a high frequency of intestinal parasites infection in shanty towns of Libreville. An unexpected higher prevalence of protozoa compared to STH is also noticed.

INTRODUCTION

Intestinal parasitic infections (IPIs) due to soil-transmitted helminth (STH) and protozoa infections have been recognized as neglected tropical diseases (NTDs) and as one of the most significant causes of illnesses among disadvantaged communities from SubSaharan Africa [1,2]. In 2010, it was estimated that IPIs result in 400 to 800 million illness cases [3]. These infections are ubiquitous with high prevalence among the poor and socioeconomically deprived communities who undergo overcrowding, poor environmental sanitation, low level of education and lack of access to safe water, warm climate, lack of personal or environmental hygiene, and poor health or nutritional status [4]. These factors increase the risk of being contaminated, trapping exposed populations in a perennial cycle of poverty [1,4]. IPIs identification is important, since some are due to pathogenic organism and, others considered up to now as non-pathogenic, seem to induce intestinal and allergic symptoms that can be chronic. Indeed, STH are recognized as pathogens inducing clinical symptoms while the pathogenic role of protozoan, other than *Entamoeba histolytica* and *Giardia intestinalis* such as *Blastocystis hominis*, remains either controversial or less investigated [6]. Previous studies reported that *B. hominis* is probably the most common human gut protozoan in the world, with >50% of prevalence in developing countries but the relationship between its presence and the clinical digestive symptoms is controversial [5,6].

Urban migration leads to the creation of precarious urban settlements, with a growth process of shantytowns characterized by poor housing conditions, narrow streets that are not paved, a lack of sanitation, clean water supply and electricity, as well as a poor accessibility to health services. This phenomenon occurs in Libreville, the capital city of Gabon where more than 60% of the total

51 population live, with a high proportion located in shanty towns. Data on intestinal parasitism are
52 scarce in this city. Those available focus on STH and were recorded more than 8 years ago in rural
53 areas while since several years, access to safe water is difficult due to frequent supply interruptions.
54 People are supplied in common water sources, such as pumps and then transport the collected
55 water to their home and store it in tanks. Thus, the risk of human and/or waterborne transmission
56 of IPIs would be actually high. Moreover, national deworming programmes are being prepared by
57 national control programs for country implementation. They consist of mass administration of
58 albendazole and praziquantel highly effective against STH whom prevalence estimates remain
59 unknown. It is therefore essential to identify the real burden and reservoirs of these parasites, to
60 facilitate the establishment of effective control strategies. Furthermore, while these programs focus
61 on preschool-aged and school-aged children from semi-urban and rural areas, important gaps
62 regarding IPIs among adults exist. During the weekly consultation performed at the Department of
63 Parasitology Mycology (DPM), patients with symptoms such as pruritus and or abdominal pain are
64 routinely screened for blood and IPIs. The present study was designed to estimate the frequency of
65 IPIs in patients from shanty towns of Libreville consulting at the DPM during a 4 months period.

66 **PATIENTS AND METHODS**

67 **Study site**

68 This was a cross-sectional and observational study performed at the Department of Parasitology
69 Mycology (DPM) of the Faculty of Medicine of Libreville, from February to April 2014. The study was
70 conducted in Libreville, the capital city of Gabon, in Central Equatorial Africa. Libreville is located on
71 the northwest coast and is built on rugged ground, with numerous hills and valleys crossed by
72 streams more or less important that flow into the estuary. The population of Libreville was
73 estimated at 850,000 inhabitants in 2014, representing almost 54% of the total population. Urban
74 density is high, with 250 to 300 inhabitants per km², and urbanization is anarchic and uncontrolled.
75 Thus, shanty towns occupy almost two third of the urban space in Libreville; the 3rd and the 6th

76 districts are shanty towns with approximately 150 000 inhabitants who live in precarious conditions.
77 In Gabon, the climate is equatorial, characterized by an alternation of two dry (December–January
78 and June–September) and two rainy (February– May and September–December) seasons. The
79 average annual rainfall is about 1,600–1,800 mm. The mean temperature is about 24°C.

80

81 **Study population and procedures**

82 Patients seen during medical consultations performed at the DPM of the Faculty of Medicine were
83 invited to participate to the study. Among the forty to sixty patients seen weekly, approximately one
84 half undergo a biological testing due to the presence of clinical signs suggestive of parasitic infection,
85 or as an additional test for a global check-up. Individuals with a negative diagnosis of filariasis and
86 those presenting symptoms likely associated with IPIs such as pruritus or urticaria, digestive
87 symptoms or a combination of both signs, usually benefit from stool examination. In the present
88 study, patients were included if they met the following criteria :to live in the 3rd or the 6th district of
89 Libreville, to provide a stool sample, not leaving their place of residence for at least six months, to
90 have available parasitological test results, to have given their oral informed consent for data use.
91 The demographic, socioeconomic and sanitary information **as well as symptoms** were collected by
92 trained field researchers, through semi-structured questionnaires. The interviewer wrote down the
93 answers and the questionnaire was checked before releasing the patient. In case of illiteracy,
94 interview was performed in local language with the help of the person accompanying the patient for
95 the consultation. All collected stool samples **were macroscopically examined**, then prospectively
96 analysed by routinely used standard methods: direct microscopic examination of a small amount of
97 stool mixed in a saline solution drop (0.9% sodium chloride), or fixed stools with 10% formaldehyde,
98 and after the merthiolate-iodine-formaldehyde coloration and concentration (MIFc) of eggs, cysts
99 and vegetative forms according to Sapero and Lawless [7]. MIFc is a simple and faster colouring
100 method which consists of concentrating approximately 1 gram of fecal sample crashed in the test

solution and stained with 5% lugol solution. The combination of MIF and lugol for the coloration has a good sensitivity for the identification, the fixation and the conservation of vegetative forms and cyst of protozoa. Iodine and eosin facilitate the recognition and differentiation by well-trained technician of the nuclear membrane and other cytoplasmic inclusion which are specific for each species. Species identification is carried out as follows. First, the marked cyst or vegetative form is detected at magnification x10 or x40; after perfectly centred and an addition of a drop of immersion oil on the slide, the parasite is recognized at magnification x 1000. On a bright background (light pink cytoplasm), cysts stand clear, the nuclei appear as clearly drawn in ink or red orange, the nuclear membrane appears in dark brown, chromatin appears darker. The siderophilic bodies remain colourless and refracting and vacuoles are mahogany. Red cells appear bright red [8].

Parasite coproculture based on the Harada and Mori technique was performed for the detection of helminth larvae [9]. Two trained laboratory technicians undertook the examination using optical microscopes. The result was considered positive if eggs or larvae (for helminths) or cysts/vegetative forms (for protozoa) were found by at least one method. They were considered negative if the entire slide or pellet did not reveal any parasite.

Ethical considerations

All tests performed were part of the routine diagnosis of IPIs in the DPM. Oral informed consent for the use of data was obtained from each participant after the study had been explained. An oral assent was sought from children aged between 12 and 18 years with the oral consent of their parents. Results were reported within three days and positive patients were seen by the physician and treated according to the national recommendation. Data analysis was performed in agreement with the Gabonese Ministry of Health.

Statistical analysis

Data collected were entered into an Excel spreadsheet and analysed using the Stata® version 12.1 software (StataCorp LP, Texas, USA). The statistical tests performed included the Pearson's Chi-square and Fischer's exact for group comparisons. Statistical significance was set at $p < 0.05$.

RESULTS

Data from 101 individuals with a median age of 40[24-54] years were analysed. Their characteristics are presented in Table 1. Almost half (n=44, 43.6%) had a secondary school level, 35 (34.7%) lived in houses made of wood or sheet metal and 53 (52.5%) used latrines. Overall, 26 patients presented digestive symptoms, as single symptoms in 10 patients (9.9%) and associated with pruritus in 16 patients with abdominal pain (n=13) or with diarrhoea (n=3).

IPIs were detected in 76 (75.2%) stool samples with a predominance of protozoa among infected ones (n=76/76; 100.0%) (Table 2). Helminth infections were always associated with protozoa species in positive samples (Table 2 and 3). The most commonly found parasite was *Blastocystis hominis* followed by *Entamoeba coli* and *Endolimax nana* (Table 2); 4 diarrheic out of the 7 stools with *Entamoeba histolytica/dispar* carried both vegetative forms and cysts. *Trichuris trichiura* (5.9%) was the more frequently detected STH species while larva of hookworm and *Strongyloides stercoralis* were rarely present (Table 2). Patients with polyparasitism carried two to five different species (Table 1). The rate of infected samples with pathogenic protozoa including *B.hominis* was higher than that of non-pathogenic (n=42/75; 56.0 % versus n=25/75; 33.3%; $p<0.01$). Cysts of *Iodamoeba butschlii* were not detected. One sample infected with *E. histolytica/dispar* had also vegetative forms of *Chilomastix mesnili*. Among the five samples infected by cysts or vegetative forms of *Entamoeba hartmanni*, three also had *B.hominis* and *E.coli*, and two *E. histolytica/dispar* and other parasites. *B.hominis* was the only parasite found in samples from 11 patients with digestive symptoms.

The prevalence of IPIs was significantly higher among individuals who live in a wooden house and among those who had no education ($p<0.01$) (Table 4). Coparasitism rate (27.8%; n=25/90) was almost twofold lower among adults compared to children (54.5%; n=6/11) but this difference was

not statistically significant ($p=0.14$). Pathogenic species were significantly more prevalent in case of illiteracy (75.0% *versus* 29.8%; $p<0.01$). STHs were more frequently found among children ($p<0.01$) and women ($p<0.01$) (Table 4). The same trend was observed in the group of patients who live in a wooden house ($p=0.06$) as well as in the group of illiterate ($p=0.09$), while no difference was found according to the type of toilets (Table 4). No socio-demographic characteristic-related differences were found for intestinal protozoa infection.

DISCUSSION

IPIs mapping is essential for the design or the improvement of MDA strategies. National deworming programmes were recently included in the package of strategies for the control of endemic parasitic neglected diseases in Gabon. Thus, data should be generated and available for the national control program. However, accurate data are scarce in the country, where only those from rural areas or very old ones are available [10,11].

The present cross-sectional study determining the prevalence of soil transmitted helminths and intestinal protozoa among 101 individuals revealed that intestinal parasitic infections are frequent in shanty towns of the urban city of Libreville. The purpose of this work was not to induce a decision at country level by policy makers. It was designed as a pilot study performed in an urban city and among adults. Indeed, they represented the majority of the included individuals; additionally, 75 out of the 101 patients did not present any digestive symptoms. Thus, as demonstrated elsewhere, they could represent a subset of the parasite reservoir living in shanty towns, providing important information for public health [12,13]. Moreover, weekly observations generated by the results of the lab diagnosis showed a low frequency of STH which are targeted by deworming programs. The present work, although obtained in a small sample size of symptomatic patients highlights the predominance of protozoan parasites. Further studies in urban and rural areas with higher sample size and in all age populations would confirm the present findings.

173 The prevalence of IPIs was 75.2%, similar to rates reported more than 20 years ago in Libreville
174 (76%), in an eastern rural area (100%) as well as recently found in a population of Tanzanian adults
175 (83%) [14-16]. Such prevalence are higher than those found at Lambaréné, a rural city located
176 almost 300km from Libreville (49%) and in other countries such as Ethiopia (45.3), Cameroun
177 (44.1%) and Rwanda (50%), this infection rate is higher. [17-20]. This difference is likely due to the
178 concomitant detection of helminths and protozoa by the MIF concentration. Indeed, studies
179 previously performed in the country focused on helminth species and Kato-Katz was used for
180 parasite detection. Moreover, study populations were children and pregnant women who are more
181 often infected by STH [14,17,21,22].

182 Commensal enteroparasites warrant a specific attention because their transmission pathways can be
183 identified by behaviours patterns and other conditions that contribute to the spread of other
184 diseases via the same route of transmission [1,23]. Indeed, knowledge of transmission patterns and
185 the prevalence of intestinal parasitism are useful when implementing preventive strategies at a
186 public health level. Prevalence rates of *E.coli*, *E.nana*, *E.hartmanni* are considerable, underlining the
187 poor living conditions of inhabitants from these areas.

188 The predominance of protozoa, more precisely of pathogenic ones, is in agreement with data from
189 Ethiopia (70.8% *versus* 5.2%) and Rwanda (54.5% *versus* 20), but in contrast with others [2,21,24,25].
190 Surveys performed 30 years ago in Gabon, showed prevalence of STH infections of 75% or higher
191 [14,16]. Frequent self-medication with antihelminthic drugs as well as deterioration of populations
192 living conditions could partly explain the observed profile of intestinal parasitism. Unplanned urban
193 expansion with environmental components such as water, food, cooking facilities, housing, and
194 health-care are also risk factors for IPIs. With the frequent water supply shortage, common sources
195 of water were created, increasing the risk of contamination during transporting and storing of
196 drinking water supply between the source and the point of use, as demonstrated elsewhere[26,27].

Otherwise, intestinal protozoa such as *E.histolytica* and *G.intestinalis* have clinical importance and impact on the quality of life of infected individuals. Their frequency is relatively low, in accordance with data obtained in other areas of the country and outside Africa. However, 4 out the 7 stools infected with *E.histolytica* contained both cysts and vegetative forms. Considering that molecular analysis often discriminate *E.histolytica* among collected cysts, actions against this pathogenic amoeba should be envisaged for therapeutic or preventive management [28]. Contamination of drinking water with *Giardia* cysts can occur during transporting and storing of water as population uses containers to transfer water from the dams or wells to the houses where water is stored to be used. Moreover, faecal contamination of water between the source and the house is reported [27].

B.hominis was the most predominant pathogen species. It was found as a mono-infection in 11 out of the 26 symptomatic patients. Its pathogenic role, although controversial, is being highlighted by several authors. Its association with clinical sign such as urticarial, abdominal pain, irritable bowel syndrome and diarrhoea in the absence of another aetiology was already demonstrated [29-30]. Screening of *B.hominis* should be envisaged in symptomatic patients, as well as during epidemiological studies.

Compared to *A.lumbricoides* and hookworm, *T.trichiura* was the most frequently detected STH. Its prevalence is high with an estimation 464.6 million people infected worldwide, mainly in sub-Saharan Africa and Asia [1]. A possible explanation is the common self-medication practice with albendazole or mebendazole in Libreville, both drugs do not have 100% efficacy against *T.trichiura* [19-20].

Our study has some limitations. The sample size was small and only one stool sample was collected from each participant. It was hospital-based, not performed at a community level. Furthermore, the intensity of IPIs was not determined because the Kato-Katz technique was not used. However, MIF coloration and concentration have a good sensitivity for common STH and protozoan species detection, including schistosomiasis, although kato-katz remains the reference method for

Schistosoma mansoni detection and quantification. This species as well as *Taenia spp* eggs are not endemic in Gabon. Additionally, in the DPM, patients are treated based on the presence of parasite whatever the parasite density. Finally, molecular analysis which could differentiate *E.histolytica* from *E.dispar*, as well as *Blastocystis hominis* strains, was not performed.

Although this study was not aimed at analysing epidemiological factors associated with IPIs carriage, and the small sample size which probably underpowered the statistical analysis, some factors were found associated either with the high prevalence of IPIs (living in a wooden house and being illiterate) or with STH carriage (age, sex, type of house) in agreement with reports from others [5,18,21]. Other factors such as poor individual hygiene, bad eating habits, low socioeconomic status, poor living conditions, insufficient knowledge and practice of correct sanitation could also be involved.

CONCLUSION

Intestinal parasitic infections are highly prevalent and should be considered as major public health concerns among the poor and socioeconomically communities living in some shanty towns in the capital city of Libreville, Gabon. Further studies should be performed to confirm the predominance of protozoa infections among IPIs and to identify IPI transmission mechanisms in order to design appropriate control strategies. If the low frequency of STH is confirmed, the design of actual deworming programmes which are based on antihelminthic mass administration should be revised.

REFERENCES

1. Hotez PJ, Kamath A. Neglected tropical diseases in sub-saharan Africa: review of their prevalence, distribution, and disease burden. PLoS Negl Trop Dis. 2009;3(8):e412.
2. Ngui R, Ishak S, Chuen CS, Mahmud R, Lim YAL. Prevalence and Risk Factors of Intestinal Parasitism in Rural and Remote West Malaysia. PLoS Negl Trop Dis. 2011;5(3):e974.
3. Pullan RL, Smith JL, Jasrasaria R, Brooker SJ. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. Parasite Vect. 2014; 7:3

- 248 4. Mehraj V, Hatcher J, Akhtar S, Rafique G, Beg MA. Prevalence and factors associated with
249 intestinal parasitic infection among children in an urban slum of Karachi. *PLoS One*.
250 2008;3(11):e3680.
- 251 5. Strunz EC, Addiss DG, Stocks ME, Ogden S, Utzinger J, Freeman MC. Water, sanitation,
252 hygiene, and soil-transmitted helminth infection: a systematic review and meta-analysis.
253 *PLoS Med*. 2014;11(3):e1001620.
- 254 6. Roberts T, Stark D, Harkness J, Ellis J. Update on the pathogenic potential and treatment
255 options for *Blastocystis* sp. *Gut Pathog*. 2014;6:17.
- 256 7. Stenzel DJ, Boreham PF. *Blastocystis hominis* revisited. *Clin Microbiol Rev*. 1996;9(4):563-84.
- 257 8. Saperio Jj, Lawless Dk. The MIF stain-preservation technic for the identification of intestinal
258 protozoa. *Am J Trop Med Hyg*. 1953;2(4):613-9.
- 259 9. Petithory JC, Audoin-Guidon, Chaumeil C, Hoti Sang. Amibes et flagellés intestinaux: leur
260 diagnostic microscopique. *Cahier de formation Biologie médicale. Bioderma*. 1998; 11.
- 261 10. Harada U, Mori, O. A new method for culturing hookworm. *Yonago acta med*. 2055; 1:177-9.
- 262 11. Gendrel D, Kombila M, Khayati A, Bourdil M, Engohan E, Richard-Lenoble D. Early onset of
263 intestinal parasitic infections in infants in Equatorial Africa. *Ann Pediatr*. 1983;30(6):453-6.
- 264 12. Zeukeng F, Tchinda VHM, Bigoga JD, Ndzi CHTES, Abonweh G, Makoge V, et al. Co-infections
265 of Malaria and Geohelminthiasis in Two Rural Communities of Nkassomo and Vian in the
266 Mfou Health District, Cameroon. *PLoS Negl Trop Dis*. 2014;8(10):e3236.
- 267 13. Abdulsalam AM, Ithoi I, Al-Mekhlafi HM, Khan AH, Ahmed A, Surin J, Mak JW. Prevalence,
268 predictors and clinical significance of *Blastocystis* sp. in Sebha, Libya. *Parasit Vectors*.
269 2013;6:86.
- 270 14. Gendrel D, Richard-Lenoble D, Kombila M, Dupont C, Moreno JL, Gendrel C, Nardou M,
271 Chaussain M. Influence of intestinal parasitism on lactose absorption in well-nourished
272 African children. *Am J Trop Med Hyg*. 1992;46(2):137-40.
- 273 15. Barda B, Ianniello D, Zepheryne H, Rinaldi L, Cringoli G, Burioni R, Albonico M. Parasitic
274 infections on the shore of Lake Victoria (East Africa) detected by Mini-FLOTAC and standard
275 techniques. *Acta Trop*. 2014;137:140-6.
- 276 16. Garin Y, Languillat G, Beauvais B, Tursz A, Larivière M. Parasitisme intestinal à l'Est du Gabon.
277 *Bull Soc Pathol Exot Filiales*. 1978 ;71(2):157-64.
- 278 17. Adegnika AA, Ramharther M, Agnandji ST, Ateba Ngoa U, Issifou S, Yazdanbakhsh M,
279 Kremsner PG. Epidemiology of parasitic co-infections during pregnancy in Lambaréné,
280 Gabon. *Trop Med Int Health*. 2010;15(10):1204-9.
- 281 18. Aklilu A, Kahase D, Dessalegn M, Tarekegn N, Gebremichael S, Zenebe S, Desta K, Mulugeta
282 G, Mamuye Y, Mama M. Prevalence of intestinal parasites, salmonella and shigella among

283 apparently health food handlers of Addis Ababa University student's cafeteria, Addis Ababa,
284 Ethiopia. BMC Res Notes. 2015;8:17.

285 19. Niyizurugero E, Ndayanze JB, Bernard K. Prevalence of intestinal parasitic infections and
286 associated risk factors among Kigali Institute of Education students in Kigali, Rwanda. Trop
287 Biomed. 2013;30(4):718-26.

288 20. Korkes F, Kumagai FU, Belfort RN, Szejnfeld D, Abud TG, Kleinman A, Florez GM, Szejnfeld T,
289 Chieffi PP. Relationship between intestinal parasitic infection in children and soil
290 contamination in an urban slum. J Trop Pediatr. 2009;55(1):42-5.

291 21. Njunda AL, Fon SG, Assob JC, Nsagha DS, Kwenti TD, Kwenti TE. Coinfection with malaria and
292 intestinal parasites, and its association with anaemia in children in Cameroon. Infect Dis
293 Poverty. 2015 Oct 6;4:43.

294 22. Bechir M, Schelling E, Hamit MA, Tanner M, Zinsstag J. Parasitic infections, anemia and
295 malnutrition among rural settled and mobile pastoralist mothers and their children in Chad.
296 Ecohealth. 2012 Jun;9(2):122-31.

297 23. LaBeaud AD, Nayakwadi Singer M, McKibben M, Mungai P, Muchiri EM, McKibben E,
298 Gildengorin G, Sutherland LJ, King CH, King CL, Malhotra I. Parasitism in Children Aged Three
299 Years and Under: Relationship between Infection and Growth in Rural Coastal Kenya. PLoS
300 Negl Trop Dis. 2015 ;9(5):e0003721.

301 24. Hürlimann E, Yapi RB, Hounbedji CA, Schmidlin T, Kouadio BA, Silué KD, Ouattara M,
302 N'Goran EK, Utzinger J, Raso G. The epidemiology of polyparasitism and implications for
303 morbidity in two rural communities of Côte d'Ivoire. Parasit Vectors. 2014 Feb 25;7:81.

304 25. Al-Delaimy AK, Al-Mekhlafi HM, Nasr NA, Sady H, Atroosh WM, Nashiry M, Anuar TS, Moktar
305 N, Lim YA, Mahmud R. Epidemiology of intestinal polyparasitism among Orang Asli school
306 children in rural Malaysia. PLoS Negl Trop Dis;8(8):e3074

307 26. Wright J, Gundry S, Conroy R (2004) Household drinking water in developing countries: a
308 systematic review of microbiological contamination between source and point of use.
309 Tropical Medicine and International Health.2004;9:106-17

310 27. Omar M, Mahfouz A, Moneim M. The relationship of water sources and other determinants
311 to prevalence of intestinal protozoal infections in a rural community of Saudi Arabia. J
312 Commun Health.1995;20: 433-40

313 28. Calegar DA, Nunes BC, Monteiro KJ, Santos JP, Toma HK, Gomes TF, Lima MM, Bóia MN,
314 Carvalho-Costa FA. Frequency and molecular characterisation of *Entamoeba histolytica*,
315 *Entamoeba dispar*, *Entamoeba moshkovskii*, and *Entamoeba hartmanni* in the context of
316 water scarcity in northeastern Brazil. Mem Inst Oswaldo Cruz. 2016;111(2):114-9.

29. Wawrzyniak I, Poirier P, Viscogliosi E, Dionigia M, Texier C, Delbac F, Alaoui HE. Blastocystis, an unrecognized parasite: an overview of pathogenesis and diagnosis. *Ther Adv Infect Dis* 2013;1(5):167-78.
30. Verma R, Delfanian K. Blastocystis hominis associated acute urticaria. *Am J Med Sci*. 2013;346(1):80-1.
31. Levecke B, Montresor A, Albonico M, Ame SM, Behnke JM, Bethony JM, Noumedem CD, Engels D, Guillard B, Kotze AC, Krolewiecki AJ, McCarthy JS, Mekonnen Z, Periago MV, Sopheak H, Tchuem-Tchuente LA, Duong TT, Huong NT, Zeynudin A, Vercruysse J. Assessment of anthelmintic efficacy of mebendazole in school children in six countries where soil-transmitted helminths are endemic. *PLoS Negl Trop Dis*. 2014 Oct 9;8(10):e3204.
32. Keiser J, Utzinger J. Efficacy of current drugs against soil-transmitted helminth infections: systematic review and meta-analysis *JAMA* 2008;299:1937-48.

Table1: Characteristics of the study population

CHARACTERISTICS

	Median	25 th –75 th Quartile
Age	40	24-54
	N	%
Adults	90	89.1
Female	66	65.3
Type of house		
- Wooden house	24	23.8
- Brick house	66	65.3
- Metal sheet house	11	10.9
Educational level		
- None	11	10.9
- Primary school	33	32.7
- Secondary school	48	47.5
- High school	9	8.9
Type of toilets		
- Interne toilets	48	47.5

352	- Latrines	53	52.5
353	Reported dewormed the last six months	15	14.8
354	Symptoms		
355	- Abdominal pain	23	22.8
356	- Diarrhoea	4	4.0
357	- Pruritus or urticaria	91	90.1

358

359

360 **Table 2 : Profile of intestinal parasitism**

361

362	PARASITE CARRIAGE	N	%
363			
364	Helminths	8	7.9
365	Protozoa	76	75.2
366	Monoparasitism	40	39.6
367	Coparasitism	36	35.6
368	Protozoa + helminths	8	7.9
369	2 parasites	24	23.8
370	3 parasites	7	6.9
371	4 parasites	4	4.0
372	5 parasites	1	1.0

373

374

375

376

377

378

379

380

381

382

383

384

Table 3 : Prevalence of parasite species detected in the study population

SPECIES	TOTAL		MONOPARASITISM		COPARASITISM	
	N	%	n	%	n	%
Helminths						
<i>Ascaris lumbricoides</i>	2	2.0	0	0.0	2	100.0
<i>Hookworm</i>	1	1.0	0	0.0	1	100.0
<i>Strongyloides stercoralis</i>	1	1.0	0	0.0	1	100.0
<i>Trichirus trichiura</i>	6	5.9	0	0.0	6	100.0
Protozoa						
<i>Blastocystis hominis</i>	42	41.6	17	40.5	25	59.5
<i>Endolimax nana</i>	28	27.7	8	28.6	20	71.4
<i>Entamoeba coli</i>	33	32.7	12	36.4	21	63.6
<i>Entamoeba hartmanni</i>	7	6.9	1	14.3	6	85.7
<i>Entamoeba histolytica/dispar</i>	7	6.9	1	14.3	6	85.7
<i>Giardia intestinalis</i>	2	2.0	1	50.0	1	50.0

409 **Table 4: Frequency of IPIs according to socio-demographic data**

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

CHARACTERISTICS		INFECTED		NON INFECTED		STH		PATHOGENIC PROTOZOA	
		N	%	N	%	N	%	N	%
Age									
-	Adults	67	64.4	23	25.6	6	6.7	36	40.0
-	Children	9	82.8	2	18.2	2	18.2*	6	54.4
Sex									
-	Female	48	72.7	18	27.3	6	9.1*	24	36.4
-	Male	28	80.0	7	20.0	2	5.7	18	51.4
Type of house									
-	Wooden house	24	100.0*	0	0.0	4	16.7	12	50.0
-	Brick house	43	65.1	23	34.9	3	4.5	23	31.8
-	Metal sheet house	9	81.2	2	10.8	1	9.1	7	63.6
Educational level									
-	None	11	100.0	0	0.0	2	18.2	9	81.8
-	Primary school	23	69.7	10	30.3	4	9.3	18	54.5
-	Secondary school	37	77.1	11	22.9	2	4.2	26	68.4
-	High school	7	77.8	2	22.2	0	0.0	5	55.6
Type of toilets									
-	Interne toilets	38	79.2	10	20.8	2	4.2	20	62.5
-	Latrines	38	71.7	15	28.3	6	11.3	29	54.7

433 * : difference statistically significant

434