

Coital abstinence and High Risk Sexual behaviour among husbands of postpartum women, in Ikenne, Southwestern Nigeria

ABSTRACT

Aims: To determine the prevalence of high risk sexual behaviour and its relationship to the duration of coital sexual abstinence in husbands of postpartum women.

Study design: Cross sectional descriptive study.

Place and Duration of Study: Ikenne Local Government Area, in Ogun State, South Western Nigeria between, December 2014 and May 2015.

Methodology: Data were obtained with the aid of semi-structured interviewee administered questionnaire from 771 husbands of postpartum women using the multi-stage sampling technique.

Results: The prevalence of High risk sex in the population was 10.6%. The duration of coital sexual abstinence was the most important risk factor ($P<.001$), while previous extramarital sexual relationship ($P<.001$, $OR=41.70$, $95\%CI=18.07-96.07$) and husband's knowledge of his own HIV status ($P=.03$, $OR=1.71$) were also significant determinants of this occurrence. Consistent condom use during unsafe sex was 6.1%, while STIs occurred in 3.7% of the participants. Significantly longer durations of coital abstinence (8.30 ± 6.24 months) were observed in men who were rural dwellers than in urban dwelling husbands (7.16 ± 6.01 months), $P=.01$. Violent behavior against the postpartum wife during the abstinence period was reported by 1.2% of the participants.

Conclusion: High risk male sexual behavior was a consequence of prolonged postpartum sexual abstinence and a predisposition to STIs among husbands of postpartum women.

Keywords: Postpartum, Male behavior, Unsafe sex, Networking, HIV, Violence

1. INTRODUCTION

All of the Nigerian tribes practice some form of sexual proscription in the postpartum period. This practice is mandatory especially during the puerperium and most of the postpartum period unlike in pregnancy where coital abstinence is only relative, [1,2]. Medically indicated coital sexual abstinence is for the first 2-3 weeks of the puerperium to allow healing of the birth canal and prevent genital tract infection [3].

Traditionally the Yoruba tribe in Nigeria had practiced agriculture as a means of livelihood and postpartum sexual abstinence helped the women to cope with both the demands of being a mother at home and a worker on the farm. The 'natural contraception' produced by this mandatory sexual abstinence made it possible [4]. The widely accepted belief that helped to uphold this practice among the Yoruba, was that a man should not have sexual intercourse with his wife when she was nursing a baby; the concept of 'lactation taboo' [1,2,5]. It was widely believed among them that the semen could reach the breast milk and poison it, thus jeopardizing the health of the new-born when intercourse occurs before the culturally recommended time [1,2]. This abstinence period lasted until the child was weaned which may take up to two years or more [2].

High risk sex has been shown to be a very potent mechanism for the spread of the HIV virus. Indeed, reports from studies in Africa have shown that the greatest risk of HIV infection for married women was from sexual intercourse within their marriage, and the greatest source of HIV infection was unprotected sex with their infected husbands [6-8]. A report in 2008 revealed that, up to 80% of new HIV infections among heterosexual urban dwellers in Africa occurred within marital or cohabiting relationships [7]. Prior HIV infection in the husband before the marital union or HIV infection contracted through extra-marital

sexual relationship while still being married was the pathway to infection [8]. The permissiveness of African tribes, including the Yoruba, in condoning or only minimally reprimanding male extra-marital sexual activity makes married men more likely than married women to engage in high risk sex [6]. In addition, concurrent multiple sexual relationships have been postulated to heighten the risk of HIV transmission; an occurrence which is commoner among African men than in men from other parts of the world [9].

Sexual abstinence for couples in the past was possible in a polygamous context where temporary separation of spouses was frequent, particularly after childbirth, alternative sexual outlets were available in their other mutually faithful wives [5].

The family setting in Southwestern Nigeria in recent times has become increasingly monogamous, occasioned mainly by the widespread adoption of new occupations and lifestyles. These changes were brought about mainly by westernization, civilization and in many instances by religion [10]. Clearly the challenges facing the average Nigerian family today is different and the increasing economic reality has greatly reduced the practice of polygamy. Modernization also brings with it the benefit of better, more reliable and effective means of contraception, the better care of, and increasing survival of children [5]. Sub-Saharan Africa today is faced with the challenges of HIV, AIDS, hepatitis and other grave consequences of STIs such as infertility, cervical cancer and liver disease [11,12]. This belief in prolonged postpartum sexual proscription however still appears to have a strong grip and may produce sexual behavior inimical to prevention of these diseases.

The postpartum period may be an appropriate time to observe married male sexual behavior during required abstinence; because unlike pregnancy where medical or even culturally indicated coital abstinence is only relative, most of the African tribes practice coital sexual abstinence, with variations only in the duration [1,5,13]. Refusal of women to have sexual intercourse with their husbands has been found to be one of the reasons for Intimate Partner Violence (IPV) [14], the relationship between IPV and postpartum sexual abstinence is however yet to be established. In addition, healthcare workers are oftentimes faced with situations where the male partner requires counseling on sexual issues related to timing the resumption of sexual intercourse with the wife who is a nursing mother and coping mechanisms when abstinence is indicated, currently opinions vary on best options.

This study was thus aimed at determining the prevalence and patterns of high risk sexual behavior in husbands of postpartum women and its relationship to the duration of sexual abstinence in Ikenne, South-western Nigeria.

2. MATERIAL AND METHODS

This study was conducted in Ikenne Local Government Area (LGA), one of the twenty LGAs in Ogun State, South Western Nigeria. This LGA, is made up of three urban towns and two rural villages. The urban towns are Iperu, Ikenne, and Ilisan, while the rural villages are Irolu and Ogere [15]. Ikenne local government spans 137.13km² with 202,980 inhabitants (National Population Commission, 2006) [16].

It was a community-based cross sectional descriptive study involving 771 husbands of postpartum women using the multi-stage sampling technique. The LGA was divided into 5 clusters according to the 5 towns, trade union groups were then selected from each town by computerized random sampling technique. This was followed by simple random sampling to select individual participant, at selected work places. The sample size was determined using the Leslie-Kish formula $n = z^2pq/d^2$ [17]. With the precision (d) set at 0.05, the prevalence of high risk sex of 48% obtained by Lawoyin et al (2002) in a study on husbands of postpartum women was used [4,18]. The value (n=384) obtained was then doubled (n=766) to remove cluster effect.

Data were obtained over a 6-month period with the aid of semi-structured pre-coded interviewee administered questionnaire designed by the authors. The questionnaire contained sections on socio-biologic data, which had mainly open-ended questions. Other sections included were durations of coital

sexual abstinence, reasons for it and coping strategies. The occurrence and patterns of high risk sexual behavior section; contained mainly structured questions, leaving some open ended questions for questions involving duration. The questionnaire was translated to the Yoruba language and back-translated to English to ensure that it retained its meaning. A pre-test of the questionnaire was done on men attending a meeting of the commercial motorcycle riders association, in Ilshah-Remo and administrative workers of Babcock University. After obtaining verbal consent (witnessed by 2 or more persons), the participants were given the questionnaire to fill. Those who could not write had their responses entered into the questionnaire for them by trained interviewers who were doctors and could speak the Yoruba language well, such interviews lasted 20 minutes averagely. Interpreters were used for the non-Yoruba speaking respondents whose answers were also audio-recorded in addition to their questionnaire being filled for them by the interviewers. The recorded answers were transcribed by independent transcribers and checked with the responses on the questionnaire for uniformity.

The collected data were analyzed using the Statistical Software Package for Social Sciences (SPSS) version 17.0 (Chicago, Illinois). Frequency tables were made and numerical data expressed as mean $\pm$ standard deviation. The results were tested for statistical significance thus: t- test was used to compare continuous variables while chi- square was used for comparison of categorical variables. The level of statistical significance was set at *P*-value of less than .05

The puerperium was defined as the period of the first six weeks following delivery of the index baby, while the postpartum period was taken as that period lasting up to one year following the delivery of the index baby, or the interval from delivery of the index baby to a subsequent pregnancy. Postpartum coital sexual abstinence was considered as the absence of penetrative peno-vaginal sexual intercourse between the respondent and his socially, culturally or legally accepted wife in the postpartum period. A husband was considered as a man 18 years old and above who was legally, customarily, religiously married to and or lives with a woman in a union whom his or her family or surrounding society has accepted as man and wife. High risk sex was considered as the occurrence of currently married or cohabitating men reporting having had coital sexual relations with a woman who is not known or socially, culturally, legally or religiously accepted as a wife [19]. It also extended to encompass sexual intercourse with someone who was susceptible to or infected with an STI whether a condom was used or not [20]. Early resumption of coitus was considered as postpartum sexual intercourse at 6 weeks or less after delivery of a baby [21]. Prolonged coital abstinence was considered as sexual intercourse occurring for the first time 1 year or more after delivery. Intimate partner violence (IPV) was considered as the occurrence of Physical, emotional or verbal abuse against the postpartum wife.

3. RESULTS AND DISCUSSION

3.1. RESULTS

Eight hundred copies of the questionnaire were administered, but 771 were returned correctly completed, for participants who met inclusion criteria, giving a retrieval rate of 96.4%.

Table 1 highlights the socio-demographic characteristics of the respondents. Participants in the rural areas were significantly younger (mean age of 36.17 $\pm$ 6.17 years, (*P*=.03) and had lower educational levels (*P*=.01). Other characteristics such as occupation (*P*=.13), religion (*P*=.90), tribe (*P*=.11), marriage setting (*P*=.72), knowledge of own HIV status (*P*=.79) and co-habitation with postpartum wife (*P*=.48) were however not significantly different, between the rural and urban respondents.

Table 1. Socio-demographic characteristics of participants

Settlement	<i>P</i> -value
------------	-----------------

Characteristics	Rural(n=308)	Urban(n=463)	Total(n=771)	(95% CI)
Age				
Mean(years)	36.17±6.17	37.31±7.50	36.85±7.01	.03(-0.01-0.13)
Educational level				
Primary	63(49.3%)	65(50.7%)	128	.01(0.01-0.02)
Secondary	192(40.0%)	288(60.0%)	480	
Tertiary	53(32.5%)	110(67.5%)	163	
Occupation				
Unemployed	3(15.8%)	16(84.2%)	19	.13(0.12-0.14)
Unskilled	64(38.1%)	104(61.9%)	168	
Skilled	192(41.7%)	268(58.3%)	460	
Professional	49(39.5%)	75(60.5%)	124	
Religion				
Christianity	255(39.9%)	385(60.1%)	640	.90
Islam	53(40.5%)	78(59.5%)	131	
Tribe				
Hausa	25(35.2%)	46(64.8%)	71	.11(0.10-0.11)
Igbo	24(30.4%)	55(69.6%)	79	
Yoruba	259(41.7%)	362(58.3%)	621	
Marriage Setting				
Monogamous	283(40.1%)	422(59.9%)	705	.72
Polygamous	25(37.9%)	41(62.1%)	66	
Cohabit with wife				
Yes	290(40.3%)	430(59.7%)	720	.48
No	18(35.3%)	33(64.7%)	51	
Awareness of own HIV status				
Yes	170(39.5%)	260(60.5%)	430	.79

No	138(40.5%)	203(59.5%)	341
----	------------	------------	-----

127 n = number of participants

128 95%CI = 95% Confidence Interval

129

130 Table 2 reveals that 63/771 (8.2%) of the couples in Ikenne LGA, had resumed coital sexual intercourse
 131 before 6 weeks. Men in the rural towns however had significantly longer durations of postpartum coital
 132 sexual abstinence than urban men (8.30 ± 6.24 months, $P = .01$, 95% CI of -0.25-2.02). It also showed
 133 that a significantly greater proportion of rural men 94/308 (30.5%) than the urban dwelling men 95/463
 134 (20.5%), abstained for periods of 12 months and above ($P = .007$). None of the participants abstained from
 135 sex with their postpartum wives for up to 3 years or longer.

136 **Table 2. Duration of postpartum coital sexual abstinence in husbands of postpartum women**

Settlement	Duration(months)			P-value (95%CI)
	<6 weeks	6 weeks -<1 yr	1-<3yrs	
Rural (n=308)	22(7.1%)	192(62.4%)	94(30.5%)	.007
Urban (n=463)	41(8.9%)	327(70.6%)	95(20.5%)	
Total (n=771)	63(8.2%)	519(67.3%)	189(24.5%)	
Mean Duration (months)				
Rural	8.30±6.24			.01 (-0.25 to -2.02)
Urban	7.16±6.01			
Total	7.62±6.16			
	Median = 6.00			
	Range = 29.5			

137

138 The result in table 3 shows that the duration of postpartum coital abstinence was not significantly
 139 influenced by age ($P = .34$), highest level of education attained ($P = .62$) or occupation ($P = .10$). In addition,
 140 the tribe ($P = .17$), religion ($P = .85$), marriage setting ($P = 0.78$) or cohabitation with index wife ($P = .76$) were
 141 also not significant determinants of the duration of sexual abstinence in the postpartum period.

142

143 **Table 3. Socio-demographic Factors affecting the duration of postpartum coital sexual abstinence**
 144 **in husbands of postpartum women**

		Duration			P-value CI)	(95% CI)
		<6 weeks (n=63)	6 weeks-<1yr (n=519)	1-<3rs (n=189)		
Factor						
Age						
Mean		38.21±6.95	36.73±7.15	36.74±6.63	36.85±7.01	.34(0.33-0.35)
Educational Level						
Primary		7(5.5%)	92(74.2%)	29(22.7%)	128	.62(0.62-0.64)
Secondary		44(9.2%)	318(66.2%)	118(24.6%)	480	
Tertiary		12(7.4%)	109(66.9%)	42(25.7%)	163	
Occupation						
Unemployed		5(26.3%)	12(63.2%)	2(10.5%)	19	.10(0.09-1.10)
Unskilled		10(6.0%)	116(69.0%)	42(25.0%)	168	
Skilled		39(8.5%)	305(66.3%)	116(25.2%)	460	
Professional		9(7.3%)	86(69.4%)	29(23.3%)	124	
Tribe						
Hausa		7(9.9%)	46(64.8%)	18(25.3%)	71	.17(0.16-0.17)
Igbo		10(12.7%)	44(55.7%)	25(31.6%)	79	
Yoruba		46(7.4%)	429(69.1%)	146(23.5%)	621	
Religion						
Christianity		53(8.3%)	428(66.9%)	159(24.8%)	640	.85(0.85-0.87)
Islam		10(7.6%)	91(69.5%)	30(22.9%)	131	
Marriage setting						
Monogamous		58(8.2%)	472(67.0%)	175(24.8%)	705	.78(0.78-0.79)
Polygamous		5(7.6%)	47(71.2%)	14(21.2%)	66	
Cohabitation with wife						
Yes		60(8.3%)	485(67.4%)	175(24.3%)	720	.76(0.75-0.77)

No	3(5.9%)	34(66.7%)	14(27.4%)	51
----	---------	-----------	-----------	----

145

146 The reproductive health factors did not exert any significant influence on the duration of postpartum
147 sexual abstinence (Table 4).

148 **Table 4. Reproductive health factors affecting the duration of postpartum coital sexual abstinence**
149 **in husbands of postpartum women**

Factor	Duration			Total (n=771)	P-value (95% CI)
	<6 weeks	6 weeks-<1yr	1-<3rs		
	(n=63)	(n=519)	(n=189)		
Mode of delivery					
Vaginal	58(8.0%)	489(67.7%)	175(24.3%)	722	.68
Caesarean section	5(10.2%)	30(61.2%)	14(28.6%)	49	(0.67-0.69)
Episiotomy/genital laceration					
Yes	8(9.6%)	52(62.7%)	23(27.7%)	83	.63
No	55(8.0%)	467(67.9%)	166(24.1%)	688	(0.62-0.64)
Postpartum obstetric complications					
Yes	6(10.1%)	39(66.1%)	14(23.8%)	59	.86
No	57(8.0%)	480(67.4%)	175(24.6%)	712	(0.85-0.87)
Duration of breastfeeding					
	17.92±3.40	17.36±3.45	18.51±3.64	17.69±3.52	.18 (0.17-0.19)
Awareness of own HIV status					
Yes	34(7.9%)	280(65.1%)	116(27.0%)	430	.22
No	29(8.5%)	239(70.1%)	73(21.4%)	341	(0.21-0.23)
Decreased					

libido (husband)

Yes	6(8.6%)	41(58.6%)	23(32.8%)	70	.21
No	57(8.1%)	478(68.2%)	166(23.7%)	701	(0.20-0.22)

The result in figure 1 reveals that 9/771 (1.2%) of the participants in this study engaged in violent behavior against their postpartum wives during the abstinence period. The pie chart also shows that although 289/771 (37.5%) did 'nothing sexual' in that interval, a larger proportion 296/771 (38.4%) engaged in non-coital, non-organic sex with the postpartum wife.




Figure 1. Strategies for coping with postpartum sexual abstinence in husbands of postpartum women in Ikenne

History of occurrence of high risk sex was volunteered by 82 respondents, giving a prevalence of 10.6% as depicted in table 5. This table also shows that progressively longer durations of coital abstinence correlated with increasing prevalence of high risk sex ($P < .001$). A sharp increase was noted after 1 year with change in prevalence from 8.1% (6 weeks to < 1 year) to 19.6% after 1 year of abstinence. The mean duration of coital abstinence was also found to be significantly longer in participants who engaged in high risk sex (11.01 ± 6.99 months), compared with those who did not (7.21 ± 5.92 months) P -value $< .001$.

Greater proportion of men who had previous extramarital sex (77.1%), were involved in high risk sex compared with 7.5% who had no prior extramarital sexual relations ($P < .001$). Also with an odds ratio $OR = 41.70$, $P < .001$, and 95%CI=18.07-96.07), husbands who had previous extramarital sex were more likely to engage in high risk sex during their wife's postpartum period. Men who knew their HIV status were also found to be more likely to engage in high risk sex at this time ($P = .03$ and $OR = 1.71$). Other factors such as age ($p = .93$), educational level ($P = .74$), religion ($P = .44$), ethnicity ($P = .91$), settlement

172 ($P=.81$), occupation ($P=.44$), cohabitation with postpartum spouse ($P=.64$) and duration of breastfeeding
 173 ($P=.47$) were not significantly different between participants who engaged in high risk sex and those who
 174 did not (Table 5).

175 **Table 5. Determinants of High Risk Sex in husbands of postpartum women**

Factor	High Risk Sex		Total	P-value (95%CI)
	Yes	No		
	n=82(10.6%)	n=689(89.4%)	771	
Age (mean $\pm$ SD in years)	36.91 $\pm$ 7.17	36.84 $\pm$ 7.00	36.85 $\pm$ 7.01	.93(-1.54 to 1.68)
Educational level				
Primary	16(12.5%)	112(87.5%)	128	.74(0.73-0.74)
Secondary	50(10.4%)	430(89.6%)	480	
Tertiary	16(9.8%)	147(90.2%)	163	
Religion				
Christianity	71(11.1%)	569(88.9%)	640	.44
Islam	11(8.4%)	120(91.6%)	131	
Ethnicity				
Hausa	8(11.3%)	63(88.7%)	71	.91(0.89-0.91)
Igbo	7(8.9%)	72(91.1%)	79	
Yoruba	67(10.8)	554(89.2)	621	
Settlement				
Rural	34(11.0%)	274(89.0%)	308	.81
Urban	48(10.4%)	415(89.6%)	463	
Occupation				
Unemployed	1(5.3%)	18(94.7%)	19	.44(0.43-0.45)
Unskilled	21(12.5%)	147(87.5%)	168	
Skilled	51(11.1%)	409(88.9%)	460	
Professional	9(8.0%)	115(92.0%)	124	
Marriage setting				

Monogamous	75(10.6%)	630(89.4%)	705	.60
Polygamous	7(10.6%)	59(89.4%)	66	
Cohabit with wife				
Yes	78(11.0%)	630(89.0%)	708	.64
No	7(10.6%)	59(89.4%)	66	
Prior extramarital Sex				
Yes	27(77.1%)	8(22.9%)	35	<.001,OR=41.70
No	55(7.5%)	679(92.5%)	734	(18.07 to 96.07)
Awareness of own HIV Status				
Yes	55(12.8%)	375(87.2%)	430	.03,OR=1.71
No	27(7.9%)	314(92.1%)	341	(1.05 to 2.77)
Duration of breastfeeding				
Mean(months)	17.43±3.43	17.72±3.53	17.69±3.52	.47(-1.10 to 0.51)
Duration of coital abstinence				
< 6 weeks	3(4.8%)	60(95.2%)	63	<.001
6 weeks -<1 year	42(8.1%)	477(91.9%)	519	
1-< 3 years	37(19.6%)	152(80.4%)	189	
Mean duration of coital				
Abstinence (months)	11.01±6.99	7.21±5.92		<.001

OR=Odds Ratio

The result in table 6 reveals that high risk sex occurred in similar proportions in both rural and urban settlements ($P=.77$). An 'old friend' was the commonest high risk sexual partner 54/82 (65.8%) in the study population. Most 64/82 (78.0%) of the respondents, used condoms occasionally during HRS. Only 5/82 (6.1%) used a condom at every HRS act, 13/82(15.9%) never used a condom. The mean interval between delivery and first high risk sex episode for the study population was 4.26 ± 3.10 months, thus a sizable proportion of participants had engaged in high risk sex before the 5th postpartum month, while abstaining from their wives (population mean duration of coital abstinence= 11.01 ± 6.99). Occurrence of STI was reported by 3/82 of the respondents who engaged in high risk sex, giving a prevalence of 3.7%, whereas only 16 responses were received for sex with persons suspected to be infected with an STI.

Table 6. Patterns of high risk sex (HRS) in husbands of postpartum women, across the Ikenne settlements

	Settlement			
Pattern	Rural (n=34)	Urban (n=48)	Total (n=82)	P-value(95%CI)
Type of partner				
Old friend	25(46.3%)	29(53.7%)	54	.32(0.31 to 0.32)
New friend	8(36.4%)	14(63.6%)	22	
Commercial sex worker	1(16.7%)	5(83.3%)	6	
Sex with infected person(STI)				
Yes	1(25.0%)	3(75.0%)	4	.75
No	3(25.0%)	9(75.0%)	12	
Condom use for HRS				
Never	6(41.2%)	7(53.8%)	13	.92(0.91 to 0.92)
Occasional	26(40.6%)	38(59.4%)	64	
Always	2(40.0%)	3(60.0%)	5	
Occurrence of STI after HRS				
Yes	1(33.3%)	2(66.7%)	3	.63
No	33(41.7%)	46(58.3%)	79	
Delivery-HRS interval				
Mean(months)	4.29±3.30	4.23±2.99	4.26±3.10	.93(-1.32 to 1.45)
Previous extramarital sex				
Yes	10(37.0%)	17(63.0%)	27	.57
No	24(43.6%)	31(56.4%)	55	

3.2. DISCUSSION

The observed mean duration of postpartum sexual abstinence, is much longer than the 2.1±0.71 months, reported in a hospital based study in Jos, North-Eastern Nigeria among women who were 14 weeks (or less) postpartum [12]. More (8.2%) of the couples in Ikenne LGA than the reported 3.5% couples in Jos had however resumed sexual intercourse before the end of the puerperium [12]. A significantly longer duration was found in this study among husbands who were rural dwellers compared to husbands who lived in the urban areas. A relatively greater proportion of rural dwelling men abstaining for periods of 12 months and longer, compared to their rural counterparts. This finding contrasts reports from a study conducted in Ghana, where no significant difference in sexual behavior existed in husbands of postpartum women between these two settlement types [22].

Although rural dwellers had longer durations of abstinence, no difference was observed from urban dwellers in the occurrence of high risk male sexual behavior. This finding is inconsonance with a report from a study in Ibadan, South western Nigeria [4].

The results from this study agree with findings of a study in Jos where socio-demographic and obstetric factors, were not predictive of the duration of coital abstinence [12]. It however contrasts what was reported, from a study involving data from 17 countries; most of them in Africa, where obstetric factors were found to positively affect this duration [23]. It is an established belief among the Yoruba of Nigeria that sexual intercourse during breastfeeding should be forbidden because it may poison the milk, with several couples readily abstaining for durations spanning most of or even the entire length of the breastfeeding period, which may be as long as 3 years [1,2,24]. The duration of breastfeeding observed in Ikenne among the respondents is however much longer than the duration of coital abstinence. These lower durations observed, may be due to changes in sexual practices and breastfeeding patterns among couples during the postpartum period, responsible for the lower mean duration than the 2-3 years that is culturally required [1,24]. In fact no couple in Ikenne LGA, had a duration of infant breastfeeding that lasted up to 3 years. These changes may be a result of the increasing adoption of monogamous marriages, better skilled, non-agricultural occupations with increasing number of 'working class' postpartum wives and less dependence on extended family networks who often provided support and monitoring in traditional abstinence settings as found in an earlier report [22].

Non-coital, non-orgasmic sex with the postpartum wife was the commonest alternative to coital sexual intercourse practiced by the participants. This appears 'un-African' as it is dissimilar from the commonly reported traditional coping mechanisms of 'sex with another existing wife' (polygamy), sex with other acquaintances married into the family setting or sex with a commercial sex worker as reported in earlier studies [4,13]. These coping methods observed were however similar in respondents from both rural and urban settings. The prevalence of intimate partner violence against the postpartum wife was observed in this study, was lower than the 4-29% reported in a multi-country survey of intimate partner violence related to pregnancy in some other developing countries.[25] Pregnancy and lactation have been shown to be major predispositions to violence against women [14, 26].

The prevalence of high risk sex observed in the Ikenne population is much lower than the earlier reported 48% observed in Ibadan, South-western Nigeria, from a study on male sexual networking during the postpartum period [27]. High risk male sex in the postpartum period was also less prevalent in Ikenne than the estimated 42.1% in husbands of pregnant women Ibadan [27]. Participants who engaged in high risk sex were noted to have significantly longer durations of abstinence than those who did not. In fact the prevalence more than doubled after 1 year of postpartum sexual abstinence. The finding from this study that the interval from delivery to the first episode of unsafe sex was shorter than the duration of sexual abstinence suggests that these men had started having extramarital sex earlier as an alternative to cope with the period of either puerperal or postpartum abstinence from sexual intercourse with their wives.

Men who had previous extramarital sexual relationships (within index marriage) were also found to be more likely to engage high risk postpartum sex. This observation was also supported by the finding that an 'old friend' was the commonest sexual partner in this study. This is consonance with an earlier report that the avenues for postpartum men during abstinence were mostly people known to or close to them such as young wives in polygamous marriages, single girls from poor families, widows and divorced women [13].

Similarly, Knowledge of HIV status had a significant influence on occurrence of unsafe sex, as men who knew their status were more likely to engage in it. A positive HIV test rate of 2.3% was found among men involved in high risk sex in Ibadan, the investigators in that study however noted that none of these men were aware of their status, reasons given for this was that HIV testing at that time was very low [26]. Testing for HIV now is quite widespread, thus it is pertinent to know which HIV status actually influenced high risk sexual behaviour during the postpartum period. Further studies among Nigerians with combined HIV testing would be useful in this regard.

Consistent condom use during high risk sexual activity observed in this population was low. The level of consistent condom use among rural dwelling men engaging in unsafe sex in Ikenne of 5.8% was however higher than the 1.7% earlier reported from a study in the rural parts of Ibadan [27]. These rates of condom use are lower than expected as condom use in Africa has been described as a 'marker for high risk sex' [20]. Consistent use of the male condom has been shown to produce up to 80% reduction in HIV incidence [28]. The association between male sexual networking and transmission of STIs including HIV has been well established [13]. A lower prevalence of STIs was observed in men who had sexual intercourse with persons with features of STI in Ikenne than the 23.7% prevalence of STI reported in men with similar exposures in an earlier report from Ibadan [27]; the higher rates of male condom use in Ikenne may be responsible for this lower rate. The inability of the study to determine which particular HIV status influenced the occurrence of high risk sexual behavior in the participants in this study would necessitate further prospective study involving laboratory HIV testing.

4. CONCLUSION

High risk sexual behaviour in husbands of postpartum women was a consequence of prolonged postpartum sexual abstinence and previous extramarital sexual relationships significantly increased the risk of this behaviour. Low rates of consistent condom use among these men during unsafe sex put them at increased risk for STIs. The mean duration of postpartum coital abstinence in Ikenne LGA is less than culturally prescribed, showing a decreasing trend.

REFERENCES

1. Sule-Odu AO, Fakoya TA, Oluwole FA, Ogundahunsi OA, Olowu AO, Olanrewaju DM, Akesode FA, Dada OA, Sofekun EA. Postpartum sexual abstinence and breastfeeding pattern in Sagamu, Nigeria. *Afr J Reprod Health*. 2008 Apr;12(1):96-100
2. Odar E, Wandabwa J, Kiondo P. Sexual practices of women within six months of childbirth in Mulago hospital, Uganda. *Afr Health Sci* 2003;3:117-23.
3. Lipscomb K, Novy MJ. The Normal puerperium. In *Current Diagnosis and Treatment in Obstetrics and Gynaecology*. Decherney AH, Nathan L, Goodwin TM, Laufer N, (eds). 10th Edition. 2007. Pgs 222-248.
4. Lawoyin TO, Larsen U. Male sexual behaviour during wife's pregnancy and postpartum abstinence period in Oyo State, Nigeria. *J Biosoc Sci*. 2002;3: 4(1):51-63.
5. Gueye M, Van de Walle E. Some joint determinants of fertility and infant and child mortality in sub-Saharan Africa. In: *African Population Conference/Congres Africain de Population*, Dakar, Senegal, November 7-12, 1988. Liege, Belgium, International Union for the Scientific Study of Population, 1988, 3(3):1-16.
6. Smith DJ, Modern marriage, men's extramarital sex, and HIV risk in Southeastern Nigeria, *American Journal of Public Health*, 2007, 97(6):997-1005
7. Dunkle K et al., New heterosexually transmitted HIV infections in married or cohabiting couples in urban Zambia and Rwanda: an analysis of survey and clinical data, *Lancet*, 2008, 371(9631):2183-2191
8. Stephenson R. Community-Level Gender Equity and Extramarital Sexual Risk-Taking among Married Men in Eight African Countries. *International Perspectives on Sexual and Reproductive Health*, 2010,36(4):178-188
9. Hunter M, Cultural politics and masculinities: multiple-partners in historical perspective in KwaZulu-Natal, *Culture, Health and Sexuality*, 2005, 7(4):389-403
10. Onah HE, Iloabachie GC, Obi SN, Ezugwu FO, Eze JN. Nigerian male sexual activity during pregnancy. *Int J Obstet Gynecol* Vol. 2002, 76(2): 219-223
11. Drake AL, Wagner A, Richardson B, John-Stewart G. Incident HIV during Pregnancy and Postpartum and Risk of Mother-to-Child HIV Transmission. A Systematic Review and Meta-Analysis 25,2014.DOI: 10.1371/journal.pmed.1001608
12. Anzaku AS, Mikah S. Postpartum Resumption of Sexual Activity, Sexual Morbidity and Use of Modern Contraceptives Among Nigerian Women in Jos. *Annals of Medical and Health Sciences Research* 2014 4(2):210-216

13. Caldwell JC, Orubuloye IO, Caldwell P. Perceived male sexual needs and male sexual behavior in Southwestern Nigeria Soc Sci Med 1997, 44(8): 1195-1207.
14. Abou-Zeid G, Somach SD (2009). Egypt Violence against Women Study: Literature Review of Violence against Women. United States Agency for International Development (USAID) and the Egyptian national Council for Women (ENCW). Available at <http://www.ilo.org/public/english/region/asro/bangkok/newsroom/pr0112.ht>
15. Ekpo UF, Odoemene SN, Mafiana CF, Sam-Wobo SF. Helminthiasis and Hygiene Conditions of Schools in Ikenne, Ogun State, Nigeria. Negl Trop Dis. 2008; 2(1): 146.
16. Adeleye OZ, Nwokoma OG. Non-timber forest products and poverty reduction policy framework in Ikenne Local Government Area, Ogun state Nigeria. IJASS 2012,1(1):1401-1420.
17. Kish, Leslie. 1965. Survey Sampling. New York: John Wiley and Sons, Inc.
18. Enzuladu EA, Agbo HA, Mohammed A, Chia L, Kigbu JH, Zoakah AI. Utilization of modern contraceptives among female traders in Jos South LGA of Plateau State, Nigeria. Int J Med Biomed Res 2012;1:224-231
19. Mitsunaga TM, Powell AM, Heard NJ, Larsen UM. Extramarital Sex Among Nigerian Men Polygyny and Other Risk Factors. J Acquir Immune Defic Syndr. 2005. 39(4): 478-488
20. Slaymaker E, Walker N, Zaba B, Collumbien M. Comparative risk assessment: Unsafe sex . In: Comparative Quantification of Health risks: Global and Regional Burden of Disease due to selected major risk factors. Ezzati M, Lopez, Rodgers A, Murray C (Eds). pp: 1177-1254. Geneva. 2004.
21. Radzah M, Shamsuddin K, Mormi M, Zahari MT, Syimah AN, Asia MN. Early resumption of sexual intercourse and its determinants among postpartum Iban mothers. Int J Reprod Contracept Obstet Gynecol. 2013, 2(2):124-129
22. Awusabo-Asare K, Anarfi JK Postpartum sexual abstinence in the era of AIDS in Ghana: prospects for change. Health Transition Review 1997 Supplement (7): 257-270
23. Borda MR, Winfrey W, McKaig C. Return to sexual activity and modern family planning use in the extended postpartum period: An analysis of findings from seventeen countries. Afr J Reprod Health 2010;14:72-9.
24. Dada O.A, Akesode F.A, Olanrewaju D.M et al. Infant Feeding And Lactational Amenorrhea in Sagamu, Nigeria. Afr J Reprod Health 2002; 6 (2):39-50.
25. Nasir K, Hyder AA. Violence against pregnant women in developing countries; a review of evidence. Eur J Public Health 2003; 105-7
26. Okemgbo CN, Omideyi AK, Odimegwu CO. Prevalence, patterns and correlates of domestic violence in selected Igbo communities of Imo State, Nigeria. Afr J Reprod Health 2002;6:10114.
27. Lawoyin TO, Osinowo H, Walker M. Sexual networking among married men with wives of childbearing age in Ibadan City, Nigeria: report of a pilot study. Afr J Med Med Sci 2004;33: 207-12.
28. Weller S, Davis K (2002) Condom effectiveness in reducing heterosexual HIV transmission. In: *The Cochrane library issue 3*. Update Software, Oxford, <http://cochrane.athens.ac.uk>.