UNDER PEER REVIEW

Assessment of schistosomiasis occurrence and risk factors influencing transmission in different surface irrigation schemes in Morogoro and Kilimanjaro Regions, Tanzania.

ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

Aims: To assess schistosomiasis occurence and identify risk factors influencing its transmission in different surface irrigation schemes in Morogoro and Kilimanjaro Regions.
Study design: Cross-sectional research design was used and Schistosomiasis recorded cases for ten years (January 2002 to December 2011) were collected from health facilities located nearby the irrigation schemes.
Place and Duration of Study: Mkindo, Chabi and Mwega (Morogoro Region) and Kikafu Chini, Lower Moshi and Njoro (Kilimanjaro Region) between October and December, 2011.
Methodology: A total of 378 respondents were selected; 240 in the questionnaire survey, 96 in the Focus Group Discussions and 42 in the Key informant interviews. In each scheme 40 farmers practicing irrigation rice farming were interviewed. Binary logistic regression was used to test association between schistosomiasis infection and a number of predictors such as: age, sex, education, wearing gumboots during irrigation activities, washing clothes and household utensils and cleaning farming tools in canals. Schistosomiasis occurence components such as activities involving water contacts, water management practices and people's behaviour were collected from different schemes through questionnaires and observations. Moreover, tables and figure were used to describe the data.
Results: Although none of the independent variables made a statistical significant contribution to logistic regression model (P>0.05), for Kilimanjaro schemes: age, education, not wearing gumboots during field works, washing clothes, household utensils and farming tools in canals were found to have bigger walds and therefore associated with schistosomiasis infections. Other risks observed were people's behavior of defecating and urinating around the scheme since there were no toilets in the schemes.
Conclusion: The study highlighted the value of assessing schistosomiasis occurrence and risk factors influencing transmission in different surface irrigation schemes in Morogoro and Kilimanjaro for better planning of schistosomiasis control initiatives and proper measures during planning and designing of irrigation schemes.

Keywords: Schistosomiasis Risk Factors, Schistosomiasis Occurance, Kilimanjaro and Morogoro Regions

1. INTRODUCTION

More than 80% of the population in Tanzania is engaged in agricultural activities that contribute about 45% of Gross Domestic Product (GDP) and about 30% of export earnings (1). Agriculture also provides employment to about 75% of the total labour force (1). However, rainfall is unreliable for crop production in many of the areas in Tanzania. Therefore, irrigation is considered the only alternative against drought and erratic rainfall, as well as a strategy to boost production. In Tanzania, about 48% of households use water from rivers for irrigation. According to the National Irrigation Master Plan (NIMP) based on the study by Sosovele et al. (2), out of 43 million hectares of arable land, 29.4 million hectares are suitable for irrigation development. Consequently, there are about 52 modern irrigation schemes (35,900 ha), 982 traditional irrigation schemes (122,600 ha), 113 improved traditional irrigation schemes (25,500 ha) and 42 schemes under water harvesting (197,000 ha), making a total of 381,000 hectares currently under irrigation (2).

Generally, irrigation has the potential to increase agricultural production and consequently improving the economic and social well-being of the rural population. However, irrigation farming necessitates

UNDER PEER REVIEW

an individual to be in constant contact with water and this has been reported to be among the predisposing factors to water-borne diseases such as schistosomiasis and protozoal diseases (3). According to Tanji and Kielen (4), most irrigation systems in Africa, create specific conditions that may favour the establishment of schistosome intermediate hosts and the subsequent transmission of schistosomiasis (4). Several studies have shown that irrigation and other water resource development projects have increased transmission of schistosomiasis and other water-related diseases (5;6;7;8;9;10; and 11).

In Tanzania, two major schistosome species are prevalent, mansoni and haematobium causing intestinal and urogenital schistosomiasis, respectively (12;13). Since late 1970s to early 1980s activities such as the construction of hydroelectric dams and development of irrigation schemes to meet demand for food and water supply altered the transmission pattern of schistosomiasis in the country. Both dams for hydroelectric and irrigation schemes are the risk areas for schistosomiasis transmission as they create a favourable environmental conditions for snail intermediate hosts (14;15;16). Until 2012, Urogenital and intestinal schistosomiasis remained the major public health problem in Tanzania (3). Further, the entire population of approximately 44 million people in Tanzania is at risk of getting schistosomiasis, and the prevalence of the disease appears to increase with the increase in population size. For example, schistosomiasis rose from 19% in 1977 to 51.5% in 2012 (3).

Despite the success of schistosomiasis control programmes in a few countries such as China and Morocco, the disease has remained one of the most serious public health problems, particularly in areas with irrigated agriculture (17). Adults in communities engaging in fishing and irrigation farming, are more exposed to cercariae-infested water than the youth, due to their occupations such as paddy cultivation in swamps or fishing in lakes (18; 19;20).In China and the Philippines respectively it was observed that, age difference in prevalence and infection intensity is influenced by school based control approaches, in which the control intervention is only centred on school children, and the adult population is not included (21;22).

In Tanzania, schistosomiasis and soil-transmitted helminths (STHs) are found countrywide with prevalence ranging between 12.7% and 87.6% for schistosomiasis and up to 100% for STHs (23). Nonetheless many efforts have been made by the Government of Tanzania on schistosomiasis control by providing praziquantel drugs to school children, the group considered to be at a higher risk, but still in some irrigation areas, the disease is considered a problem. Studies on assessment of schistosomiasis prevalence through hospital/dispensary data, water contact activities during irrigation and risk factors that influence schistosomiasis transmission in irrigation areas are inadequate. Previous studies have addressed children behaviour related to water contact such as swimming and playing in water (24). This study therefore intended to assess occurrence of schistosomiasis in the chosen irrigation schemes through health facilities/dispensary data and by identifying risk factors associated with transmission in different surface irrigation schemes in the two regions of Tanzania.

2. MATERIAL AND METHODS

2.1 Study Areas

Six surface irrigation schemes were included in the study. These are Mwega, Mkindo and Chabi located in Morogoro and Lower Moshi, Kikafu Chini and Njoro located in Kilimanjaro. The schemes differ in terms of their designs and construction of infrastructure (Salehe et al., 2012). Farmers from the five schemes ~~that is,~~ three from Morogoro (Mkindo, chabi and mwega) and two from Kilimanjaro (Lower Moshi and Kikafu Chini) use water from rivers as a source of irrigation while farmers in Njoro use water from natural springs originating from Njoro forests. A detailed description of the schemes, climate, topography and information about administrative boundaries of the schemes, the two regions and five districts where the schemes are located can be found in Salehe and Hassan (25) and Salehe et al. (26).

2.2 Study designs and Sampling Procedures

Both cross sectional and longitudinal study designs were used for gathering information. Data were collected between August to October 2012 using quantitative methods through questionnaires and a combination of qualitative methods including observations, key informant interviews and Focus Group

UNDER PEER REVIEW

Discussions (FGDs). A total of 378 participants were involved. 336 participants were chosen randomly in all six schemes from lists of farmers practicing irrigation rice farming, provided by the village government offices and 42 participants were randomly chosen from local pharmacies and dispensaries in the study areas. Out of 378 participants, 240 in all six schemes were included in the questionnaire survey, 40 participants from each scheme. 96 participants in all the schemes were included in 12 FGDS, two FGDs with 16 participants from each scheme, i.e. 8 men in the first group and 8 women in the second group. Men and women groups were formed in order to let women feel free to express themselves, because according to the African culture women are sometimes shy to speak in front of men. Also interviews were conducted with 42 key informants in all the schemes. About 7 key informants in each scheme were interviewed, 4 health personnel from dispensaries and 3 workers of the local pharmacies. Information on activities involving water contact pattern, water management practices, people's habits and behaviours were directly observed in the six irrigation schemes three in Morogoro and three in Kilimanjaro Regions, during October 2011 to October 2012. Moreover schistosomiasis prevalence retrospective data from January 2002 to December 2011 were collected from Mkindo, Malolo and Carmel dispensaries in Morogoro Region and from Pasua, Dr. Nderingo and Mabogini dispensaries in Kilimanjaro Region to compare schistosomiasis prevalence between years in schemes within and between the two Regions.

2.3 Data analysis

Descriptive statistics were presented in tables and graphs to give information on water management practices and diseases, and the mean number of infected people per year in the study areas. Data from the FGDs and key informant interviews were interpreted and summarized into themes based on what was reported by participants. A binary logistic regression model was used to test whether there is any association between schistosomiasis infection and a number of predictors such as: age, sex, education, wearing gumboots during irrigation activities, washing clothes and household utensils in irrigation canals and cleaning farming tools in the irrigation canals.

2.4 Research Permit and Ethical approval

Research permit was given by the Vice Chancellor, Sokoine University of Agriculture (SUA), the Regional Administrative Officers of Morogoro and Kilimanjaro Regions and the District Administrative Officers of Mvomero, Kilosa, Hai and Moshi District. Ethical clearance was obtained from the Medical Research Coordinating Committee of the Tanzania National Institute for Medical Research (NIMR). Verbal consent was obtained from each of the participating individuals after explaining the purpose and importance of the study prior to commencement of interviews. Participation in the study was on a voluntary basis. All the information collected from the participants was kept under the custody of the researcher as confidential and the study participants were anonymized.

3. RESULTS AND DISCUSSION

3.1 Health Status in the studied areas

3.1.1 Diseases, place for treatment and source of domestic water supply

More than 50% of participants interviewed through the questionnaire survey in all the schemes (Morogoro and Kilimanjaro), mentioned schistosomiasis most severe public health problem in their areas followed by malaria (Table 1). All the participants (100%) in Kilimanjaro schemes, more than three quarters of the participants in Mkindo scheme and more than half of the participants in Chabi and Mwega schemes go to hospitals/dispensaries for treatment. All interviewed respondents (100%) in Mkindo, more than half of participants in Lower Moshi (60%) and more than three quarters (86%) in Njoro scheme mentioned that they get domestic water from the taps. In addition, half of the participants in Mwega and about three quarters in Kikafu chini use pumped wells as the source of domestic water supply. Rivers are used as a source of domestic water supply by all farmers in Chabi and by more than a quarter (30%) in Mwega schemes.

Study results, further show that malaria and schistosomiasis are the major diseases affecting the majority of the respondents in all the six schemes. The results also indicate that participants in Kilimanjaro schemes are aware of the importance of attending hospitals in case of illness may be because dispensaries and hospitals are within the scheme areas. However, in the Lower Moshi

UNDER PEER REVIEW

scheme not everyone could have access to clean water because there was a schedule of distributing clean water for domestic use and only those who had high income could have access to clean water as opposed to those with low income who were not ready/could not afford to buy water for home consumption. The only alternative for people with low income was to use water from the irrigation canals for all household activities.

Table 1. Diseases, place for treatment and season experiencing schistosomiasis infections (n=240)

Variables		Schemes					
		Morogoro			Kilimanjaro		
		Mkindo %	Chabi %	Mwega %	Kikafu %	Lower moshi %	Njoro %
Waterborne diseases commonly found in the study areas	Diarrhea	0.0	2.0	2.0	0.0	0.0	0.0
	Schistosomiasis	54.0	84.0	84.0	54.0	84.0	62.0
	Malaria	30.0	6.0	14.0	28.0	12.0	26.0
	Typhoid	12.0	8.0	0.0	18.0	4.0	12.0
	I don't know	0.0	0.0	0.0	0.0	0.0	0.0
	Others	4.0	0.0	0.0	0.0	0.0	0.0
Access to medical facilities	Hospital	87.5	55.0	67.5	100.0	100.0	100.0
	Hospital and local pharmacy	12.5	12.5	22.5	0.0	0.0	0.0
	Hospital and traditional healers	0.0	32.5	7.5	0.0	0.0	0.0
	Local pharmacy	0.0	0.0	2.5	0.0	0.0	0.0
Seasons for schistosomiasis infections	Long rains	52.0	48.0	32.0	34.0	48.0	48.0
	Short rains	6.0	0	0	0	0	0
	Dry season	12.0	14.0	16.0	12.0	24.0	4.0
	Intermediate dry season	8.0	14.0	20.0	8.0	2.0	0.0
	All seasons	8.0	22.0	24.0	10.0	26.0	4.0
	I don't know	14.0	2.0	8.0	36.0	0.0	44.0

3.2 Risk Factors

3.2.1 Toileting during field working hours

Regarding to places for relieving during working hours, majority (75%) of participants during questionnaire survey declared that they relieve around the scheme area while minority (25%) go back home (Table 2). Some of the participants especially in women's FGDs insisted that they just kneel down in the same place where they are working and relieve for the case of urinating and some said that they just make a small hole enough to drop the stool. Participants also insisted by saying that ''it is not possible for someone to go back home in case one wants to relieve during working hours, every one of us relieve in the scheme area.'' Moreover, during FGDs farmers were asked if toilets were to be built in the scheme areas would they make use of them. Participants especially in male Focus Group Discussions in Lower Moshi, Chabi, Mwega and Kikafu Chini schemes said they would make use of the toilets, while in the female Focus Group Discussions in Chabi, Mwega and Kikafu Chini

UNDER PEER REVIEW

said that it would not help because people's behaviour are different, some farmers would still not make use of the toilets.

The results show that lack of toilets and people's behaviour leads to schistosomiasis transmission in the studied areas. The result also implies that farmers have to be sensitized continuously on the importance of using toilets and how schistosomiasis can be controlled if toilets are to be built and utilized effectively. The findings are in line with the study done in Zimbabwe (27) who reported that, construction of toilets in a way that every farmer in the field gets closer to it helped in minimizing schistosomiasis transmission.

3.2.2 Schedule for canal maintenance (cleaning)

Table 2 indicates that more than 75% of the participants in Mkindo, Mwega and Kikafu Chini schemes clean irrigation canals once in a month, while more than half of the participants in Chabi do clean canals twice in a month. However, half of the participants in Lower Moshi and more than quarter of the participants in Njoro schemes declared to have been removing vegetation in canals once in two months. When farmer were asked during Focus Group Discussions whether they understood why was it important for them to remove vegetation in canals and drain excess water in the field, their response are in two categories. First majority of them gave two reasons, which are: (i) they remove vegetation in order to allow easy flow of water from canals to their paddy plots and, (ii) they remove excess water to let their crops dry during harvesting. The second view, which comprise the minority (from Mkindo and Lower Moshi schemes), related the removal of vegetation in canals for control of diseases.

Results implies that participants in the two schemes, Morogoro improved traditional scheme (Mkindo) and Kilimanjaro modern scheme (Lower Moshi), were aware of the reason of removing vegetation probably because they received training from Mkindo Farmers Center and Kilimanjaro Agriculture Training Centre respectively.

3.2.3 Risk factors and likelihood of having schistosomiasis infection: comparison for schemes in the two regions

Direct logistic regression was performed to assess the impact of a number of factors on the likelihood of respondents in all schemes studied to report that they had schistosomiasis infection. The model contained six independent variables (sex, age, education, cleaning farming tools in canals, wash clothes and household utensils in canals and wear gumboots during irrigation farming activities). The full model containing all the predictors was not statistically significant for the schemes in the two regions.

Table 2: Risk Factors Influencing Schistosomiasis Transmission

Variables		Schemes					
		Morogoro			Kilimanjaro		
		Mkindo %	Chabi %	Mwega %	Kikafu %	Lower moshi %	Njoro %
A place for relieving during field work hours	Go back home	16.0	18.0	10.0	18.0	20.0	34.0
	Around the scheme area	84.0	82.0	90.0	82.0	80.0	66.0
	I don't know	14.0	2.0	8.0	36.0	0	44.0
Schedules for canal cleaning	Once a month	82.5	46.2	100.0	77.5	32.5	72.5
	Twice a month	15.0	53.8	0.0	22.5	17.5	27.5
	Once in two months	2.5	0.0	0.0	0.0	50.0	82.5

UNDER PEER REVIEW

Sources of domestic water supply	Irrigation canal	0.0	28.0	20.0	0.0	34.0	0.0
	Tap	100.0	0.0	0.0	10.0	60.0	86.0
	Pumped well	0.0	0.0	50.0	78.0	6.0	0.0
	Rivers	0.0	100.0	30.0	12.0	0.0	0.0
	Springs	0.0	0.0	0.0	0.0	0.0	14.0

3.2.4 Binary logistic regression outputs and the odds of being infected by schistosomiasis

Case processing summary was one of the important outputs of binary logistic regression model (Table 3a). In the tables it shows that 100% were included in the analysis in each Region representing the whole samples of 120 of respondents interviewed in each region. There were no missing cases in both Regions.

Table 3a. Case processing summary

Schemes in Morogoro

Un-weighted cases (n = 120)		%
Selected Cases	Included in analysis	100
	Missing cases	0
	Total	100
Unselected cases		0

Schemes in Kilimanjaro

Un-weighted cases (n = 120)		%
Selected Cases	Included in analysis	100
	Missing cases	0
	Total	100
Unselected cases		0

Table 3a. Case processing summary

Schemes in Morogoro

Un-weighted cases (n = 120)		%
Selected Cases	Included in analysis	100
	Missing cases	0
	Total	100
Unselected cases		0

UNDER PEER REVIEW

Schemes in Kilimanjaro

Un-weighted cases (n = 120)		%
Selected Cases	Included in analysis	100
	Missing cases	0
	Total	100
Unselected cases		0

The Omnibus test is another output of binary logistic regression model. It tests for the capability of all predictors (independent variables) in the model jointly to predict the response (dependent) variable. From the output results as seen in table 3b it shows that there were no significant differences in all the data entered into the model in the two Regions.

Table 3b. Omnibus test of model coefficients

Schemes in Morogoro Region

		Chi-square	Df	p-value.
Step 1	Step	9.055	6	0.171
	Block	9.055	6	0.171
	Model	9.055	6	0.171

Schemes in Kilimanjaro Region

		Chi-square	Df	p-value.
Step 1	Step	3.407	6	0.756
	Block	3.407	6	0.756
	Model	3.407	6	0.756

Moreover, the model summary, which is presented in Table 3c showing Cox & Snell R square and Nagelkerke R square, was chosen as an important output of the binary logistic regression model. The Cox-Snell R2 and Nagelkerke R2 are attempts to provide a logistic analogy to R2 in OLS regression; hence are called pseudo R2. Nagelkerke R2 is a modification of Cox-Snell R2 to assure that Cox-Snell R2 varies from zero to one, as does R2 in OLS regression. If Cox-Snell R2 is not modified, its maximum value is usually less than 1, making it difficult to interpret.

Table 3c. Model summary

Schemes in Morogoro Region

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	147.534	0.073	0.100

Schemes in Kilimanjaro Region

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	160.238	0.028	0.038

UNDER PEER REVIEW

Nagelkerke R2 is normally higher than Cox-Snell R2 and is the most-reported of the pseudo R2 estimates. Therefore, based on the results in Table 3d which show that Nagelkerke R2 was 0.100 and 0.038 in Morogoro and Kilimanjaro Regions respectively, it means that the independent variables entered in the model explained 10 and 3.8% of variance in the dependent variable for schemes in Morogoro and Kilimanjaro respectively.

Hosmer and Lemeshow Test is another output of logistic regression model which tests the goodness-of-fit of a logistic regression model. In this study, the value of the Hosmer and Lemeshow chi-square obtained was 7.193 for schemes in Morogoro and 3.783 for schemes in Kilimanjaro Region, and it was not significant (p= 0.576 and p=0.876) for schemes in Morogoro and Kilimanjaro respectively, as observed in Table 3c. Normally, where the Hosmer and Lemeshow chi-square value is greater than 0.05, it represents the goodness of fit of the model. Therefore it implies that the model's estimate which was used in this study which contained six explanatory variables in each region fitted the data at an acceptable level.

Table 3d. Hosmer and Lemeshow Test

Schemes in Morogoro Region

Step	Chi-square	Df	p-value
1	7.193	8	0.576

Schemes in Kilimanjaro Region

Step	Chi-square	Df	p-value
1	3.783	8	0.876

Wald statistics is another useful output. Wald coefficients associated with individual independent variables help us realize the relative importance of each independent variable. A bigger Wald statistic implies that the independent variable associated with it has high contribution to the occurrence of the dependent variable. In this case Education in both Morogoro and Kilimanjaro schemes highly contribute to occurrence of the dependent variable since they have bigger Walds.

In a B coefficient, a negative sign shows that, that particular variable decreases the logit of the dependent variable (i.e. it decreases the probability of that event (in this case schistosomiasis infection) will be realized, and vice versa. For example in Table 3e: age, sex and the activity of washing clothes and household utensils in canals in Morogoro schemes; and sex, education and the activity of cleaning farming tools in irrigation canals in Kilimanjaro schemes reduced the chances of farmers to be infected by schistosomiasis since their B values are associated with negative signs. In other words, the other variables increase chances of farmers to be infected since they have positive signs. Therefore in Morogoro schemes education, the activity of bathing in irrigation canals and the behaviour of not wearing gumboots during field works in paddy plots had impact on schistosomiasis infection.

Odds Ratio

The odds ratio is the predicted change in the odds for a unit increase in the corresponding independent variable. Odds ratios less than 1 correspond to decreases in the odds whereas odds ratios more than 1.0 correspond to increases in the odds. An odds ratio equals to 1.0 means that the respective independent variable has no effect on the dependent variable whereas an odds ratio close to 1.0 means that the respective independent variable almost has no effect on the dependent variable. Based on the above Exp(B) value for farmers infected by schistosomiasis, results of the model explains that important variables that contributed positively to schistosomiasis infection were education, bathing in canals and not wearing gumboots and age for Morogoro schemes, and washing of clothes and household utensils and wearing gumboots for Kilimanjaro schemes. The magnitude of

UNDER PEER REVIEW

effects of other independent variables on the dependent variable (schistosomiasis infection) is shown in Table 3e. In Morogoro schemes farmers with formal education were more likely to avoid activities that could have exposed them to infections about four times than those with no formal education (illiterates). Also farmers who wore gumboots and those who did not bath in canals were likely to avoid infections one and two times respectively, than those who did not wear gumboots and those who bath in canals.

3.2.5 Risk factors identified during observations

In Mwega scheme, people of different age groups were observed taking bath in the canals. Since the scheme area is close to residential areas, all activities that involve water are done in the irrigation canals (see plate 1 and 2). It was observed that some people especially those that reside near to the canals (4 to 15 m away) were seen in the morning hanging towels on their shoulders, holding tooth brushes and soap in their hands going to the canal for bathing. The same activity of washing and bathing directly in the canals was observed in the evening where groups of men on one side of the main canal and other groups of women on the other side of the main canal taking a bath half naked (for women) and naked (for men). In addition, to the above observations, improper cropping pattern were observed in the plots. One plot was found with paddy while the next plot had maize, beans or onions. This result imply that there is improper water management practice carried out in this plots since all the crops mentioned above have different water demands and therefore there will be standing pools of water.

UNDER PEER REVIEW

Table 3e: Likelihood of Reporting Schistosomiasis Infection by Region

Variables	B	S.E.	Wald	Df	P-value	Exp (B)	95.0% C.I For Odds Ratio Lower	Upper
Morogoro Schemes								
Age	0.024	0.016	2.245	1	0.134	0.976	0.945	1.008
Sex	-0.582	0.398	2.139	1	0.144	0.559	0.252	1.219
Education	1.278	0.891	2.057	1	0.152	3.591	0.626	20.606
Wash clothes house-hold utensils in canals	-0.32	0.416	0.600	1	0.438	0.725	0.321	1.637
Bath in canals	0.277	0.401	0.477	1	0.490	1.320	0.601	2.899
Wear gumboots during irrigation activities	0.515	0.721	0.511	1	0.475	1.673	0.408	6.869
Constant	1.441	0.976	2.179	1	0.140	4.226		
Kilimanjaro schemes								
Age	0.009	0.017	0.274	1	0.601	1.009	0.976	1.043
Sex	-0.305	0.405	0.564	1	0.452	0.737	0.333	1.632
Education	-1.268	1.175	1.164	1	0.281	0.281	0.028	2.817
Wash clothes and house-hold utensils in canals	0.308	0.406	0.574	1	0.449	1.360	0.614	3.016
Cleaning farming tools in irrigation canals	-0.338	0.419	0.650	1	0.420	0.714	0.314	1.621
Wear gumboots during irrigation activities	0.137	0.414	0.110	1	0.740	1.147	0.509	2.584
Constant	-0.523	0.920	0.322	1	0.570	0.593		

P is significant at 0.05 level (two tailed test)

UNDER PEER REVIEW

Plate 1: A child swimming in the canal

Plate 2: A Woman washing near the main canal

Even though farmers in Mkindo had access to clean water from taps, still a number of people infected with schistosomiasis from the dispensary data were high compared to other schemes. The researcher was therefore interested to know why the number of infected people in Mkindo was so high for the ten years in which data were collected compared to other scheme areas, and therefore Mkindo as a case to be investigated. Risk factors that were observed in Mkindo irrigation scheme were as follows: (i) standing pools of water almost for the whole scheme area; (ii) the main irrigation canal and the main drainage canal were full of water and vegetation, (iii) the drainage canal is also used as irrigation canal, (iv) the division boxes (structures) were also chocked with vegetation and snails were found sticking on the vegetation; (v) the drainage canals were not serving the purpose of removing excess water from the plots since the plots were found with standing water even during harvesting period (see plate 3 and 4).

UNDER PEER REVIEW

Plate 3: Plots filled with water during harvesting period

Plate 4: Canals chocked with vegetation

Another observation took place in Chabi scheme. It was observed that: (i) the village did not have a domestic water supply system, therefore water for domestic purposes was collected from the canals and Chabi river, (ii) there was a very big swamp called ''tete'' in respondents local language. The swamp stores water used for irrigation purposes and it was chocked with aquatic weeds (Plate 5).

Observations from Lower Moshi scheme were as follows: (i) people were found washing, swimming especially children and collect water for domestic use in canals; (ii) some farmers were found drinking water from the canals during field working hours; (iii) some were washing their limbs and farming tools in the canals after farm work.

Plate 5: A water storage structure full of vegetation growth

The following were considered as the main causes and risk factors of schistosomiasis transmission during observations in Kikafu Chini scheme: (i) some of the earth canals are constructed near residential houses, (ii) the canals used to irrigate bananas, vegetables and coffee and people do wash clothes and bath in the canals; (iii) pit latrines constructed close to irrigation canals at the middle stream; (iv) children do bath close to these canals and the downstream dwellers do fetch water for domestic use.

UNDER PEER REVIEW

Plate 6: Pit latrine built close to a canal

Plate 7: people washing clothes and households in a canal

The same risks were observed in Njoro scheme where; people were found fetching water for washing and bathing from the irrigation canals. In addition children were swimming, and farmers were found washing their limbs and farming tools after field work in the canals.

These results suggest that social activities such as washing household utensils, bathing, swimming, collecting water for household use and washing of limbs and farming tools in irrigation canals may have contributed to exposing farming communities to schistosomiasis infections and transmission in the study areas. These results concur with what has been reported in either studies (28;29;30). Moreover the results further shows that lack of toilets in the studied schemes and excessive vegetation growth in irrigation and drainage canals have also lead to prominence of schistosomiasis infection. These results are supported by Tsafack (31) and Hilali et al. (32). Very few farmers particularly men in the four schemes (Mkindo, Kikavu Chini, Lower Moshi and Njoro) were observed wearing rubber/gum boots while working in the water. This also indicates that farmers are exposed to schistosomiasis infections during working hours of the day.

None of the independent variable tested made a significant statistical contribution to the model when testing association between independent variables and reporting schistosomiasis infection. However, wald statistics showed that education had significant contribution to occurrence of schistosomiasis infection in the two regions. Other variables which contributed positively to schistosomiasis infections were bathing in irrigation canals, behaviour of not wearing gumboots during field works in paddy plots and washing clothes and household utensils and farming tools in canals.

3.3 Schistosomiasis Occurrence

3.3.1 Information on schistosomiasis at community level

During interviews with key informants (Health personnel and local pharmacists) from Mabogini and Carmel dispensaries, it was revealed that the number of people visiting dispensaries and pharmacies for schistosomiasis clinical test and treatment has been declining compared to the past years (from year 2009 to 2011). However, key informants in Mkindo still believe that the number of people infected by schistosomiasis per year is still high. Also schistosomiasis recorded data from dispensaries in the studied areas (Mabogini, Carmel mission, Malolo and Pasua dispensaries) reveals the same (Figure 1).

UNDER PEER REVIEW

Further discussions with the key informants in all the schemes, shows that people of all ages are equally affected by schistosomiasis. In addition, when the key informants were asked why children are equally affected like adult farmers while they spent most of their time in schools, the key informants in the five schemes with exception of those in Mkindo scheme answered that children do play and swim in irrigation canals after school hours. On the contrary informants in Mkindo pointed out that children play in the standing pools of water during the rainy season and some accompany their parents to work in the rice fields during weekends and holidays.

Regarding the question on whether schistosomiasis infected people attend treatment in the early stages or late stages of the disease the key informants pointed out that some do visit the dispensaries for clinical tests and get treatment from the dispensary whenever they themselves or their children saw symptoms related to schistosomiasis.However, others just visit the local pharmacy complaining that they have signs of schistosomiasis without prescription from the doctor. The local pharmacists in Mkindo village insisted that most of those visiting the pharmacy without doctor's prescription were male adults who shy off visiting dispensaries for clinical tests because they relate signs of blood in urine with other sexually transmitted diseases (STDs).

Responding on why schistosomiasis is still a health problem in the study areas the key informants in Mwega, Chabi, Kikafu Chini and Lower Moshi claimed that the majority of people in these areas depend on water from the irrigation canals for their domestic uses (drinking, cooking, washing, bathing and for animals). Informants in Mwega scheme further argued that worse enough some people do clean cooking pots and wash clothes in canals and some bath in the canals and faeces have been seen several times floating in canals indicating that some people use the canals as toilets. Key informants in Kikafu Chini and Mwega added that parents leave their children to play in the canals. Moreover, informants in Kikafu Chini maintained that the canals are not regularly cleaned and some households drain dirty water from toilets into the canals.

3.3.2 Schistosomiasis occurrence in the study area

Study results further show that February, May, June and December constituted the highest number of people infected by schistosomiasis in scheme areas in Morogoro Region while January through May also maintained higher level of infection in scheme areas in Kilimanjaro Region (Fig. 1). The pattern for Kilimanjaro was irregular in the period from June to December. It was also evident that in all months across ten years (2002-2011) schistosomiasis recorded cases were higher in Morogoro health centers compared to Kilimanjaro, meaning that schistosomiasis reported cases were higher in the Morogoro schemes while the cases were least in Kilimanjaro schemes.

According to the secondary data for a ten year period (2002-2011), there were a total of 4272 schistosomiasis patients who received treatment from health facilities in/near Mkindo, Malolo and Chabi schemes in Morogoro while 1778 patients were diagnosed to have schistosomiasis infections in the Mabogini, Pasua and Dr.Kisanga's health facilities in Kilimanjaro (Table 4). Compartively, Mkindo health center (in Morogoro) had the highest number of recorded schistosomiasis cases followed by Mabogini health center (in Kilimanjaro) while Pasua health center (also in Kilimajaro) had the least cases of the disease (Table 4). In addition, there were more recorded patients in year 2007 in all six health facilities compared to other years.

UNDER PEER REVIEW



Figure 1. Mean monthly occurence of schistosomiasis recorded cases from different health centres in Morogoro and Kilimmanjaro across ten- year period from 2002 to 2011.

Only data for 2010 and 2011 were accessed in Kikafu Chini from Dr. Kisanga's dispensary, simply because there were no records of patients visiting the dispensary. Therefore, the researcher adviced the Doctor to start keeping records for schistosomiasis cases from the year 2010 to 2011. In addition, FGDs were conducted with some farmers to get information on schistosomiasis prevalence. From the FGDs, it was revealed that apart from Dr Kisanga's dispensary people used to get treatment from Dr Mbonea's dispensary which is located 5 kms away from Kikafu Chini village.

The result implies that, the number of infected people by schistosomiasis in Malolo, Carmel, Pasua and Mabogini dispensaries decreased from the year 2010 to 2011 probably due to health education people receive from health personnel from the above mentioned health facilities. In addition, the reason as to why people living in Mkindo village had high number of schistosomiasis cases in all the ten years data collected may be contributed by poor drainage system and vegetation growth in the conveyance systems which encouraged snails breeding. According to Salehe et al. (33), poor knowledge of schistosomiasis is another reason which causes increase in schistosomiasis transmission in Mkindo scheme as only 45% of respondents in the scheme had better knowledge on schistosomiasis predisposing factors. From this result, it indicates that majority of farmers in Mkindo scheme are not aware of the risk factors influencing schistosomiasis transmission. Moreover, the presence of a swamp chocked with vegetation in Morogoro traditional scheme (Chabi) may have contributed to schistosomiasis transmission as snails breed and feed on weeds.

UNDER PEER REVIEW

Table 4: Number of people infected by schistosomiasis per year in the six health facilities in Morogoro and Kilimanjaro Regions

Region	Health facilities in the scheme areas	Total number of schistosomiasis cases in different years										
		2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	Total
Morogoro	Mkindo	189	0	0	440	730	635	0	800	298	276	3368
	Malolo	0	0	0	0	42	50	31	29	0	0	152
	Carmel	0	98	0	68	0	347	78	123	34	4	752
	Total	189	98	0	508	772	1032	109	952	332	280	4272
Kilimanjaro	Mabogini	268	0	242	274	0	179	141	67	64	49	1284
	Pasua	112	84	33	8	25	79	23	48	29	8	449
	Dr. Kisanga	0	0	0	0	0	0	0	0	29	16	45
	Total	380	84	275	282	25	258	164	115	122	73	1778

4. CONCLUSION AND RECOMMENDATIONS

4.1 CONCLUSIONS

From the above results the study concludes that, occurrence of schistosomiasis decreased from year 2008 in Mabogini, Carmel, and Malolo dispensaries while the number of cases was still high in Mkindo dispensary. Schistosomiasis mass treatment programmes which were provided to primary school children since year 2004 might have helped in minimizing the number of cases.

Major risk factors observed were people's behaviour of swimming, bathing and washing of clothes and farming tools in irrigation canals. Other risks observed were: limited availability of toilets which enhanced schistosomiasis transmission, poor maintenance of irrigation and drainage canals to remove vegetation and snails were found attached to vegetation covers in the canals in all studied schemes.

4.2 RECOMMENDATIONS

From the study it is recommended that the Government through the Ministry of Health and Social Welfare and the National Institute for Medical Research to provide praziquantel tablets to irrigation farming communities at least once in every year. This should be done through mass treatment programmes. The more farmers receive treatment in irrigated areas, the lesser transmission of the disease and the fewer the snails will be infested by the parasites.

The study also recommends that irrigation department through the local government at District level should: i) keep on creating awareness to farmers about schistosomiasis, its effects, transmission and the way people can protect themselves from infections. ii)provide education to farmers on the necessity of cleaning canals to remove vegetation cover and proper draining of excess water in the fields and canals; iii) sensitize farmers to raise funds for construction of simple affordable toilets and educate them on the importance of using toilets.

The study further recommends that the Ministry of Agriculture, Livestock and Fisheries Development through the department of irrigation and technical services shoud do the following:
(i) Integrate irrigation and health related issues during farmers training.

(ii) Include budget for construction of toilets and clean water for domestic use during planning and designing of irrigation projects.

UNDER PEER REVIEW

REFERENCES

1. United Republic of Tanzania.The National Irrigation Policy Draft. Ministry of Water and Irrigation. Dar es Salaam, Tanzania. 70pp; 2009.

2. Sosovele, H., Boesen, J. and Maganga, F. Social and Environmental Impact of Irrigation Farming in Tanzania, Selected Cases. Dar es Salaam University Press Ltd, Dar es Salaam, Tanzania. 184 pp; 2005.

3. Mazigo, H.D., Nuwaha, F., Kinung'hi, S.M., Morona, D., Moira, A.P., Wilson, S.,Heukelbach, J. And Dunne, D.W. Epidemiology and control of human schistosomiasis in Tanzania. Parasites and Vectors. 2012; 5: 274.

4. Tanji, K.K. and Kielen, N.C. Agricultural drainage water management in arid and semi-arid areas. Food and Agriculture Organisations of the United Nations (FAO) Rome, Italy. 202pp; 2002.

5. Deacon, N.H.G. A review of the literature on schistosomiasis and its control. Technical note OD/TN3. Hydraulics Research. Wallingford, United Kingdom. 16pp; 1983.

6. Oomen, J.M.V., De Wolf, J. and Jobin, W.R. Health and irrigation. Incorporation of disease control measures in irrigation, a multi faceted task in design, construction, operation. International Institute for Land Reclamation and Improvement.Wageningen,The Netherlands. 119 pp; 1988.

7. Talla, I., Kongs, A., Verle, P., Belot, J., Sarr, S. and Coll, A.M. Outbreak of intestinal schistosomiasis in the Senegal River Basin. Annals of the Belgian Society of Tropical Medicine. 1990; 70 (3): 173-180.

8. Hunter, J.M., Rey, L., Chu, K.Y., Adekolu-John, E.O. and Mott, K.E. Parasitic diseases in water resources development. The need for intersectoral negotiation. Publications of the World Health Organization, Geneva, Switzerland. 13pp;1993.

9. Grosse, S. Schistosomiasis and Water Resources Development: A Re-evaluation of an Important Environment-Health Linkage. University of Michigan Press United States of America. 32pp;1993.

10.Steele, P., Konradsen, F. and Imbulana, K.A.U.S. Irrigation, health, and the environment: A literature review with examples from Sri Lanka. International Irrigation Management Institute (IIMI). Colombo, Sri Lanka. 29 pp; 1997.

11. Ofoezie, I.E. Human health and sustainable water resources development in Nigeria: Schistosomiasis in artificial lakes. Natural Resources Forum. 2002; 26: 150-160.

12. Ross, A.G.P., Li, Y.S., Sleigh, A.C., Li, Y., Williams, G.M., Wu, W.Z., Lou, X., He, Y. and McManus, D.PEpidemiologic features of Schistosoma japonicum among fishermen and other occupational groups in the Dongting Lake region. (Human Province) of China. American Journal of Tropical Medicine and Hygiene. 1997; 57(3): 302-308.

13. Steinmann, P., Keiser, J., Bos, R., Tanner, M. and Utzinger, J.Schistosomiasis and water resources development: systematic review,meta-analysis,and estimates of people at risk. Lancet Infectious Disease. 2006; 6: 411–25.

14. Sturrock, R.F. The development of irrigation and its influence on the transmission of bilharziasis in Tanganyika. Bulletin of the World Health Organisation. 1965;32: 225- 236.

15. McCullough, F.S., Eyakuze, V.M., Msinde, J. and Nditi, H. Water resources and bilharziasis transmission in the Misungwi area, Mwanza district, north-west Tanzania. East Africa Medical Journal. 1996; 45(5): 295-308.

16. Matovu, D.B. and Nditi, H.P. The distribution and intensity of schistosomiasis in Tanzania mainland. Proceedings of the Annual Scientific Conference of the Medical Association of Tanzania. Dar Es Salaam, Tanzania. 12 pp; 1980.

17. Engels, D., Chitsulo, L., Montresor, A. and Saviolli, L. The global epidemiological situation of schistosomiasis and new approaches to control and research. Acta Tropica. 2002; 82(2): 139-146.

UNDER PEER REVIEW

18. Rugemalila, J.B. The impact of urinary schistosomiasis on the health of two community populations living in endemic areas in Tanzania.Tropical and Geographical Medicine. 1979; 31(3): 375-380.

19. Jordan, P. and Webbe, G. Schistosomiasis. Epidemiology, Treatment and Control.Heinemann Medical Books Ltd, London, United Kingdom.415 pp; 1982.

20. Lwambo, N.J.S. Transmission of urinary schistosomiasis in Sukumaland, Tanzania. 1. Snail infection rates and incidence of infection in school children. Journal of Helminthology. 1988; 62: 213-217.

21. Olveda, R.M., Tiu, E., Fevidal, P. Jr., de Veyra, F., I catlo, F. C. Jr. and Domingo, E.O. Relationship of prevalence and intensity of infection to morbidity in schistosomiasis japonica: a study of three communities in Leyte, Philippines. American Journal of Tropical Medicine and Hygiene. 1983; 32:1312-1321.

22. Ross, A. G., Bartley, P.B., Sleigh, A.C., Olds, G.R., Williams, G.M. and McManus, D.P. Schistosomiasis. N Engel Journal of Medical. 2002; 346:1212-1220.

23. Lwambo, N.J., Siza J.E., Brooker, S., Bundy, D.A. and Guyatt, H. Patterns of concurrent hookworm infection and schistosomiasis in schoolchildren in Tanzania. Transactions of Royal Society of Tropical Medicine and Hygiene. 1999; 93(5): 497- 502.

24. Ndyomugyenyi, R., and Minjas, J.N. Urinary schistosomiasis in schoolchildren in Dar-es-Salaam, Tanzania, and the factors influencing its transmission. Annals of tropical medicine and parasitology. 2001; 95: 697-706.

25. Salehe, F.S and Hassan, S.N. Socio-economic Effects of Schistosomiasis on Irrigation Rice Growers in Morogoro, Tanzania. American Journal of Experimental Agriculture. 2012; 2(3): 395-406

26. Salehe, F.S., Mtambo, M.M.A., Tarimo, A.K.P.R., Mattee, A.Z and Hassan, S.N. Schistosomiasis: A comparative study of its socio- economic consequences in different types of irrigation schemes in Tanzania.Global Advanced Research Journal of Agricultural Science. 2012); 1(8): 240-249.

.

27.Chimbari, M.J. Enhancing Schistosomiasis Control Strategy for Zimbabwe: Building on Past Experiences. Journal of Parasitology Research. 2012; 93(8): 1795-1801.

28. Van Koppen, B., Moriarty, P. and Boelee, E. Multiple-use water services to advance the millennium development goals. Research Report 98. International Water Management Institute. Colombo, Sri Lanka. 53pp; 2006.

29. Jones, C.R. WHO Report on the Health Component in the Volta Lake Project. Unpublished World Health Organisation document AFR/PHA/1 15: AFR/SCHIST/27. Accra, Ghana. 9 pp;1973.

30. Nicholas, M., Sekesai, M., Munyaradzi, P. M., Noah, H. P, Davison, S., Gibson,H., Masceline, J. M., Farisai T., Godfrey, M., Vivian, C., Kimberly, C. B., Francisca, M., Nirbhay, K. and Takafira, M. Knowledge attitudes and practices of grad three primary school-children in relation to schistosomiasis, soil transmitted helminthiasis and malaria in Zimbabwe. BMC Infectious Diseases. 2011;11: 169.

31. Tsafack, F.Ecology of schistosome intermediate hosts in the Benue Upper-Valley area. Public Health. 1997; 12: 46-55.

32. Hilali, A.M.H., Dessouqui, L.A., Wassila, M., Daffalla, A.A. and Fenwick, A.Snails and aquatic vegetation in Gezira irrigation canals. Journal of Tropical Medicine and Hygiene. 1985; 88: 75–81.

UNDER PEER REVIEW

33. Salehe, F.S., Mattee, A.Z., Tarimo, A.K.P.R., Ensink, J.H.J. and Mtambo, M.M.A. Differences in Schistosomiasis Knowledge among Irrigation Rice Farming Communities in Different Irrigation Schemes in Tanzania. International Journal of TROPICAL DISEASE and Health. 2013; 3 (1): 13-24.

UNDER PEER REVIEW