

Clinical Practice Articles

Low Body Mass Index Helps Detect HIV Infection in Uganda

ABSTRACT

Background: The HIV infection has been and is still a big issue in the world and especially in sub-Saharan Africa with UN statistics showing that 65% of the people infected with HIV. Although the Ugandan Ministry of Health (MOH) has implemented the Nutritional Education Program at hospital level, the actual nutritional condition of HIV positive people has not documented yet. The context prompted us to execute this investigation to examine the nutritional condition and the eating habits of HIV positive people compared to HIV negative people in Uganda.

Place and Duration of Study: Kampala, Busia, Mbende, Mbende General Hospital HIV Clinic, and Masafu General Hospital HIV Clinic in Uganda. 24th August 2013 and 2nd September 2013.

Methods: Eighty HIV positive patients visited two hospitals, and one hundred and eleven individuals who self-reported as HIV negative in Kampala, Mbende and Busia. Participant's weight, height and body size were measured using an electric weight scale, and a measuring tape were used to calculate BMI using standard equations. After calculating BMI scores for low BMI were filtered according to the WHO criteria (BMI<18.5). BMI scores were categorized by gender.

Results: There was a significant difference in BMI mean values between HIV positive and negative people. Although HIV positive and negative subjects consumed a similar amount of food every day, the mean BMI of HIV positive people is lower than that of HIV negative people. We saw similar results for all age groups.

Conclusion: HIV infection in the non-symptomatic carrier stage may influence the nutritional condition and eating habits for people in Uganda. Therefore, BMI measurement could be a very useful method for assessing chronic malnutrition.

Keywords: HIV; AIDS; BMI; Nutritional condition

1. INTRODUCTION

The purpose of this study is to prove to the Uganda government that HIV positive individuals show significantly lower BMI compared to HIV negative individuals. This means there is a cost-effective method to help hospital staff to identify HIV positive individuals faster. According to the most recent study of World Health Organization (WHO), 36.7 million people were living with HIV by the end of 2016 world-wide, and almost two-thirds of them are residing in Sub-Saharan Africa [1].

Uganda is the one of the countries in Sub-Saharan Africa that has been severely affected by HIV. Although the Ugandan government has been conducting several anti-HIV measures

including a large-scaled HIV awareness campaign in early 2000's [2], the domestic HIV infection rate has increased in recent years [3]. The reasons for the increased infection rates in the area are due to a lack of public knowledge regarding preventative measures [4]. The other issue is that the HIV detection rate is very slow, because Uganda citizens do not have access to affordable HIV test. If affordable testing methods for HIV becomes available the HIV infection could be detected early, and infection prevention could be promoted in Uganda.

Low Body Mass Index (BMI) and body weight loss are known to be directly related to the CD4 (Antibody) reduction [5], and development of some diseases associate with HIV, such as tuberculosis (TB) [6, 7].

BMI is easily measured and calculated with minimum tools and knowledge; therefore, we conducted a research to find a relationship between BMI and HIV status in Uganda, based on an assumption that BMI of HIV carrier is lower than non-HIV carriers. If any relationship between those is confirmed, BMI could be an affordable and beneficial method to identify the possibility of an HIV infection.

Since a meal frequency links to weight gain and loss, we also studied a relationship between a meal frequency in HIV positive and negative [8].

2. STUDY POPULATION AND METHODS

2.1 Study Subjects

Eighty HIV positive patients visited two hospitals in Mbende and Busia, and one hundred and eleven individuals who self-reported as HIV negative in Kampala, Mbende and Busia. All HIV positive participants were diagnosed based on the result of blood test conducted by those hospitals prior to this study. Only HIV positive individuals who presented no symptoms of AIDS and AIDS-related diseases World Health Organization (WHO) defines participated in this study. Subjects were enrolled if they were 18 years old or older, had an average income level (US \$50-\$100) and no dietary restriction for religion. These included both males and females.

2.2 Informed Consent & Confidentiality

All subjects were informed in advance the purpose of this study, and participation in the study was voluntary. Anonymity, confidentiality and privacy of data were also explained and guaranteed. Written consent was required by all applicants.

2.3 BMI Assessment

Participant's weight, height and body size were measured using an electric weight scale, and a measuring tape brought from Japan were used to calculate BMI using standard equations. After calculating BMI scores for low BMI were filtered according to the WHO criteria (BMI<18.5). BMI scores were categorized by gender.

2.4 Lifestyle Factors

A standard and structured questionnaire was used to collect information on demographics and lifestyle factors. The data collection period was 24th August 2013 to 2nd September 2013. The questionnaire prepared both in English and a local language spoken in the region. Content of the questionnaire includes participants' gender, age, occupation, HIV status, a chance of pregnancy, family structure, family history of HIV disease, smoking and drinking habits, medical history and daily meal frequency. Daily meal frequency data was classified

into 3 categories; 'twice or less,' '3 times' and '4 times or more,' and then compared with participant's HIV status.

2.5 Statistical Method

To validate the hypotheses on relationships between BIM and HIV status, the Student's t-test was used. Furthermore, multiple regression analysis adjusted confounders including age, medical history smoking and drinking habits, a chance of pregnancy and family history of HIV disease was applied. For examination of meal frequency and HIV status, the chi-square test and the Fisher's exact test were employed. All statistical analysis was conducted with the level of significance set at 0.05.

3. RESULTS

There were 80 HIV-positive individuals (39 males and 41 females), and 111 HIV-negative individuals (61 males and 50 females) enrolled in this study.

Table 1 shows the basic characteristics of the participants that were analyzed with a comparison between HIV-positive and HIV-negative group. BMI was significantly lower in the HIV-positive than HIV negative participants, both males ($p=0.007$) and females ($p=0.02$). This difference remained significant in a multivariate analysis including age and other factors such as medical history smoking and drinking habits, a chance of pregnancy and family history of HIV disease.

The relative frequency distribution of HIV status and gender is shown in Table 2. Participants in low BMI (<18.5) were 7 males with HIV positive (17.9%), 3 males with HIV negative (5.0%), 8 females with HIV positive (19.5%), and 1 female with HIV negative (2.0%).

The results of linear regression analysis using BMI and other explanatory variables such as age, HIV status, medical history smoking and drinking habits, a chance of pregnancy and family history of HIV disease are shown in Table 3. The analysis revealed a negative relationship between BMI score and HIV status regardless of gender ($p=0.02$ in males and $p=0.045$ in females), even after adjusted for age ($p=0.01$ in males and $p=0.03$ in females).

Table 4 shows the correlation between HIV status and meal frequency. There was a certain tendency that HIV-positive participants eat more frequently a day than HIV-negative participants regardless of gender. However, the only relationship between HIV status in male and meal frequency was statistically significant ($p=0.04$). Besides, no significant association between participants in low nutritional condition ($BMI<18.5$) and daily meal frequency was confirmed regardless of gender.

Table 1: Basic attributes

	Male:N=100 mean (SD)			Female:N=91 mean (SD)		
	HIV(+) n=39	HIV(-) n=61	P value	HIV(+) n=41	HIV(-) n=50	P value
BMI (kg/m2)	20.6±2.4	22.1±3.0	0.007	21.6±3.10	23.3±3.7	0.02
BMI<18.5 (%)	7 (17.9)	3 (5.0)	0.04	8 (19.5)	1 (2.0)	0.01
Age (years)	35.3±5.3	27.2±5.8	<0.0001	32.9±5.8	27.7±6.1	0.0001
Height (cm)	168.7±6.5	170.8±6.3	0.12	159.2±7.8	161.4±6.1	0.13
Weight (kg)	58.6±7.8	64.5±9.3	0.001	55.0±10.5	60.9±11.5	0.013
Number of meals (/day)	3.0±0.7	2.6±0.7	0.04	2.9±0.6	2.8±0.4	0.06
Smoking (%)	6 (15.4)	9 (14.8)	1.0	1 (2.0)	0 (0.0)	0.45
Drinking (%)	0 (0.0)	8 (13.1)	0.02	0 (0.0)	2 (4.0)	0.56
Chronic illness (%)	3 (8.7)	9 (14.8)	0.36	9 (21.9)	3 (6.0)	0.03
Family history of HIV	7 (17.9)	9 (14.8)	0.88	4 (9.8)	4 (8.0)	0.94

Table 2: Relative Frequency Distribution of HIV status and gender

	Male:N=100		Female:N=91	
	HIV(+):N=39	HIV(-):N=61	HIV(+):N=41	HIV(-):N=50
BMI	(%)	(%)	(%)	(%)
16.5-18.5	7 (17.9)	3 (4.9)	8 (19.5)	1 (2.0)
18.5-20.5	17 (43.6)	16 (26.2)	10 (24.4)	8 (16.0)
20.5-22.5	5 (12.8)	20 (32.8)	7 (17.1)	16 (32.0)
22.5-24.5	7 (17.9)	10 (16.4)	7 (17.1)	13 (26.0)
24.5-26.5	2 (5.1)	7 (11.5)	6 (14.6)	3 (6.0)
26.5-28.5	1 (2.6)	3 (4.9)	3 (7.3)	4 (8.0)
28.5-30.5	0	0	0	1 (2.0)
30.5-32.5	0	2 (3.3)	0	3 (6.0)
34.5-36.5	0	0	0	1 (2.0)

Table 3: Results of linear regression analysis composed of HIV status and other variable as an explanatory variable

	Male:N=100			Female:N=91		
	Coefficient	SD	P value	Coefficient	SD	P value
Adjustment of age						
HIV status	-1.76	0.69	0.01	-1.79	0.79	0.03
Age	0.03	0.05	0.6	0.02	0.06	0.76
Linear regression analysis						
HIV status	-1.62	0.71	0.02	-1.79	0.79	0.045
Age	0.03	0.05	0.54	0.02	0.06	0.78
Chronic illness	1.71	0.88	0.05	-0.78	1.13	0.49
Drinking	0.36	1.06	0.74	0.07	2.54	0.98
Smoking	-0.8	0.79	0.31	-1.69	3.59	0.64
Family history of HIV	0.33	0.68	0.63	-0.68	0.65	0.32
Chance of pregnancy				-0.29	2.12	0.89

158

Table 4: correlation between HIV status and meal intake situation

Male:N=100						
Number of meals/day	All			BMI<18.5		
	HIV(+)	HIV(-)	P value	HIV(+)	HIV(-)	P value
	(%)	(%)	Chi-squared test	(%)	(%)	Chi-squared test
Twice or less	7 (17.9)	25 (41.0)	0.04	2 (28.5)	0 (0.0)	0.3
3times	26 (66.7)	32 (52.4)		3 (43.0)	3 (100.0)	
4times or more	6 (15.4)	4 (6.6)		2 (28.5)	0 (0.0)	
Total	39 (100.0)	61 (100.0)		7 (100.0)	3 (100.0)	

Female:N=91						
Number of meals/day	All			BMI<18.5		
	HIV(+)	HIV(-)	P value	HIV(+)	HIV(-)	P value
	(%)	(%)	Chi-squared test	(%)	(%)	Chi-squared test
Twice or less	10 (24.4)	10 (20.0)	0.06	2 (25.0)	0 (0.0)	0.3
3times	25 (61.0)	39 (78.0)		4 (50.0)	1 (100.0)	
4times or more	6 (14.6)	1 (2.0)		2 (25.0)	0 (0.0)	
Total	41 (100.0)	50 (100.0)		8 (100.0)	1 (100.0)	

159

160

161

4. DISCUSSION

A limit of this study is a lack of information of daily meal portion and exercise amount. Since these factors could greatly impact BMI, we cannot state that low BMI of HIV-positive individuals caused by HIV infection or others such as daily meal portion and exercise amount [9].

166

167

5. CONCLUSION

168

169

In this study, a significant association between HIV infection and BMI was confirmed. Regardless of gender, individuals with HIV infection have lower BMI scores than individuals without HIV infection. The outcomes of this study shows that measuring and monitoring BMI score could be a key to catch symptoms of HIV infection early. According to DHS (Demographic and Health Survey) in Uganda, only 30% of HIV-positive individuals in Sub-Saharan Africa has access to anti-Retroviral (ARV) medication used for HIV treatment [10]. While HIV infection and AIDS development have promoted in those countries, effective countermeasures for them are needed. Of course, it is important to keep in mind that other symptoms such as fever, diarrhea or vomiting caused by HIV infection could breakout even in a widow period. Nevertheless, we believe that measuring and monitoring BMI as an easy and economical method promotes early detection and prevention of HIV. Lastly, we hope the findings in this study will contribute to future studies and improve care of HIV-positive individuals in Uganda, including early detection and access to testing opportunity and treatment.

182

183

184

CONCENT

185

As per standard of *School of Public Health*, Teikyo University, patient's written consent has been collected and preserved by the authors.

187

188

189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240

ETHICAL APPROVAL

As per standard of *School of Public Health*, Teikyo University, written approval of Ethics committee has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

6. REFERENCES

1. WHO (World Health Organization), Global Health Observatory data (2016)
Available: <http://www.who.int/gho/hiv/en/>
2. Dora Panagides, Rick Graciano, Peter Atekyereza, Lilia Gerberg and Mickey Chopra (1992). A review of nutrition and food security approaches in HIV and AIDS programmes in Eastern and Southern Africa. EQUINET DISCUSSION PAPER. NUMBER 48.
3. Results from the 2011 Uganda AIDS Indicator Survey (2011).
http://health.go.ug/docs/UAIS_2011_FACT_SHEET.pdf#search='Uganda+HIV++rate+increase'. Accessed date: 2014-1-9
4. Dr.Elizabeth Madraa (2004).Nutritional care and Support for people Living with HIV/AIDS in Uganda. Guidelines for Service Providers STD/AIDS Control Program Ministry of Health.
5. van der Sande MA, Schim van der Loeff MF, Aveika AA, Sabally S, Togun T, Sarge-Njie R, Alabi AS, Jaye A, Corrah T, Whittle HC (2004). Body mass index at time of HIV diagnosis: a strong and independent predictor of survival. *J Acquir Immune Defic Syndr*: 1288-94
6. Maro, T. Lahey, T. MacKenzie, L. Mtei, M. Bakari, M. Matee, K. Pallangyo, and C. F. von Reyn (2010). Low BMI and falling BMI predict HIV-associated tuberculosis: a prospective study in Tanzania. *NIH Public Access Int J Tuberc Lung Dis*. 2010 November ; 14(11): 1447–1453.
7. Snehal Shah,Christopher Whalen,Donald P. Kotler, Harriet Mayanja,Alice Namale George Melikian,Roy Mugerwa, and Richard D. Semba (2001).Severity of Human Immunodeficiency Virus Infection Is Associated with Decreased Phase Angle, Fat Mass and Body Cell Mass in Adults with Pulmonary Tuberculosis Infection in Uganda. *J Nutr. Nov.* 131(11):2843-7.
8. Science Daily, Loma Linda University Adventist Health Sciences Center (July 20,201).
Available: <https://www.sciencedaily.com/releases/2017/07/170720094844.htm>
9. John Curry, Esther Wieggers, Alessandra Garbero, Shannon Stokes and John Hourihan (2006). Gender,HIV/AIDS and Rural Livelihoods Micro-Level Investigations in Three African Countries. UNITED NATIONS UNIVERSITY Research Paper.No.2006/110.
10. Nobuyoshi Watahiki, Nobuyuki Hyou (2009). Towards Improving Health Seeking Behavior in Population and Health Related Areas in Developing Countries. Department of Education and Training Technology, National Institute of Public Health.