

Original Research Article

TITLE: Clinical profile and outcome of paediatric severe malaria in a North-Eastern Nigerian tertiary hospital.

ABSTRACT

Aims: To examine the clinical manifestations and outcome of severe malaria among children admitted into the Federal Medical Centre Azare, Nigeria.

Study design: The study was a cross sectional study.

Place and duration of study: The study was conducted at the department of Paediatrics, Federal Medical Centre, Azare, Bauchi state, Nigeria from 1st August to 31st October, 2013.

Methodology: Children aged 6 months to 12 years diagnosed with severe malaria were consecutively recruited from the Emergency Paediatric Unit. The information obtained included age, sex, weight, presenting complaints and associated symptoms. Elicited clinical signs, as well as outcome were also documented. The data was analysed using SPSS version 16.0.

Results: Ninety- eight patients diagnosed with severe malaria were recruited into the study. There were 57 (58.2%) boys and 41 (41.8%) girls giving a male to female ratio of 1.4: 1. The mean age of the patients was 4.2 ± 2.8 years, with a median and modal age of 3 years. Ninety patients (91.8%) survived and were discharged home while 8 died, giving a case fatality rate of 8.2%. Cough ($P = .04$), breathlessness ($P = .001$), reduction in urine output ($P = .003$), loss of consciousness ($P = .01$) and abdominal swelling ($P = .03$) were significantly more common in children aged 3 years and below. There were also statistically significant relationships between

age and nutritional status ($P=.03$), hepatomegaly ($P=.02$), coma ($P<.001$) as well as severe anaemia ($P=.03$). Binary logistics regression analysis showed that coma ($P=.03$) and dehydration ($P=.02$) were the most likely to predict death.

Conclusion: Prompt identification and management of fluid and electrolyte deficits may significantly reduce mortality in children with severe malaria managed at the Federal Medical Centre, Azare, Nigeria.

Keywords: Clinical profile, Severe malaria, Children, Coma, Dehydration, Outcome, Azare.

46

47

48

49

50 INTRODUCTION

51 For over 4000 years humanity has experienced the ravages of malaria. [1] Notwithstanding that
52 its incidence estimates have decreased globally in the last decade, malaria remains an important
53 cause of mortality particularly in children younger than 5 years in sub-Saharan Africa. [2] Of the
54 429,000 malaria related deaths recorded in 2015, 92% occurred in sub-Saharan Africa with two
55 countries; Nigeria and the Democratic Republic of Congo accounting for over a third of the total
56 deaths. [2]

57 Severe malaria remains the deadliest form of the disease and for the most part affects children
58 less than 5 years of age in endemic areas. [2- 4] The clinical manifestations and prognostic
59 factors of severe malaria have largely been described particularly in African children. However,
60 disease pattern and the comparative impact on outcome by the various symptoms and signs may
61 vary with age, access to health care, endemicity, genetics and geographical location. [5-9]

62 Understanding of the clinical spectrum and relative outcomes of severe malaria amongst children
63 in our locale may aid in prompt diagnosis and appropriate treatment as well as identify gaps in
64 knowledge requiring further research. Thus we carried out a cross sectional study aimed at
65 examining the clinical manifestations and outcome of severe malaria among children admitted
66 into the Emergency Paediatric Unit (EPU) of the Federal Medical Centre (FMC) Azare in North-
67 Eastern Nigeria. To our knowledge this is the first of such studies to be conducted in this part of
68 Nigeria.

METHODOLOGY

The present study was conducted at the EPU of the Federal Medical Centre Azare, Bauchi state from 1st August to 31st October, 2013 (peak of the rainy season and period of highest transmission). The centre is a tertiary facility which serves as a referral hospital for the populations of Bauchi, Yobe and Jigawa states of Northern Nigeria. Prior approval was obtained from the research ethics committee of the hospital.

The patients (aged 6 months to 12 years) were consecutively recruited from the EPU excluding those who died on arrival. Confirmation of severe malaria diagnosis was done with microscopy (*Plasmodium falciparum* (PF) parasitaemia confirmed with blood film microscopy) or rapid diagnostic tests (RDT) in the presence of any of the World Health Organization (WHO) case definitions for severe malaria. [4,10] Information was obtained with the use of a structured questionnaire. Informed consent was obtained verbally from the caregivers. The information included age, sex, weight, presenting complaints and associated symptoms. The duration of hospital stay, as well as the outcome was also documented. The axillary temperatures were measured with a mercury thermometer and recorded. Hypothermia was defined as axillary temperature $< 35^{\circ}\text{C}$ while hyperpyrexia was defined as axillary temperature $> 38.5^{\circ}\text{C}$.

Anthropometric measurements were taken and recorded. Other information obtained included hydration status, nutritional status (using the modified Wellcome classification), liver span and splenic size in the presence of splenomegaly. Consciousness level was determined with the modified Glasgow Coma Scale. The RDT was done using the paracheck Pf[®] rapid test kit for *P. falciparum* malaria (a rapid test for the detection of *Plasmodium falciparum* specific histidine rich protein-2). Results were documented as positive or negative as indicated by the coloured bands on the control windows; test ('T') and control ('C') respectively. Haemoglobinuria was

defined as the presence of dark or brownish coloured urine and severe anaemia as haemoglobin concentration $< 5\text{g/dl}$. [6] All the patients were managed using the WHO treatment guidelines. [11] The collected data was entered into SPSS version 16.0. Categorical data were compared using the Chi-square test. A *P* value less than 0.05 was regarded as being statistically significant. The binary logistics regression analysis was applied to determine the most likely predictors of outcome utilizing 95% confidence interval.

RESULTS

Ninety- eight cases of severe malaria were recruited into the study. The patients were aged 6 months to 12 years. There were 57 (58.2%) boys and 41 (41.8%) girls giving a male to female ratio of 1.4: 1. The mean age of the patients was 4.2 ± 2.8 years, with a median and modal age of 3 years. Fifty percent (50%) of the patients were less than 1 year of age while 2 % were between the ages of 10 years and 12 years. Of the 98 patients 90 (91.8%) survived and were discharged home while 8 died, amounting to a case fatality rate (CFR) of 8.2%. Table 1 displays the age and sex distribution of the patients.

Table 1. Age and sex distribution of children with severe malaria.

Age (years)	Sex		Total (%)
	Male (%)	Female (%)	
< 1	27(27.6)	22(22.4)	49(50.0)
1-3	16(16.3)	12(12.2)	28(28.6)
4-6	10(10.2)	4(4.1)	14(14.3)

7-9	3 (3.1)	2 (2.0)	5(5.1)
10-12	1 (1.0)	1 (1.0)	2(2.0)
Total	57 (58.2)	41 (41.8)	98(100)

Table 2 shows the distribution of presenting symptoms by age. All the patients had fever, 50 (51%) had presenting complaint of refusal to feed made by their caregivers. Twenty-five (25.5%) of these children were aged less than 1 year while none of them was between ages 10 and 12 years. The symptoms of cough ($P = .04$), breathlessness ($P = .001$), reduction in urine output ($P = .003$), loss of consciousness ($P = .01$) and abdominal swelling ($P = .03$) had statistically significant relationships with age.

The relationships between the clinical signs elicited in the patients and their ages are outlined in Table 3. Thirty-five (35.7%) patients had hyperpyrexia, while 3 (3.1%) had hypothermia. Twenty-eight (28.5%) patients were dehydrated at presentation with the majority; 14(14.3%) being less than age 1 year. Of the 28 (28.5%) children who presented with features of dehydration, 14 (14.3%) were aged 6 months to 11 months whereas 1(1%) was aged 10 years-12 years. There were statistically significant relationships between age and nutritional status ($P = .03$), hepatomegaly ($P = .02$), coma ($P < .001$) as well as severe anaemia ($P = .03$).

125

126

127

128 Table 2. Distribution of presenting symptoms by age

Symptoms	Age (years)					Total	<i>P</i>
	<1(%)	1-3(%)	4-6(%)	7-9(%)	10-12(%)		
Fever	49(50)	28(28.6)	14(14.3)	5(5.1)	2(2.0)	98(100)	-
Refusal to feed	25(25.5)	17(17.3)	5(5.1)	3(3.1)	0	50(51.0)	.39
Vomiting	23(23.5)	11(11.2)	6(6.1)	3(3.1)	2(2.0)	45(45.9)	.50
Diarrhoea	11(11.2)	11(11.2)	1(1.0)	1(1.0)	0	24(24.5)	.43
Cough	20(20.4)	10(10.2)	2(2.0)	1(1.0)	0	33(33.7)	.04
Breathlessness	20(20.4)	6(6.1)	0	0	0	26(26.5)	.001
Reduction in urine output	13(13.3)	1(1.0)	0	0	0	14(14.3)	.003
Coca-Cola coloured urine	5(5.1)	5(5.1)	2(2.0)	1(1.0)	0	13(13.3)	.65
Bloody urine	1(1.0)	1(1.0)	1(1.0)	0	0	3(3.1)	.73
Jaundice	11(11.2)	6(6.1)	2(2.0)	1(1.0)	0	20(20.4)	.43
Convulsion	32(32.7)	17(17.3)	10(10.2)	3(3.1)	1(1.0)	63(64.3)	.87
Loss of	14(14.3)	10(10.2)	8(8.2)	3(3.1)	2(2.0)	37(37.8)	.01

consciousness

Abdominal	10(10.2)	3(3.1)	0	0	0	13(13.3)	.03
-----------	----------	--------	---	---	---	----------	-----

swelling

129

130

131 Table 3. Distribution of clinical signs by age.

Clinical signs	Age (years)					Total	<i>P</i>
	<1(%)	1-3(%)	4-6(%)	7-9(%)	10-12(%)		
Hypothermia	2(2)	0	1(1)	0	0	3(3.1)	.12
Hyperpyrexia	13(13.3)	13(13.3)	5(5.1)	2(2.0)	2(2.0)	35(35.7)	.12
Dehydration	14(14.3)	7(7.1)	3(3.1)	3(3.1)	1(1.0)	28(28.5)	.57
Oedema	4(4.1)	0	0	1(1.0)	0	5(5.1)	.64
Nutritional status							
Normal	13(13.3)	14(14.3)	8(8.2)	3(3.1)	2(2.0)	40(40.8)	
Underweight	31(31.6)	13(13.3)	6(6.1)	2(2.0)	0	52(53.1)	
Marasmus	2(2.0)	1(1.0)	0	0	0	3(3.1)	
Kwashiorkor	1(1.0)	0	0	0	0	1(1.0)	
Marasmic kwashiorkor	2(2.0)	0	0	0	0	2(2.0)	.03
Hepatomegaly	24(24.5)	11(11.2)	2(2.0)	2(2.0)	0	39(39.8)	.02
Splenomegaly	7(7.1)	2(2.0)	1(1.0)	0	0	10(10.2)	.18
Coma	7(7.1)	8(8.2)	7(7.1)	3(3.1)	2(2.0)	27(27.6)	<.001
Severe anaemia	18(18.4)	7(7.1)	1(1.0)	1(1.0)	0	27(27.6)	.03

132

133

134

135 Table 4 relates outcome to the patients' symptoms. None of the symptoms studied reached
 136 statistical significance in their relationship with outcome. However, 6 (6.1%) of the 8 (8.2%)
 137 patients who died had convulsions. Table 5 shows the relationship between some clinical signs
 138 and outcome. Twenty – seven (27.6%) patients presented with severe anaemia and 3 (3.1%) died.
 139 Coma and dehydration had statistically significant relationships with outcome. The ability of the
 140 individual clinical signs to predict death from severe malaria was tested using the binary logistics
 141 regression model and as shown in Table 6, coma and dehydration ($P = .03$ and $P = .02$
 142 respectively) were the most likely to independently predict death in this study.

143 Table 4. The association of outcome with clinical symptoms

144

Symptoms	Outcome		No of patients N=98(%)	X^2	P
	Survived (%)	Died (%)			
Fever	90(91.8)	8(8.2)	98(100)	-	-
Jaundice	18(18.4)	2(2.0)	20(20.4)	.11	.74
Convulsion	57(58.2)	6(6.1)	63(64.3)	.43	.51
Haematuria	2(2.0)	1(1.0)	3(3.1)	2.59	.11

Cough	30(30.6)	3(3.1)	33(33.7)	.01	.81
Vomiting	41(41.8)	4(4.1)	45(46.4)	.05	.83
Diarrhoea	20(20.4)	4(4.1)	24(24.5)	3.03	.08

145

146 Table 5. The association between outcome and clinical signs

147

Clinical signs	Outcome		No of Patients N=98(%)	X^2	<i>P</i>
	Survived (%)	Died (%)			
Severe anaemia	24(24.5)	3(3.1)	27(27.6)	.43	.51
Haemoglobinuria	11(11.2)	2(2.0)	13(13.3)	1.03	.31
Hypothermia	2(2.0)	1(1.0)	3(3.1)	2.59	.11
Hyperpyrexia	33(33.7)	2(2.0)	35(35.7)	.43	.51
Coma	22(22.4)	5(5.1)	27(27.6)	5.28	.02
Splenomegaly	10(10.2)	0	10(10.2)	.98	.32
Hepatomegaly	37(37.8)	2(2.0)	39(39.8)	.98	.32
Severe malnutrition	5(5.1)	1(1.0)	6(6.1)	.03	.86
Dehydration	23(23.4)	5(5.1)	28(28.5)	6.71	.01
Oliguria	12(12.2)	2(2.0)	14(14.3)	.81	.37

148

149
150
151
152
153
154

155 Table 6. Binary logistics regression analysis of signs of severe malaria on outcome

Clinical signs	β	X^2	P	95% Confidence Interval
Coma	-1.64	4.69	.03	.04, .88
Dehydration	1.48	5.52	.02	1.28, 15.16
Hyperpyrexia	-.55	.46	.51	.11, 3.02
Hypothermia	1.84	1.62	.15	.51, 78.19
Severe anaemia	-.50	.41	.51	.13, 2.73
Severe malnutrition	.08	.03	.86	.44, 2.70
Oliguria	.70	.70	.38	.39, 12.00
Haemoglobinuria	.87	.88	.32	.43, 13.39
Splenomegaly	-18.90	1.80	.18	.00
Hepatomegaly	-.76	1.12	.33	.10, 2.16

156
157
158
159

DISCUSSION

Our study has demonstrated that severe malaria is more prevalent among preschool children. This is in conformity with multiple studies done across sub-Saharan Africa. [12-15] However, what is at odds with findings from other studies is the observation that 50% of our patients were aged 6 - 11 months. [8, 12, 16] Several reasons have been adduced to justify the low prevalence of severe malaria in children younger than 1 year. These include; breast milk is low in para-aminobenzoic acid (PABA) which is vital for the replication of PF, high foetal haemoglobin levels impeding the growth of PF and acquired passive immunity from immune mothers in endemic areas. [17,18] The reasons for the contrary observation in the present study is unclear. Yet it has been demonstrated that in endemic areas children born during the low-transmission season possess significantly higher risk of being infected in comparison to those born during the high transmission season. [19] This may explain our finding given the fact that most of the patients less than age 1 year were born between the months of December and April which is part of the low transmission season in the Sahel region of sub-Saharan Africa. [20] Secondly, some authors have suggested that maternally acquired antibodies may be a marker for risk of infection instead of that of protective immunity. [21,22]

Our case fatality rate (CFR) of 8.2% is lower than the estimated average for sub-Saharan Africa. [13] It is also lower than the 10.3% reported in parts of the Niger-delta region of Nigeria, [23] but higher than the 5.2% recorded in Osogbo, South West Nigeria. [14] The later study was done retrospectively and consisted of a small sample size (58) hence the findings are likely to be less representative than that of the present study - done prospectively with a larger sample size. Be that as it may, the fact that the patients who died on arrival were excluded from our study may have largely confounded the CFR.

There were notable age-related differences in the clinical features of our patients. Cough was a significantly frequent symptom in children less than age 5 years and non-existent in those older than 6 years. It was found at rates higher than that reported at Osogbo, South Western Nigeria but lower than that reported in Uganda. [16,24] Respiratory symptoms have long been known to be associated with malaria, furthermore, studies have recognized small airway obstruction, impaired gas exchange and increased pulmonary phagocytic activity in malaria patients exhibiting pulmonary symptoms.[25-27] These may explain the advent of respiratory symptoms in children with malaria.

Breathlessness, reduction in urine output, loss of consciousness and abdominal swelling were also more common in children aged 3 years and less, whereas hepatomegaly, severe anaemia and coma were more frequent in those aged 5 years and below. These are consistent with findings from previous studies. [24] No clear explanation has been given for the age-associated differences in the clinical presentation of severe malaria. However, there are indications that several factors may be responsible including; lower levels of complement regulatory proteins, significantly smaller red cell mass as well as differential organ sequestration of PF in younger children in comparison to older children putting them at greater risk of certain manifestations. [28-30]

Coma and dehydration were independently associated with an increased risk of death in this study. This is in keeping with findings from multiple studies. [5,12,31] Nevertheless, it is inconsistent with the findings reported in Ado-Ekiti, South Western Nigeria. [8] The sequestration of parasitized erythrocytes in the microvasculature of the brain is fundamental to the aetiology of coma in severe malaria, deepening coma and subsequent affection of the respiratory centre may lead to death. [32] The significant association of dehydration with death

here may have stemmed from the cumulative prevalence of diarrhoea and vomiting (8.2%) among our patients and failure to promptly manage fluid and electrolyte deficits. Severe anaemia was not independently associated with an increased risk of death in our study. This may be largely due to the fact that blood transfusions were promptly administered to patients diagnosed with severe malarial anaemia. Blood transfusions have been shown to be effective at preventing death in severe malarial anaemia. [33]

CONCLUSION

We have demonstrated that cough, breathlessness, reduction in urine output, abdominal swelling and loss of consciousness are common symptoms of severe malaria in preschool children. Severe anaemia, hepatomegaly and coma are also frequent in this age group. The predictors of death as found in this study are coma and dehydration. We recommend prompt identification and management of fluid and electrolyte deficits in children presenting with severe malaria to the Federal Medical Centre, Azare. In addition, parents should be continuously educated on the significance of uninterrupted feeding of their children when they are ill and the import of home management of diarrhoea and vomiting with oral rehydration solution before presentation to the hospital.

ETHICAL APPROVAL

Approval was obtained from the research ethics committee of the Federal Medical Centre Azare, Bauchi state, Nigeria before commencement of the study.

COMPETING INTERESTS

The authors hereby declare that there were no competing interests involved in this study.

REFERENCES

1. Cox FEG. History of the discovery of the malaria parasites and their vectors. Parasit Vectors. 2010; 3: 5. Available: <http://www.parasitesandvectors.com/content/3/1/5>. (Accessed 17 October 2017).
2. World Health Organization. World Malaria report 2016: Summary. WHO Geneva; 2017 (WHO/HTM/GMP/2017.4). Available: <http://www.who.int/malaria/publications/world-malaria-report-2016/report/en/>. (Accessed 16 October 2017).
3. Dondrop AM, Lee SJ, Faiz MA, Masha S, Price R. The relationship between age and the manifestations of and mortality associated with severe malaria. Clin Infect Dis. 2008; 47: 151-7.
4. World Health Organization. Guidelines for the treatment of malaria. 3rd Ed. WHO Geneva; 2015. Available:

http://apps.who.int/iris/bitstream/10665/162441/1/9789241549127_eng.pdf?ua=1&ua=1.

(Accessed 17 October 2017).

5. Von Seidlein L, Olaosebikan R, Hendriksen ICE, Lee SJ, Adedoyin OT, Agbenyega T, et al. Predicting the clinical outcome of severe falciparum malaria in African children: findings from a large randomized trial. CID. 2012; 54(8): 1080-90.
6. World Health Organization. Severe falciparum malaria. Trans R Soc Trop Med Hyg. 2000; 94(Suppl 1): 1-90.
7. Pankoui Mfonkeu JB, Gouado I, Fotso Kuate H, Zambou O, Grau G, Combes V, et al. Clinical presentation, haematological indices and management of children with severe and uncomplicated malaria in Douala, Cameroon. Pak J Biol Sci. 2008; 11(20): 2401-6.
8. Oluwayemi OI, Brown BJ, Oyedeji OA, Adegoke SA, Adebami OJ, Oyedeji GA. Clinical and laboratory predictors of outcome in cerebral malaria in suburban Nigeria. J Infect Dev Countr. 2013; 7(8): 600-7.
9. Manning L, Laman M, Law I, Bona C, Aipit S, Teine D, et al. Features and prognosis of severe malaria caused by *Plasmodium falciparum*, *Plasmodium vivax* and mixed *Plasmodium* species in Papua New Guinea children. PLoS ONE. 2011; 6(12): e29203. Available: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0029203>. (Accessed 17 October 2017).
10. World Health Organization. Basic malaria microscopy (part 1: learner's guide) 2nd Ed. WHO 2010. Available: <http://www.who.int/malaria/publications/atoz/9241547820/en/>. (Accessed 17 October 2017).
11. World Health Organization. Management of severe malaria: A practical handbook. 3rd Ed. WHO Geneva; 2012. Available:

http://apps.who.int/iris/bitstream/10665179317/1/9789241548526_eng.pdf. (Accessed 17 October 2017).

12. Orimadegun AE, Fawole O, Okereke JO, Akinbami FO, Sodeinde O. Increasing burden of childhood severe malaria in a Nigerian tertiary hospital: Implication for control. *J Trop Pediatr*. 2007; 53(3): 185-9.
13. Maitland K. Severe malaria in African children- The need for continuing investment. *N Engl J Med*. 2016; 375: 2416-7.
14. Oyedele OA, Oluwayemi IO, Afolabi AA, Bolaji O, Fadero FF. Severe malaria at a tertiary paediatric emergency unit in South West Nigeria. *Research Journal of Medical Sciences*. 2010; 4(6): 352-6.
15. Kunuanunua TS, Nsibu CN, Bodi JM, Tshibola TK, Bura MM, Magoga K, et al. Severe malaria in children: A descriptive report from Kinshasa, the Democratic Republic of Congo. *J Trop Pediatr*. 2015; 61: 272-8.
16. Oninla SO, Ogunro PS, Oninla OA, Kayode OV. Childhood cerebral malaria in Nigeria: Clinical features, treatment and outcome. *IJTDH*. 2016; 12(4): 1-12.
17. Amaratunga C, Lopera-Mesa TM, Brittain NJ, Cholera R, Arie T, Fujioka H, et al. A role for fetal haemoglobin and maternal immune IgG in infant resistance to *Plasmodium falciparum* malaria. *PLoS ONE* 2011; 6(4): e14798. Available: <https://doi.org/10.1371/journal.pone.0014798>. (Accessed 19 October 2017).
18. Jiya MN, Airede KL, Ahmed H. Cerebral malaria: Presentation and outcome in children in Sokoto. *Nig Med Pract*. 2006; 50: 55-6.

19. Riley EM, Wagner GE, Ofori MF, Wheeler JG, Akanmori BD, Tetteh K et al. Lack of association between maternal antibody and protection of African infants from malaria infection. *Infect Immun*. 2000; 68(10): 5856-63.
20. Samdi LM, Ajayi JA, Oguche S, Ayanlade A. Seasonal variation of malaria parasite density in paediatric population of North Eastern Nigeria. *Global Journal of Health Science*. 2012; 4(2): 103-9.
21. Kitau AY, Urassa H, Wechsler M, Smith T, Vounatsou P, Weiss NA, et al. Antibodies against *Plasmodium falciparum* vaccine candidates in infants in an area of intense and perennial transmission: Relationships with clinical malaria and with entomological inoculation rates. *Parasite Immunol*. 1999; 307-17.
22. Wagner G, McGuinness D, Koram K, Bennett S, Nkrumah FK, Riley E. High incidence of asymptomatic malaria infections in a birth cohort of children under 1 year of age in Ghana, detected by multicopy gene polymerase chain reaction. *Am J Trop Med Hyg*. 1998; 59: 115-23.
23. Imananagha K. The presenting complaints of children with severe malaria. *Ann Trop Med Public Health*. 2010; 3: 68-71.
24. Idro R, Bitarakwate E, Tumwesigire S, John CC. Clinical manifestations of severe malaria in the highlands of South Western Uganda. *Am J Trop Med Hyg*. 2005; 72(5): 561-7.
25. Taylor WRJ, Canon V, White NJ. Pulmonary manifestations of malaria. *Treat Respir Med*. 2006; 5(6): 419-28.
26. Anstey NM, Jacups SP, Cain T, Pearson T, Ziesing PJ, Fisher DA, et al. Pulmonary manifestations of uncomplicated *falciparum* and *vivax* malaria: cough, small airways

obstruction, impaired gas transfer, and increased pulmonary phagocytic activity. J Infect Dis. 2002; 185: 1326-34.

27. Basset Q, Machevo S, O'Callaghan-Gordo C, Sigauque B, Morais L, Diez-Padrisa N, et al. Distinguishing malaria from severe pneumonia among hospitalized children who fulfilled Integrated Management of Childhood Illness criteria for both diseases: A hospital-based study in Mozambique. Am J Trop Med Hyg. 2011; 85(4): 626-34.

28. Waitumbi JN, Donvito B, Kisserli A, Cohen JHM, Stoute JA. Age-related changes in red blood cell complement regulatory proteins and susceptibility to severe malaria. J Infect Dis. 2004; 190(6): 1183-91.

29. Molineaux L. Plasmodium falciparum malaria: Some epidemiological implications of parasite and host diversity. Ann Trop Med Parasitol. 1996; 90(4): 379-93.

30. Marsh K, Snow RW. Host-parasite interaction and morbidity in malaria endemic areas. Philos Trans R Soc Lond B Biol Sci. 1997; 352: 1385-94.

31. Kwenti TB, Kwenti TDB, Latz A, Njunda LA, Akenji TN. Epidemiological and clinical profile of paediatric malaria: A cross sectional study performed on febrile children in five epidemiological strata of malaria in Cameroon. BMC Infect Dis. 2017; 17: 499-511.

32. Newton CR, Hien TT, White N. Cerebral malaria. J Neurol Neurosurg Psychiatry. 2000; 69: 433-41.

33. English M, Ahmed M, Ngando C, Berkley J, Ross A. Blood transfusion for severe anaemia in children in a Kenyan hospital. Lancet. 2002; 359: 494-5.

338

339

340

341

342

343

344

345

346

347

348

349