

Growth Performance, Metabolic Efficiency and Nutrient Utilization of BALB/C Mice Infected with *Leishmania major* Fed with Standard Rat Pellets and *Annonaceae* Fruit Pulp Pellets

ABSTRACT

This study was aimed at evaluating the clinical, biochemical, and hematological changes in male BALB/C mice infected with *Leishmania major* fed with standard rat pellets (RP) and *Annonaceae* Fruit Pulp pellets (AFPP) in different experimental exposures. The results of the study showed good palatability, acceptability and normal behaviour of the mice during the whole experimental period except in the infected groups. Furthermore, the results achieved with low levels of *Annonaceae* fruit pulp pellets crude proteins (3.81 ± 0.14) and lipids (2.95 ± 0.18) were comparable with that of RP at a level of 25.28 ± 0.04 and 18.60 ± 0.06 respectively. The growth parameter, metabolic efficiency and feed utilization parameters between the rat pellets and *Annonaceae* fruit pulp pellets in non-infected, infected non-treated and infected treated groups of mice did not differ. At the same time haematological changes, liver and kidney function parameters and lipid profile of the mice that were feed with rat pellets and *Annonaceae* fruit pulp pellets in non-infected, infected non-treated and infected treated were also comparable same. Other parameter measured in the study such as weight of organs (liver and spleen) parasite burden (LDU) followed the same trend. However, some of the infected non-treated groups had parameters beyond the normal ranges than the non-infected and infected treated groups at the end of the experimental period. It can be concluded that *Annonaceae* fruit pulp pellets can be utilized as source of raw material in the manufacture of animal feeds and nutraceuticals upon further research in higher animal model.

Key Words: BALB/C, *Leishmania major*, *Annonaceae*, Growth performance, metabolic efficiency, feed utilization

INTRODUCTION

One of the world's most devastating neglected tropical diseases of documented epidemiological; and experimental public health importance is Leishmaniasis [1]. In developing countries where it is manifested, it often co-exists with chronic malnutrition, one of the main risk factors for its development [2, 3, 4]. In most of the study conducted, only a few emphasized on the relationship between leishmaniasis progression and malnutrition. Malnutrition is a serious health problem that remains common in many parts of the world [5, 6].

In the developing world, Leishmaniasis combined with malnutrition is a public health problem. The most frequent form of malnutrition is Protein energy malnutrition (PEM). Globally, about 800 million people, including over 150 million children under 5 years of age, most of them in developing countries, are affected by PEM [7]. The excess morbidity and mortality associated with malnutrition is due to the impairment of sufferers' defense mechanisms, which predisposes them to infectious diseases [8]. Further, the poor understanding of the impact of PEM on immune response against *Leishmania* infection coupled with neglected nutritional status of the host [9, 10], worsen the situation. Studies have revealed that good protein energy (PE) promote adaptive immune response in mice infected with *L. chagasi* [11]. The BALB/C mice provide a unique opportunity to study human cutaneous leishmaniasis (CL) in its active form due to their susceptibility to *L. major* infection. As a model, it reproduces clinical and pathological features of human CL.

Food insecurity, especially in the developing world, has necessitated research in alternative source of food substances. Therefore, natural products, notably those derived from plants, have been used to help mankind sustain its nutrition and health since the dawn of medicine. The importance of plants in agriculture and medicine has stimulated significant scientific interest [12]. However, a restricted range of plant species, with regard to their medicinal importance, has experienced detailed scientific inspection, resulting into comparatively insufficient knowledge concerning their potential role in nutraceutical application [13]. Therefore, to attain a reasonable perception of medicinal values/benefits, comprehensive investigations on the role of plants in the management of human ailments are needed.

In a nutraceutical landscape, plants with a long history of use in ethno medicine are a rich source of active phyto-constituents that provide medicinal or health benefits against various ailments and diseases [14]. One such plant family with extensive traditional use in Kenya and mostly grown in the coast region is *Annonaceae* family which include; *A. cherimola*; Mill (cherimoya), *A. muricata*; L (soursop), *A. squamosa*; L (sugar apple) *A. senegalensis*; L (wild soursop) and *A. reticulata*; L (custard apple). The fruits of these plants contain numerous bioactive substances [15]. These fruits are also possible contributors of the carotenoids, vitamins, mineral salts, fibres and bioactive compounds [16, 14]. Thus they can be incorporated as components of diets to provide a delicate balance of food security. The fact that they are used as food by other parts of the world [16] with Kenyans having similar gastronomic habits [17] and also exhibit medicinal potency is a significant aspect of modern trends in research focusing on food-medical interface (neutraceuticals).

Nonetheless, these plants are neglected especially as sources of alternative foods for reasons not well understood. The present paper reports the influence of diet made from *Annonaceae* fruit pulp on growth performances, metabolic efficiency and nutrients utilization on BALB/C mice infected with *L. major*. This might reflect on proper

utilization of these neglected crops for development of nutraceutical products. It might also contribute towards curbing food insecurity in the region where they are grown.

MATERIALS AND METHODS

Plant material

The ripe fresh fruits of Annonaceae (*A. muricata* and *A. squamosa*) were collected from farms in coast province (Kilifi and Kwale Counties) of Kenya. The harvested fruits collected were then transported to the Jomo Kenyatta University of Agriculture and Technology (JKUAT) in the Department of Food Science Technology, Food Biochemistry laboratories. In the laboratory, the fruits pulps were dried using a constant temperature and humidity chamber (Tokyo Thermo Tech Co. Ltd, Japan) set at 25°C. Thereafter, feed pellets were made from the dried fruit pulp. The pellets made were taken to the Centre for Biotechnology Research and Development (CBRD), Kenya Medical Research Institute (KEMRI) leishmaniasis laboratories for *in vivo* trials.

Animal feeds

Standard rat pellets (Rate pellets®, Unga Feeds Ltd, Kenya) and Annonaceae fruit pulp (AFP) were used in the study. To ensure palatability of the AFP formulated feed, cubes were made from the powdered AFP by mixing with wheat flour as a binder after addition of water using a pellet-making machine. Proximate composition analysis of the feeds was performed as described in the Association of Official Analytical Chemists (AOAC) manual [18].

Parasites

Metacyclic promastigotes of *L. major* (strain IDUB/KE/83=NLB-144) maintained by cryopreservation and *in vitro* culture, and periodic passage in BALB/c mice at KEMRI were cultured in Schneider's insect medium (Gibco, Invitrogen), supplemented with 20% fetal calf serum (FBS) (Life Technologies), glutamine (2mM), penicillin G (100U/mL), and streptomycin (100µg/mL). The parasites were harvested in the stationary phase after 8–10 days of culture, centrifuged (at 1000rpm for 5 min), washed twice with Schneider's insect medium, counted, and used to inoculate mice.

Animals

Male BALB/c mice (3-4 weeks old) obtained from Kenya Medical Research (KEMRI) animal house were fed with standard rat pellets (Rate pellets®, Unga Feeds Ltd, Kenya), Annonaceae fruit pulp (AFP) and water ad libitum and kept under standard temperature (26°C and 60% humidity) and a natural light-darkness cycle. All *in vivo* experiments with mice were performed according to the established bioethical standards of the KEMRI's Animal Care and Use Committee (ACUC), Scientific and Ethics Review Unit (SERU).

Experimental Design

The mice were equally and randomly allocated into experiment groups assigned to groups of six mice per group as follows; Non-Infected fed with Rat pellets (NI-RP), Infected Non-Treated fed with Rat pellets (INT-RP), Infected Treated fed with Rat pellets (IT-RP), Non-Infected fed with Annonaceae fruit pulp (NI-AFP), Infected Treated fed with Annonaceae fruit pulp (IT-AFP), Infected Non-Treated fed with Annonaceae fruit pulp (INT-AFP). After the different experimental exposures, the mice were closely monitored on a daily basis for agility, hair ruffling, appetite, vomiting, urine colour, skin turgor, ocular tension, limb paralysis, convulsions and roll-over movements [19]. Body weight and temperature were measured on a weekly basis. After lesion have developed, density of parasites was determined by counting the number of amastigotes form of parasite from smears made from ulcer stained with Leishman's stain [20]. The infected footpads were measured using a direct reading vernier caliper and lesion size calculated [21]. Means of weekly readings were calculated to facilitate comparison of lesion progression. Growth performance and nutrient utilization was assessed at the end of the experimental period (12 weeks) described by as follows; Body mass gain (BMG) = $[(\text{FBM}) - (\text{IBM})]/\text{IBM}] \times 100$, Specific growth rate (SGR% per day) = $[(\text{FBM}) - (\text{IBM})/\text{number of trial days}] \times 100$, Metabolic growth rate (MGR g/kg^{0.8}/day) = $(\text{BMG})/[(\text{IBM}/1000)^{0.8} + (\text{FBM}/1000)^{0.8}]/2]/\text{duration of the trial days}$, Feed conversion ratio (FCR) = Dry feed fed (g)/BMG (g), Protein efficiency ratio (PER) = fresh BMG (g)/Crude Protein (CP) fed (g), Protein productive value (PPV)% = $[(\text{final mice body protein in g} - \text{initial mice body protein in g})/\text{total protein consumed in g}] \times 100$ and Apparent lipid conversion (ALC)% = $[(\text{final mice body lipid in g} - \text{initial mice body lipid, g})/\text{total crude lipid consumed in g}] \times 100$ [22]. At the end of the experimental period (90 days), the animals were fasted for about 4 hours with free access to water and sacrificed by inoculation with 100µL of pentobarbitone sodium (Sagatal®). 2 mL of blood sample was collected for determination of haematological and biochemical parameters, renal and liver functions and blood metabolites using a Retroflon® Plus automated analyzer. Liver and spleen impression smears were used to quantitate the parasite loads [23].

Data analysis

Haematology and biochemical parameters, blood metabolites, growth performance parameters and body weights data were analyzed using mean separation by GenStat program [24]. Mean separation was done through Fischer least significance difference. Comparisons between two treatments were done by means of unpaired Student's t-test and significance established by ANOVA. A difference of $P < 0.05$ was considered statistically significant.

RESULTS

Proximate composition analysis and palatability of the experimental feeds

Good palatability, acceptability and normal behaviour were observed during the whole experimental period except in the infected group. In the non-infected group, minimal

residual feeds were observed in the cages. The feed composition parameters in % Dry Weight Basis (DWB) of the two experimental feeds (Table 1) differ significantly ($p < 0.05$). The two experimental feeds were analyzed for proximate feed composition parameters in % Dry Weight Basis (DWB); carbohydrates, crude lipids, crude proteins, moisture content, ash content, dry matter and fibre content (Table 1). The analyzed proximate composition parameter of the two feeds; RP and AFPP differed significantly ($P < 0.05$). The RP had a higher value of crude protein and crude lipids, 25.28 ± 0.04 and 18.60 ± 0.06 respectively against the AFPP; 3.81 ± 0.14 and 2.95 ± 0.18 respectively. However, the AFPP had higher values of carbohydrates, moisture content, ash content, dry matter and fibre content of 40.12 ± 0.38 , 6.20 ± 0.12 , 5.93 ± 0.50 , 93.80 ± 0.12 and 59.56 ± 2.64 against the RP of 27.14 ± 2.05 , 2.17 ± 0.98 , 3.51 ± 0.04 , 1.42 ± 0.24 and 25.67 ± 1.28 respectively. Good palatability, acceptability and normal behaviors were observed during the whole experimental period in all groups of BALB/C mice except in the infected groups.

Table 1: Proximate composition analysis of diets used in feeding BALB/C mice

Parameter	Experimental diets	
	Rat Pellets (RP)	Annonaceae Fruit Palp Pellets (AFPP)
Crude Proteins	$25.28^a \pm 0.04$	$3.81^b \pm 0.14$
Crude Lipids	$18.60^a \pm 0.06$	$2.95^b \pm 0.18$
Carbohydrate	$27.24^a \pm 2.05$	$40.12^b \pm 0.38$
Moisture content	$2.17^a \pm 0.98$	$6.20^b \pm 0.12$
Ash content	$3.51^a \pm 0.04$	$5.93^b \pm 0.50$
Dry matter	$1.42^a \pm 0.24$	$93.80^b \pm 0.12$
Fibre content	$25.67^a \pm 1.28$	$59.56^b \pm 2.64$
Mean values ($n=3$) $\pm$ SEM. Values appended by different small letters within a row are significantly different ($P < 0.05$).		

Growth performance and nutrient utilization

The body mass changes in the different groups of BALB/C mice measured weekly. Generally, the BMG was observed in the two NI groups of BALB/C mice; NI-RP and NI-AFPP. However, the BMG in the NI-RP group was higher (56.10 ± 1.49) compared to that of NI-AFPP (49.11 ± 1.39). A significant decrease in body weight was observed in both the IT-RP and IT-AFPP before treatment (Figure 1). However, this reversed after treatment with Pentostam in week 4 with the IT-RP and IT-AFPP groups having almost the same BMG values of 27.31 ± 0.66 and 27.56 ± 1.37 respectively at the end of the experimental period. There was also a significant decrease in BMG in both INT-RP and INT-AFPP groups ($P < 0.05$) (Figure 1) with the INT-AFPP group having greater decrease in BMG (-40.71 ± 0.68) than the INT-RP group (-34.62 ± 0.79).




Figure 1: Body weight changes of BALB/c mice infected with *L. Major* fed with the experimental diets for 12 weeks

Other growth performance, metabolic efficiency and nutrient utilization parameters such as IBM, FBM, SGR, FCR, PER, PPV and ALC followed the same trend. Higher values of these growth performance, metabolic efficiency and nutrient utilization parameters were observed in the RP groups as compared to the AFPP groups except in the INT where low values were observed in the RP groups than in the AFPP groups. Statistically, there was a significant difference in most of the parameters ($p < 0.05$) among the different groups in different experimental treatment in the growth performance, metabolic efficiency and nutrient utilization parameters. The onset of the diseases decreased all the growth performance, metabolic efficiency and nutrient utilization parameters investigated (Table 2). Slightly high SGR was observed in the NI-RP group (13.24 ± 0.11) than in the NI-AFPP group (11.60 ± 0.23). However, in the infected treated (IT) groups, both the RP and AFPP groups had almost the same SGR. Although both the INT groups had a negative SGR, the negativity was more in the AFPP group than the RP group.

211 Table 2: Growth performance and nutrient utilization in BALB/C mice infected with *L. major* fed with experimental diets
 212 of 12 weeks

Parameter	Rat Pellets (RP)			Annonaceae Fruit Pulp Pellets (AFPP)		
	NI-RP	INT-RP	IT-RP	NI-AFPP	INT-AFPP	IT-AFPP
IBM	19.89 ^a ±0.11	20.18 ^b ±0.11	20.18 ^b ±0.11	19.99 ^a ±0.19	19.76 ^a ±0.07	19.93 ^a ±0.13
FBM	31.02 ^b ±0.21	13.18 ^d ±0.13	25.68 ^a ±0.10	29.74 ^b ±0.12	11.71 ^c ±0.13	25.38 ^a ±0.19
BMG	56.10 ^c ±1.49	-34.62 ^a ±0.79	27.31 ^b ±0.66	49.11 ^e ±1.39	-40.71 ^d ±0.68	27.56 ^b ±1.37
SGR	13.24 ^c ±0.30	-8.34 ^a ±0.22	6.54 ^b ±0.14	11.60 ^d ±0.23	-9.58 ^a ±0.17	6.49 ^b ±0.29
FCR	3.05 ^b ±0.07	-4.93 ^c ±0.11	6.28 ^a ±0.18	3.51 ^b ±0.10	-4.16 ^c ±0.07	6.56 ^a ±0.32
PER	0.33 ^a ±0.01	-0.21 ^b ±0.01	0.16 ^c ±0.04	0.29 ^a ±0.01	-0.24 ^b ±0.004	0.16 ^c ±0.01
PPV	0.66 ^a ±0.06	0.02 ^b ±0.01	0.46 ^a ±0.16	0.46±0.01	0.01 ^b ±0.04	0.46±0.01
ALC	52.67 ^a ±3.31	22.17 ^b ±2.60	40.17 ^c ±6.60	49.36 ^a ±4.49	18.7 ^d ±6.60	38.70 ^e ±6.60
Mean values (n=3) ± SEM. Values appended by different small letters within a row are significantly different (P < 0.05).						

214 Haematological changes

215 Hematological changes are presented in Table 3. A decrease in Hemoglobin (Hb) below
216 the normal range was observed in all experimental groups except in NI groups. Red
217 Blood Cells (RBC), Hematocrit (HCT) or Packed Cell Volume (PCV), Mean Capsular
218 Volume (MCV), Mean Corpuscular Haemoglobin (MCH) and Mean Corpuscular
219 Hemoglobin Concentration (MCHC) decrease in all the experimental groups. However,
220 a significant decrease ($p < 0.05$) in Hb was observed in the INT groups as compared to
221 the NI groups. After treatment with pentostam in the IT groups, Hb increased was
222 significant ($p < 0.05$), although it was still below the physiological range. The RBC,
223 HCT, MCV, MCH and MCHC followed the same trend as Hb. Total white blood cells
224 (WBC) or total leucocytes count (TLC) increased significantly ($p < 0.05$) in infected
225 groups compared to NI groups. Differentially, Leukocytes, platelets (PLT) and
226 leukocyte populations (neutrophils, lymphocytes, eosinophils, monocytes and
227 basophils) increased significantly in the INT groups compared to the NI groups. The
228 increase in these parameters was within normal range in all the experimental groups.
229 However after treatment in the IT groups, these parameters increased returning almost
230 to the values in the NI groups. On comparison, the groups showed at least one
231 significant difference among the averages of Hb ($p=0.001$), HCT ($p=0.05$), MCV
232 ($p=0.001$), MCH ($p=0.001$), RBC ($p=0.001$), leukocytes ($p=0.008$), lymphocytes ($p=0.013$),
233 neutrophils ($p=0.029$), monocytes ($p=0.002$) and basophils ($p=0.005$).

[illegible]

Liver function tests

The function test changes are presented in Table 4. Alanine aminotransferase (ALT) also called Serum glutamic pyruvic transaminase (SGTP), aspartate aminotransferase (AST) also called Serum glutamic oxaloacetate transaminase (SGOT), lactate dehydrogenase (LDH), sorbitol dehydrogenase (SDH) creatine kinase (CK) and bilirubin activity increased significantly in the INT groups compared to the NI groups. After treatment, the activity ALT (SGTP), AST (SGOT), LDH, SDH, CK and bilirubin decreased almost returning to that of NI groups. The total protein, albumin, globulin and ALkP decreased significantly in INT animals compare to the NI animals. Most of the liver function tests were found to be within normal range in all IT and NI groups. A significant increase in ALT (SGTP) beyond the physiological range (183.49 ± 3.18) was observed in INT-AFPF group. Also a higher activity of AST (SGOT) beyond the physiological range (218.20 ± 3.56 and 184.87 ± 5.99) was observed in INT-RP and INT-AFPF groups respectively. The activity of amylase increased beyond the physiological range whereas that of LDH decreases below the range in NI-RP (24.33 ± 0.67), NI-AFPF (17.90 ± 0.69) groups but increased significantly beyond the physiological range (40.17 ± 0.29) in the INT-AFPF group. After comparison, there was at least one significant difference among the averages of all biochemical parameters used to ascertain liver function of the mice.

253 Table 4: Liver functions in BALB/C mice infected with *L. Major* fed with RP and AFPP for 12 weeks

Parameter	Rat Pellets (RP)			Annonaceae Fruit Pulp Pellets (AFPP)			Ref. Range
	NI-RP	INTC-RP	IT-RP	NI-AFPP	INT-AFPP	IT-AFPP	
Glucose (mg/L)	122.71±2.12	123.87±1.95	124.37±1.91	126.17±0.45	124.27±0.65	125.20±0.41	106.00 – 278.00
T-Bilirubin (g/ dL)	9.02±0.17	11.18±0.10	9.52±0.29	10.20±0.06	12.68±0.35	9.68±0.07	7.12 -12.05
C-Bilirubin (g/ dL)	2.57±0.11	3.43±0.05	2.57±0.11	3.07±0.26	4.23±0.35	2.80±0.08	0.00 – 2.00
U-Bilirubin (g/ dL)	3.67±0.17	4.83±0.13	3.17±0.13	3.83±0.16	4.33±0.20	3.00±0.11	0.00 – 3.00
ALT (IU/L)	77.49±2.40	115.99±1.02	98.66±1.31	83.49±3.18	183.49±3.18	81.82±2.86	41.00 – 131.00
AST (IU/L)	119.87±2.16	218.20±3.56	111.53±0.69	134.87±5.86	184.87±5.99	129.87±5.66	28.00 – 191.00
GGT (IU/L)	25.43±0.55	28.77±0.48	24.60±0.46	23.77±0.60	31.43±0.56	23.77±0.60	20.00 – 40.00
ALKP (IU/L)	132.36±1.53	127.36±1.33	134.03±1.21	157.36±4.40	140.70±1.70	155.70±3.03	118.00 – 187.00
T-Protein (g/ dL)	51.77±0.90	45.94±0.04	47.12±0.33	42.95±0.51	35.02±0.51	45.27±0.51	43.00 – 70.00
Albumin (g/ dL)	26.46±0.49	23.34±0.58	35.27±0.51	24.50±0.32	22.27±0.40	26.35±0.14	27.00 – 46.00
Globulin (g/ dL)	34.56±0.64	25.22±0.47	35.34±0.35	26.34±0.38	23.18±0.42	26.68±0.10	27.00 – 42.00
Alb/Glob ratio	0.77 ^d ±0.02	0.93 ^d ±0.03	1.98 ^{bc} ±0.05	1.10 ^a ±0.02	1.06 ^a ±0.02	2.12 ^b ±0.04	0.00 – 1.09
CPK (IU/L)	3.48 ^b ±0.22	4.02 ^c ±0.08	1.79 ^a ±6.86	1.79 ^a ±0.07	2.46 ^d ±0.05	1.60 ^a ±0.04	2.50 – 3.70
Amylase (U/L)	796.97 ^b ±17.47	1580.31 ^a ±54.65	896.97 ^d ±7.90	813.64 ^e ±23.68	1413.64 ^e ±36.61	896.97 ^f ±10.69	210.43 -323.57
LDH (IU/L)	24.33 ^a ±0.67	34.66 ^b ±0.39	27.17 ^c ±0.68	17.90 ^d ±0.69	40.17 ^e ±0.29	24.83 ^a ±0.52	26.8 – 34.00
SDH (IU/L)	29.15 ^a ±0.90	34.50 ^c ±0.53	31.16 ^{ab} ±0.54	23.83 ^d ±0.39	43.67 ^e ±0.44	29.17 ^a ±0.25	27.00 - 37.00
Mean values (n=3) ± SEM. Values appended by different small letters within a row are significantly different (P < 0.05).							

Kidney function tests and lipid profiles

The kidney function tests and lipid profile changes are presented in Table 5. Creatinine, urea and blood urea nitrogen (BUN) increased significantly ($p<0.05$) in the INT group compared to NI groups. In all the IT groups, creatinine, blood Urea and BUN levels decreased significantly ($p<0.05$) compared to the INT groups. Creatinine levels beyond the physiological range were observed in all BALB/C mice groups while BUN decreased in NI-AFPP (25.16 ± 0.11) and IT-AFPP (23.82 ± 0.10) groups. However, the increased urea levels in all the BALB/C mice groups were found to be within normal range. Serum sodium, chloride, potassium and phosphorous increased significantly in the INT groups compared to the NI groups although values remained within the physiological ranges. After treatment, a decrease in these ions was observed in the IT groups. The cholesterol, triacylglycerides (TGs), high density lipoproteins (HDL), low density lipoprotein (LDL) and cholesterol/HDL ratio increased significantly ($p<0.05$) in the INT groups in NI groups indicative of liver damage and development of cardiovascular diseases. All lipid profile values were within the physiological range except HDL in INT-RP (2.73 ± 0.11), IT-RP (1.23 ± 0.07), INT-AFPP (3.03 ± 0.06) and IT-AFPP (1.90 ± 0.13) groups suggesting a relationship between leishmaniasis and cardiovascular diseases. Statistically, there was no significant difference between lipid profiles of NI-RP and NI-AFPP (0.09) and INT-RP and INT-AFPP ($p=0.08$) but there was significant differences between NI and INT ($p=0.003$) and INT and IT ($p=0.001$) groups.

275 Table 5: Kidney functions and lipid profile in BALB/C mice infected with *L. Major* fed with RP and AFPP for 12 weeks

Parameter	Rat pellets (RP)			Annonaceae Fruit Pulp Pellets (AFPP)			Ref. Range
	NI-RP	INT-RP	IT-RP	NI-AFPP	INT-AFPP	IT-AFPP	
Creatinine (mg/L)	2.57±0.16	3.32±0.15	2.49±0.19	2.12±0.11	2.87±0.25	2.48±0.10	0.50 - 0.80
Blood Urea Nitrogen	26.66±0.16	31.32±0.15	27.49±0.19	25.16±0.11	27.66±0.25	23.82±0.10	7.00 - 26.00
Urea (mg/L)	7.88±0.11	9.11±0.11	7.38±0.18	6.88±0.22	10.03±0.06	6.88±0.15	7.00 - 26.00
Sodium (meq/L)	143.48±0.74	147.13±1.04	142.96±0.46	135.63±1.33	143.63±0.93	137.29±0.78	125.30 - 187.40
Chloride (meq/L)	116.23±0.82	121.90±1.49	117.40±0.64	121.23±1.10	131.23±1.56	122.90±1.91	109.6 - 138.80
Potassium (meq/L)	6.16±0.13	6.49±0.17	5.49±0.19	7.14±0.06	6.82±0.17	6.32±0.21	7.30 - 12.07
K/Na Ratio	0.04±0.00	0.04±0.00	0.04±0.00	0.05±0.00	0.05±0.00	0.05±0.00	0.06 - 0.09
Phosphorus (meq/L)	3.77±0.09	3.72±0.09	3.65±0.08	4.22±0.16	5.22±0.19	4.22±0.02	8.20 - 14.70
Calcium (mg/dL)	2.59±0.06	2.30±0.03	2.74±0.04	2.76±0.11	2.68±0.09	3.43±0.06	9.40 - 12.70
Cholesterol (g/dL)	3.75±0.18	7.13±0.16	4.81±0.16	4.22±0.14	7.13±0.16	5.31±0.26	11.10 - 24.60
Triglycerides (g/dL)	1.16±0.04	2.48±0.05	1.68±0.05	1.34±0.06	3.11±0.08	1.81±0.03	1.050 - 5.35
HDL Cholesterol (g/dL)	0.77±0.02	2.73±0.11	1.23±0.07	1.00±0.09	3.03±0.06	1.90±0.13	0.40 - 1.00
LDL Cholesterol (g/dL)	1.34±0.11	3.10±0.09	2.12±0.15	1.51±0.11	3.10±0.09	2.12±0.15	
Cholesterol/ HDL Ratio	4.23±0.08	2.97±0.39	4.37±0.07	4.65±0.07	2.97±0.14	4.30±0.13	
Mean values (n=6) ± SEM. Values appended by different small letters within a row are significantly different (P < 0.05).							

276

277

Lesion size and liver and splenic length, size and parasite burden

The lesion sizes (mm²) for the INT and IT groups were determined and are presented in Figure 2. Generally, there was an increase in the size of the lesions with the progression of the disease. After the initiation of treatment on the IT groups, the lesion sizes decreased. However, the rate of decrease of the size of the lesions was higher in the IT-RP group than in the IT-AFPP group (Figure 2). The lesion sizes of the INT groups increased significantly until the end of the experiment, with the INT-AFPP having larger lesion sizes than the INT-RP group. The hepatic and splenic length (mm) and size (mg) for the NI, INT and IT groups (Table 6) determined at the end of the experiment varied significantly ($p < 0.05$). The relative mean weight and length of the liver and the spleen increased with increasing the time of infection. A significant differences ($p < 0.05$) was observed between the weight and length of infected (both IT and INT) and non-infected (NI) BALB/C mice groups. However, no significant difference ($P > 0.05$) was observed between NI-RP and NI-AFPP. The same trend was observed between ITC-RP and IT-AFPP.

The parasite density of *L. major* amastigotes in liver and spleen for infected not treated (INT) and IT groups of BALB/C mice (Table 6) determined after the experimental period (12 weeks) showed significant variations. In all the IT groups of BALB/C mice, parasite load decreased significantly ($p < 0.05$) as compared to the INT groups. The reduction in parasite load was significantly ($p < 0.05$) more in the IT-RP group as compared to the IT-AFPP group. After the experimental period (12 weeks), hepatomegaly and splenomegaly was observed on the INT groups of BALB/C mice. Impression smears of liver and spleen on the INT groups of BALB/C determined at the end of the experimental period (12 weeks) showed dissemination of *L. major* amastigotes inside and outside macrophages.




Figure 2: Lesion sizes of BALB/C mice infected with *L. major* and feed with RP and AFPP on 12 weeks.

309 **DISCUSSION**

310 The results of the present investigation demonstrate that AFPP may be a good dietary
311 protein source for BALB/C mice feed because growth performance and feed utilization
312 of Annonaceae group was similar to control group (RP fed group). No significant
313 difference in growth performance (BMG, SGR and MGR) and feed utilization
314 parameters (FCR, PER and ALC) show that digestion and absorption of nutrients from
315 RP and AFPP were similar. Nutrient utilization values observed in this study signify
316 excellent utilization of the diets. Growth performance of RP fed group was similar to
317 AFPP group indicating that nutrients and energy availability from the AFPP for the
318 protein synthesis were similar to animal protein (RP). Further, the growth performance
319 and feed utilization parameter of the infected and treated groups feed with RP and
320 AFPP returned to normal. This shows that AFPP can be used as an alternative source of
321 feed for BALB/C mice.

322 In terms of haematological changes, lesion size and organ length, size functioning and
323 parasite burden, *Leishmania major* is an aetiological agent of cutaneous leishmaniasis,
324 which is a parasite of the skin on humans. However, in BALB/C mice, it attacks visceral
325 organs in addition to the local lesion at the point of inoculation [25]. The parasites can
326 be deposited in major visceral organs involved in the synthesis of major macromolecule
327 after the digestion, absorption and transport. Nonetheless, BALB/C mice model was
328 chosen for this study as a suitable model for the investigation of interaction between
329 malnutrition and leishmaniasis. In earlier studies, malnourishment contributed to
330 higher parasite loads found in the blood, skin, bone marrow, lymph node, liver and
331 spleen and favoured the development of leishmaniasis [6]. Based on these observations,
332 this model provides an excellent opportunity to elucidate the factors implicated in
333 severe malnutrition. Moreover, as it has been shown in this work, the model can be
334 used for further investigation on the relationship between severe malnutrition and
335 leishmaniasis

336 Malnutrition decreases weight-for-age in proportion to the level of dietary protein
337 deficiency [11]. Cure of the disease depends upon the development of an effective
338 immune response that activates macrophages [26], which can be promoted and
339 enhanced by good nutrition. During Leishmaniasis, there is a profound
340 immunosuppression in the host that promotes the survival of parasites [1]. In this
341 study, it is revealed that treatment of infected animals coupled with good nutrition
342 brought the levels of most biological parameters such as Hb and total leucocytes count
343 (TLC) to normal range as compared to the abnormal levels in infected animals.

344 Kidney function tests include estimation of urea, blood urea nitrogen and creatinine.
345 Renal abnormalities caused by *Leishmania* have been well documented in experimental
346 animal studies and are comprised of interstitial and glomerular abnormalities [27].
347 Sodium, chloride and potassium and K/Na ratio values give an indication of
348 electrolyte/water balance, whereas high levels of calcium implies thyroid or
349 parathyroid, intestine, pancreas, kidney and bone metastasis. Although in the current

study there were elevated levels of sodium, chloride and potassium and K/Na ratio, the role of infection caused by *L. major* on damage of these organs causing the elevated levels of these parameter is yet to be established. The reason may be that these parameters can vary with mouse strain/stock, age, sex; blood sampling method, environmental conditions pathogen status and the laboratory as well as nutrition [28].

The task of establishing a range of reference values for rodents is very difficult. This is because of very many variables, such as gender, age, genetic variation, diet and environmental conditions to which these animals are subjected, must be considered [29]. In this study, mice of the same sex and age infected with the same strain of *L. major* were used. Therefore, they may provide a useful starting point to investigate the effect of leishmaniasis and nutrition on hematological parameters. Since significant variation of biological parameters may occur between individual mice strain, stock, laboratories and method of sampling, individual laboratories should establish normal reference values for their facility [30].

The most important organ concerned with majority of biochemical activities in the human body is the liver. Since it has a great capacity to detoxify toxic substances and synthesize useful biological molecule, damage inflicted by hepatotoxic agents is of grave consequence. To assess the damage caused to the liver, activities of enzymes alanine aminotransferase (ALT) also called Serum glutamic pyruvic transaminase (SGTP), Aspartate aminotransferase (AST) also called Serum glutamic oxaloacetate transaminase (SGOT), Alkaline phosphatase (ALP) and sorbitol dehydrogenase (SDH) and the concentration of bilirubin were measured. The increased level of ALT, AST, ALP, and bilirubin is conventional indicator of liver injury. In leishmaniasis liver damage due to high parasitic load in infected groups resulting in elevated levels of ALT (SGTP), AST (SGOT), ALP and SDH an increased levels of bilirubin was observed in this study. Albumin is used to detect liver damages whereas globulin and total protein content is used to detect and immunoglobulin status, a key indicator in fighting of infections in organisms which is affected by nutrition. The stabilization of serum bilirubin, ALT (SGPT), AST (SGOT), and ALP levels is a clear indication of the improvement in the functional status of the liver cells [31]. The increase in AST (SGOT) and ALT (SGPT) levels has been reported in the visceral leishmaniasis (VL) patients [32] which is depicted in this study for cutaneous leishmaniasis (CL) using mice model. In the current study, there was no significant elevation in the levels of liver function tests in all the treated groups as compared to infected controls.

The mononuclear phagocyte system (spleen, liver, bone marrow, intestinal mucosa and mesenteric lymph nodes) is the normal habitat of leishmania parasite [33]. However, the parasite may be found in endothelial cells of the kidneys, suprarenal capsules, lungs, meninges and in cerebrospinal fluid [34]. In leishmaniasis there can be possibilities of multiple organ damage as indicated by elevated levels of non-specific markers such as lactate dehydrogenase (LDH) indicating possible damage of liver, heart, skeletal

muscles and lungs and creatine kinase (CK) commonly elevated in heart and skeletal muscle damage and muscular dystrophies.

These effects are consequences of the stimulation of the immune system by *L. major*, which promotes the inflammatory components of atherosclerosis, which are primarily the parasite-activated macrophages [35]. Several studies suggest that pathogenic bacteria, viruses and protozoa, contribute to the atherogenesis process [36, 37, 38, 39, 40]. The increased inflammation caused by these pathogens promotes macrophage activation and migration to the atheroprone site. Alternatively, proatherogenic status may be attributed to the systemic oxidative stress induced by infection, which enhances lipoprotein or endothelium oxidation. Pathogens involved in atherosclerosis development usually induce a systemic infection instead of a localized infection such as an *L. major* infection. However, even localized infections, such as odontologic ones, may be associated with the development of atherosclerosis [41].

In this study, microscopic examination of stained impression smear of liver and spleen showed the density of amastigotes in two organs that demonstrated the pathological effect of parasite. The results of this study have demonstrated that the infected mice show the hepatosplenomegaly sign of pathological effect of *L. major* promastigote in the infected mice. The weight of liver was increased with increasing days of infection in the INT mice compared with NI mice on 12 weeks. For both the liver and spleen, NT mice had the highest LDU. Significantly ($p < 0.05$) higher parasite load in the liver and spleen occurred in NT mice and the lowest LDU occurred in the IT mice. There were no significant difference between the following groups, NI-RP and NI-AFPP ($p > 0.05$), IT-RP and IT-AFPP ($p > 0.05$), INT-RP and INT-AFPP ($p > 0.05$), indicating that there is no difference between the two feeds.

CONCLUSION AND RECOMMENDATION

From the study it can be revealed that there was no difference in growth performance, metabolic efficiency and nutrients utilization parameters between RP and AFPP among the same treatment of BALB/C mice. This suggests that Annonaceae fruit pulp can be used as an alternative source of raw material in the manufacture of animal feeds. It has been observed that the data obtained from present investigation related to different growth performance, metabolic efficiency and nutrients utilization make Annonaceae fruit a good candidate for manufacture of nutraceuticals. Thus, there is a need to study these parameters in the another animal model to validate the findings of this study for utilization of Annonaceae fruits in the nutraceutical manufacturing industry.

REFERENCE

1. Serafim, T.D., Malafaia, G., Silva, M.E., Pedrosa, M.L, Rezende, S.A. (2010). Immune response to *Leishmania (Leishmania) chagasi* infection is reduced in malnourished BALB/c mice *Mem Inst Oswaldo Cruz*, Rio de Janeiro 105(6): 811-817.

2. Badaró R, Jones TC, Lorenço R, Cerf BJ, Sampaio D, Carvalho EM, Rocha H, Teixeira R, Johnson WD (1986) A prospective study of visceral leishmaniasis in an endemic area of Brazil. *J Infect Dis* 154: 639-649.
3. Cerf BJ, Jones TC, Badaró R, Sampaio D, Teixeira R, Johnson WD JR (1987) Malnutrition as a risk factor for severe visceral leishmaniasis. *J Infect Dis* 156: 1030-1033.
4. Maciel BL, Lacerda HG, Queiroz JW, Galvão J, Pontes NN, Dimenstein R, McGowan SE, Pedrosa LF, Jerônimo SM (2008) Association of nutritional status with the response to infection with *Leishmania chagasi*. *Am J Trop Med Hyg* 79: 591-598.
5. Blössner M, Onis M (2005) Malnutrition: Quantifying the health impact at national and local levels, World Health Organization, Geneva, pg 51.
6. Carrillo E, Jimenez M, Sanchez C, Cunha J, Martins CM, Seva AP, Moreno J (2014) Protein Malnutrition Impairs the Immune Response and Influences the Severity of Infection in a Hamster Model of Chronic Visceral Leishmaniasis. *PLoS ONE* 9(2): e89412. Doi: 10.1371/journal.pone.0089412
7. WHO (2002) Reducing risks, promoting healthy life, The World Health Report, chapter 4. Available: <http://www.who.int/whr/2002/en/>. Accessed 2017 Jan 27.
8. Gross RL, Newberne PM (1980) Role of nutrition in immunologic function. *Physiol Rev* 60: 188-302.
9. Malafaia G (2009) Protein-energy malnutrition as a risk factor for visceral leishmaniasis: a review. *Parasite Immunol* 31: 587-596.
10. Malafaia G, Serafim TD, Silva ME, Pedrosa ML, Rezende SA (2009) Protein-energy malnutrition decreases immune response to *Leishmania chagasi* vaccine in BALB/c mice. *Parasite Immunol* 31: 41-49.
11. Anstead GM, Chandrasekar B, Zhao W, Yang J, Perez LE, Melby PC (2001) Malnutrition alters the innate immune response and increases early visceralization following *Leishmania donovani* infection. *Infect Immun* 69: 4709-4718.
12. Moghadamtousi SZ, Goh BH, Chan CK, Shabab T, Kadir HA (2013) Biological activities and phytochemicals of *Swietenia macrophylla* king. *Molecules* 18: 10465-10483.
13. Moghadamtousi SZ, Kamarudin, MNA, Cha, CK, Goh BH, Kadir HA (2014) Phytochemistry and biology of *Loranthus parasiticus* merr, a commonly used herbal medicine. *Am. J. Chin. Med.* 42: 23-35.
14. Chimbevo ML, Malala JB, Anjili CO, Orwa J, Mibei EK, Ndeti CM, Muchiri FW, Sifuna PO, Oginga FO, Otundo DO, and Karanja SM (2017) Annonaceae Fruits

- 465 Growing in Coast Region of Kenya as an Alternative Source of Dietary Carotenoids.
466 International Journal of Food Science and Biotechnology 2(5): 114-120
- 467 15. Rupprecht JK, Hui YH, McLaughlin JL (1990). Annonaceous acetogenins: A review.
468 *J. Nat. Prod* **53**: 237-278.
- 469 16. Pinto AC, de Q, Cordeiro de Andrade SRM, Ferreira FR, Filgueiras HA, de C., Alves
470 RE, Kinpara DI (2005) *Annona species*. International Centre for Underutilised Crops,
471 University of Southampton, Southampton, UK. ISBN 0854327851.
- 472 17. Maundu PM, Ngugi GW, Kabuye CHS (1999) Traditional Food plants of Kenya.
473 KENRIK, National museums of Kenya, Nairobi, Kenya.
- 474 18. AOAC (2000) Official Methods of Analysis. Association of Official Analytical
475 Chemists, Washington DC.
- 476 19. Carvalho LJ, Ferreira-da-Cruz M, Daniel-Ribeiro FM, Pelajo-Machado CT, Henrique
477 LL (2006) *P. berghei* ANKA infection induces thymocyte apoptosis and thymocyte
478 depletion in CBA mice. *Memórias do Instituto Oswaldo Cruz* 101: 5-9
- 479 20. Schnur LE, Zuckerman A, Montillio B (1973) Dissemination of leishmaniasis to the
480 organs of Syrian hamsters following intrasplenic inoculation of promastigotes. *Exp.*
481 *Parasitol.* 34: 432-447.
- 482 21. Nolan TJ, Farrell JP (1987) Experimental infections of the multimammate rat (*Mastomys*
483 *natalensis*) with *L. donovani* and *L. major*. *American Journal of Tropical Medicine*
484 *and Hygiene*, 36(2): 264-269.
- 485 22. Kumar V, Akinleye AO, Makkar HPS, Angulo-Escalante MA, Becker K (2012)
486 Growth performance and metabolic efficiency in Nile tilapia (*Oreochromis niloticus*
487 L.) fed on a diet containing *Jatropha platyphylla* kernel meal as a protein source. *J*
488 *Anim Physiol Anim Nutr* 96(1): 37-46.
- 489 23. Bradley DJ, Kirkley J (1977) Regulations of *Leishmania* population within the host.
490 The variable course of *Leishmania donovani* infection in mice. *Clinical Experimental*
491 *Immunology*. 30: 119-129
- 492 24. Payne RW, Murray DA, Harding SA, Baird DB, Soutar DM (2011) An introduction
493 to GenStat for Windows, 14th Ed. VSN International, Hemel Hempstead, UK.
- 494 25. Makwali1 JA, Wanjala1 FME, Kaburi JC, Ingonga J, Byrum WW, Anjili CO (2012)
495 Combination and monotherapy of *Leishmania major* infection in BALB/c mice using
496 plant extracts and herbicides. *Vector Borne Dis* 49: 123-130
- 497 26. Kaur S, Kaur G, Sachdeva H, Kaur J (2013) *In Vivo* Evaluation of the Antileishmanial
498 Activity of Two Immunomodulatory Plants, *Embllica Officinalis* And *Azadirachta Indica*
499 In Balb/C Mice. *International Journal of Ayurvedic and Herbal Medicine* 3(1): 1066-
500 1079

- 501 27. Salgado FN, Ferreira TM, Costa JM (2003) Involvement of the renal function in
502 patients with visceral leishmaniasis (kala-azar). Rev Soc Bras Med Trop 36(2): 217-21.
- 503 28. NAS (1995) Nutritional requirements of laboratory animals. 4th Revised edition.
504 National academy press. Washington DC
- 505 29. Santos EW, Oliveira DC, Hastreiter A, Silva GB, Beltran JSO, Tsujita M, Crisma AR,
506 Neves SMP, Fock RA, Borelli P (2016) Hematological and biochemical reference
507 values for C57BL/6, Swiss Webster and BALB/c mice. Braz. J. Vet. Res. Anim. Sci
508 53(2): 138-145.
- 509 30. Sackow MA, Danneman P, Brayton C (2005) The laboratory Mouse
- 510 31. Achliya GS, Wadodkar SG, Dorle AK (2004) Evaluation of hepatoprotective effect of
511 Amalkadi Ghrita against carbon tetrachloride-induced hepatic damage in rats. J
512 Ethnopharmacol 90 (2-3): 229-232.
- 513 32. Mathur P, Samantaray JC, Samanta P (2008) High prevalence of functional liver
514 derangement in visceral leishmaniasis at an Indian tertiary care centre. Clin
515 Gastroenterol Hepatol 6(10): 70-1172.
- 516 33. Jarallah HM (2016) Pathological Effects of *Leishmania Donovanii* Promastigotes on
517 Liver and Spleen of Experimentally Infected BALB/C Mice Medical Journal of
518 Babylon. 13 (1): 134 – 140.
- 519 34. Beaver PC, June RC, Cupp EW (1984) Clinical Parasitology. 9th Ed. Lea and Febiger;
520 Philadelphia 70-77.
- 521 35. Fernandes LR, Ribeir, ACC, Segatt, M, Santos LFFF, Amaral J, Portugal LR, Leite JIA
522 (2013) Leishmania major Self-Limited Infection Increases Blood Cholesterol and
523 Promotes Atherosclerosis Development. Cholesterol Article ID 754580.
- 524 36. Murray LJ, Bamford KB, Reilly DPJO, McCrum EE, Evans AE (1995) Helicobacter
525 pylori infection: relation with cardio- vascular risk factors, ischaemic heart disease,
526 and social class". British Heart Journal 74 (5): 497–501.
- 527 37. Doenhoff MJ, Stanley RG, Griffiths K, and Jackson CL (2002) An anti-atherogenic
528 effect of Schistosoma mansoni infections in mice associated with a parasite-induced
529 lowering of blood total cholesterol. Parasitology 125 (5): 415–421,

- 530 38. Burnett MS, Durrani S, Stabile E, et al., (2004) "Murine cytomega-
531 loves virus infection
532 increases aortic expression of proatherosclerotic genes," *Circulation*, 109 (7): 893–
897.
- 533 39. Portugal LR, Fernandes LR, Cesaretal GC (2004) "Infectionwith *Toxoplasma gondii*
534 increases atherosclerotic lesion in ApoE- deficient mice," *Infection and Immunity*.
535 72: 3571– 3576.
- 536 40. Mussa FF, Chai H, Wang X, Yao Q, Lumsden AB, Chen C (2006) *Chlamydia*
537 *pneumoniae* and vascular disease: "an update". *Journal of Vascular Surgery* 43:
538 1301–1307.
- 539 41. Meurman H, Sanz M, Janket SJ (2004) "Oral health, athero- sclerotic, and
540 cardiovascular disease," *Critical Reviews in Oral Biology and Medicine*. 15: 403–413.
- 541