

Review Paper

Past and Current Findings in Antimalarial Drug Resistance Molecular Markers in Endemic Areas of Africa

Cynthia Ufuoma Adjekukor^{1*}

¹Covenant University Bioinformatics Research (CUBRe), Covenant University, Ota, Ogun State, Nigeria.

ABSTRACT

Antimalarial drug resistance is the major challenge in the treatment of malaria all over the world. *Plasmodium* species are the parasite that causes malaria. *Plasmodium falciparum* is the most prevalent specie found in sub-Saharan Africa that records the highest infections and death caused by malaria worldwide. Resistance to *P. falciparum* is caused by mutations in some target genes of the parasite, which includes *Plasmodium falciparum*: Na⁺/H⁺ exchanger (*Pfnhe-1*), chloroquine resistance transporter (*Pfcr1*), dihydropteroate synthase (*Pfdhps*), dihydrofolate reductase (*Pfdhfr*), multi drug resistance 1 gene (*Pfmdr1*), cytochrome b, multidrug resistance-associated protein 1 (*Pfmrp1*), cg2 (*Pfcg2*), Ca²⁺-ATPase (*PfATPase6*) and kelch 13 gene. Most of these mutations are single nucleotide polymorphisms and has led to the decrease in susceptibility of some drugs like chloroquine, quinine, mefloquine, amodiaquine, sulphadoxine/pyrimethamine, lumefantrine and artemisinins in the treatment of malaria. The aim of this review was to survey on the existing antimalarial drug resistance in endemic areas of Africa and suggests a way forward in combating drug resistant malaria in this region.

Keywords: Malaria; *Plasmodium falciparum*; antimalarial drug resistance; Africa

1. INTRODUCTION

Malaria remains one of the most deadly diseases responsible for high rate of morbidity or mortality in tropical regions of the world. In 2010, the World Health Organisation (WHO) estimated that malaria causes over 240 million illness and 0.86 million deaths annually [1]. This disease is caused by the vector; Anopheline mosquito that transmits the protozoan parasite *Plasmodium* [2]. *Plasmodium falciparum* is the most deadly species of *Plasmodium* and it is the major cause of drug resistant malaria in Sub-Saharan Africa, Southeast Asia and the Amazon region [3]. While the *Plasmodium vivax*, which causes about 25-40% of deaths, is mainly found in South America and Asian countries and it is responsible for relapsing malaria [1]. The emergence of drug resistant parasites have threatened the efficacy of the available drugs in the treatment of malaria especially in sub-Saharan Africa where *P. falciparum* is responsible for 90% of deaths, thus increasing the morbidity and mortality rates of malaria [4]. Molecular markers can be used to monitor antimalarial drug resistant parasites that are caused by genetic mutation in the target gene of the parasites [5]. This review describes the different molecular markers and state of drug resistant malaria in Africa and the way forward to combat the issue of antimalarial drug resistant.

2. LITERATURE SEARCH

A comprehensive literature search was done about prevalence of antimalarial drug resistance and molecular markers of antimalarial drug resistance of *P. falciparum* isolated from African patients and various studies done in Africa by using PubMed (<http://www.ncbi.nlm.nih.gov>) and Google Scholar (www.scholar.google.com).

3 MOLECULAR MARKERS OF ANTIMALARIAL DRUG RESISTANCE

Artemisinin-based combination therapy (ACT) is now recommended as the first-line drug for the treatment of uncomplicated malaria because of the prevalence of resistant parasites [6]. The replacements of ACTs with ineffective failing treatments (chloroquine, sulphadoxine-pyrimethamine) and also artesunate with quinine specifically for treating severe malaria have reduced the morbidity and mortality rate of malaria [7]. Although, ACTs is the new remedy for the treatment of malaria caused by *Plasmodium falciparum*, but resistance to these drug has led to an increase in the rate of treatment failure and this is now prevalent and expanding in Southeast Asia [8]. Some genes have been identified and proven to be involved in *P. falciparum* antimalarial drug resistance which include: the genes encoding the *Plasmodium falciparum* Na⁺/H⁺ exchanger (*Pfnhe-1*), *Plasmodium falciparum* chloroquine resistance transporter (*Pfcr1*), *Plasmodium falciparum* dihydropteroate synthase (*Pfdhps*), *Plasmodium falciparum* dihydrofolate reductase (*Pfdhfr*), *Plasmodium falciparum* multi drug resistance 1 gene (*Pfmdr1*), cytochrome b, [9] *Plasmodium falciparum* multidrug resistance-associated protein 1 (*Pfmrp1*), *Plasmodium falciparum* cg2 (*Pfcg2*), *Plasmodium falciparum* Ca²⁺-ATPase (*PfATP6*) and kelch 13 gene. Single Nucleotide Polymorphisms (SNPs) in these markers are very important in monitoring or determining new drug resistant patterns [10].

Artemisinin Combination Therapies (ACTs)	Trade Names
Amodiaquine + Artesunate*	Efonrex, Artesmodia, Camosunate
Artesunate* + Sulphadoxine/pyrimethamine	Amalar plus, Tesunate SP, CALSUNATE SP
Artesunate* + Mefloquine	Larinate-MF, Mefliamplus, Artequin
Pyronaridine + Artesunate*	Pyramax
Artemether* + Lumefantrine	Coartem, Lumartem, Lumapil
Dihydroartemisinin* + Piperaquine	Pymal, Pipart, Terocan
Artemisinin* + Piperaquine	Artequick

Table 1 Artemisinin Combination Therapies (ACTs) and examples of their trade names

Note the * ones are the artemisinin derivative drugs

3.1 *Plasmodium falciparum* Na⁺/H⁺ exchanger (*Pfnhe-1*)

Pfnhe is located on chromosome 13 fragments and have additional loci at chromosome 9 in the parasite. It is the second biggest eukaryotic Na⁺/H⁺ exchanger after that of the apicomplexan *T. gondii* [11]. *Pfnhe-1* is a 226 KDa protein that is predicted to have 12 trans membrane segment with a signal peptide cleavage site and functions to intensify the cytosolic pH (CytpH) and to make up for acidosis that is caused by anaerobic glycolysis [12]. The genetic changes that is associated with changes in cytosolic pH, which is linked to chloroquine/quinine resistance still remains unclear [11].

Pfnhe-1 has been described to have three point polymorphisms at codon 790, 894 and 950 with three different repeat sequence of microsatellite variants MsR1, ms3580, and ms4760 that may be responsible for drug susceptibility or resistance [13]. The ms4760 is the most studied in Africa due to the strain selectivity in geographical areas and an increase/decrease in amino acid motif DNNND or/and an increase/decrease in that of NHNDNHNNDDDD motifs have been linked to quinine susceptibility in some studies. A study done in Republic of Congo that involved 74 *Plasmodium falciparum* isolates discovered that the polymorphisms in the number of either the DNNND or NHNDNHNNDDDD were not in connection with quinine susceptibility while another study done with samples from different countries found out that an increase in the DNNND led to an increase in IC₅₀ of quinine while an increase in the NHNDNHNNDDDD repeats led to a decrease in IC₅₀ of quinine [14, 15]. In addition, some authors discovered that in Uganda and Kenya coast that there was a decrease in quinine susceptibility in the presence of *Pfmdr1* mutations combined with two DNNND repeats, while in

the presence of 1 *Pfmdr1* mutations with 3 DNNND repeats and 2 *Pfmdr1* with 3 DNNND repeats led to the return of quinine susceptibility [16].

3.2 *Plasmodium falciparum* chloroquine resistance transporter (*Pfcr*)

Pfcr is a 48KDa protein that is predicted to have 10 transmembrane spanning domains with 424 amino acids and it is found in the digestive vacuole membrane of the parasite [17]. It is found on chromosome 7 of the *P.falciparum*. When mutated, this transporter protein will lower the volume of accumulating drugs in the digestive vacuole of *P.falciparum* due to enhanced efflux of drugs [18].

Eight point mutations M74I, N75E, K76T, A220S, Q271E, N326S, I356T and R371I have been identified in the chloroquine resistance transporter isolates from both Asia and Africa and in the chloroquine resistance Dd2 strain, however the I356T was not found in some isolates [19]. Although, there are eight point mutations of chloroquine resistance transporter, K76T is the major mutation that is closely linked to increase clinical and treatment failure in the use of chloroquine [5]. There are three haplotypes that are present in codon 72-76 of *Pfcr*, which are the wild type CVMNK and the chloroquine resistant type CVET and SVMNT. Among the three haplotypes present in codon 72-76 of the *Pfcr*, the CVET is the most important mutant haplotypes that is present in Africa [20]. Also, there have been reports that the mutation at codon A220S is associated to chloroquine resistance in African countries, while two specific mutations at codon A144T and 160Y is only linked to chloroquine resistance in Philippines [21].

Some studies done in Africa has linked the parasite clearance rate at *Pfcr* 76 to the age of the host thus suggesting that the age of the host is the best surrogate marker for immunity in resistance to chloroquine and amodiaquine. One of such studies done in Mali of K76T mutation confirmed that the infants had a slightly higher clearance rate than young children, which led to the belief that host immunity plays an important role in clearance of parasites [22]. Another one done in south western part of Nigeria showed that amodiaquine selects the same marker as chloroquine (*Pfcr* 76 and *Pfmdr1* 86), probably due to their similar chemical structure and mode of action and also confirmed the age of the host as the best surrogate marker for immunity in the resistance to amodiaquine in West Africa [23].

3.3 *Plasmodium falciparum* dihydropteroate synthase (*Pfdhps*) and *Plasmodium falciparum* dihydrofolate reductase (*Pfdhfr*)

Sulpha drugs such as sulphadoxine binds *Plasmodium falciparum* dihydropteroate synthase (*Pfdhps*) which is an enzyme in the folate cycle that is important for DNA synthesis while drugs like pyrimethamine binds to *Plasmodium falciparum* dihydrofolate reductase (*Pfdhfr*) which is also an enzyme in the folic acid pathway [24]. The mechanisms for resistance for these two enzymes are that when mutation occurs; it modifies the protein structure and thus reducing its binding affinity for the drugs [24].

WHO recommended sulphadoxine/pyrimethamine (SP) as the Intermittent Preventive treatment (IPTp) for pregnant women and children under age 5 years [25]. *Plasmodium falciparum* resistance to the antifolate drugs is caused by mutations in the dihydropteroate synthase (*Pfdhps*) and dihydrofolate reductase (*Pfdhfr*) gene respectively [26].

A lot of studies have reported the prevalence of *Pfdhps* and *Pfdhfr* in Africa, this section highlighted on few of such studies. A study done in Kombwa lowland and Kisii and Kakamega highlands in Kenya, discovered that more than 80% of the samples had the quintuple mutants (51I, 59R, 108N for *Pfdhfr* and 437G/540E for *Pfdhps*) in this area [5]. Also, another study done on children under age 5 years in southern Rwanda, 75% and 93% of the isolates had the *Pfdhfr* triple and the *Pfdhps* double/triple mutations, and 69% were found with the quintuple/sextuple *Pfdhfr/Pfdhps* mutations that has been linked to high level of sulphadoxine/pyrimethamine resistant, while L164 mutation was absent in all isolates, but in a study done in Uganda from 2003 – 2012, five mutations (*Pfdhfr* N51I, C59R, S108N; and *Pfdhps* A437G, K540E) that connotes resistance to antifolate drugs were very prevalent, where as two additional allele (*Pfdhfr* 164L and *Pfdhps* 581G) linked to antifolates resistance showed moderate prevalence only from 2008-2011 while in Republic of Benin that involved 181 pregnant women, it was

discovered that SP maintained efficacy in the treatment and prevention of placental malaria infection in pregnant women [27, 28, 25]. Then in Luanda, Angola, the ratio of the *Pfdhfr* wild type C59 to the mutant type 59R was 60.6% to 20.6% with the presence of 18.8% mixed infections while that of *Pfdhps* 540E was 6.3% with mixed infections of 5.4% [26]. And in Tanzania, the clinical and molecular findings led to the conclusion that SP has a short useful therapeutic life in Tanzania after its introduction because of chloroquine resistance [29]

While some studies confirmed that the efficacy of SP is low due to the prevalent of resistant parasite to the drug, other studies tried to check for the efficacy of chlorproguanil-dapsone; the new antifolate drug introduced after sulphadoxine/pyrimethamine. A study in Blantyre, Malawi, which involved aged 3months to 6 years children, revealed that resistance in SP did not impair the clinical efficacy of chlorproguanil-dapsone [30]. Two other works done in Africa, one in Kenya, and the other in Tanzania confirmed this and proposed that the chlorproguanil-dapsone may act as a salvage therapy for sulphadoxine/pyrimethamine resistance [31, 32].

3.4 *Plasmodium falciparum* multi drug resistance 1 gene (*Pfmdr1*)

Pfmdr1 gene is found in chromosome 7 of the parasite and it is an antigen binding-cassette transporter that is localised in the digestive vacuole membrane [24]. It is a gene that encodes the P-glycoprotein transporter belonging to the ATP-dependent transporter family, which is responsible for the multidrug resistance in some human tumors [33]. This gene is located in the *P. falciparum* food vacuole. At least 5 mutations on *Pfmdr1* have been identified to be associated with resistance which include: N86Y, Y184F, S1034C, N1042D and D1246Y [34]. *Pfmdr1* N86Y, Y184F, and D1246Y are very common in Africa, where as S1034C and N1042D are rare in Africa but very common in Asia [28]. Mutations in *Pfmdr1* N86Y and D1246Y that are common in Africa have been linked to decreased sensitivity to drugs like amodiaquine and chloroquine while mutations in the 1034C and 1042D that are common outside Africa have been associated with increased sensitivity to mefloquine, lumefantrine and artemisinins [35].

Point mutations in the *Pfmdr1* N86Y codon has been linked to be associated with chloroquine, but the presence of the wild type N86 and increased in the copy number of *Pfmdr1* is linked to the resistance of mefloquine and quinine drugs [36, 10]. Also, the increase in gene copy number is also linked to parasite resistance to lumefantrine, artemisinin and halofantrine [37]. In a study that involved 296 children with age range 0.5 – 10years that was done in Gabon (Lambaréné), Ghana (Kumasi) and Kenya (Kisumu), the prevalence of wild type to mixed variants of the N86Y allele was 40% to 12%, 73% to 17% and 82% to 8% respectively, while the prevalence of the wild type to the mixed genotype of Y184F allele was 25% to 2%, 30% to 79%, 44% to 7% respectively, and that of the double mutant N86Y/Y184F to D1246Y was 45% to 1%, 9% to 1% and 2% to 5% respectively, with only 1% of the samples having the triple mutant N86Y/ Y184F/ D1246Y allele in both Ghana and Kenya [38]. \ Also, it was reported that patients with infected parasite that carried the N86 allele or increased *Pfmdr1* copy number have a greater risk of treatment failure to Artemether-Lumefantrine than patients that have the 86Y allele or a single copy number of *Pfmdr1* gene [39].

3.5 Cytochrome b

Atoquavone is the major inhibitor of cytochrome bc 1 (cytbc1) complex; that is the major respiratory enzyme found in the mitochondria membrane of *Plasmodium falciparum* [40]. *In vitro* resistance of *P.falciparum* to atoquavone have been linked to mutation in the *Pfcytb* region with the codons ranging from 271-274 especially the Y268S [40]. Also mutations at 268S have been linked to malarone treatment failure, although treatment failure can also be reported in the absence of this mutation [35].

3.6 *Plasmodium falciparum* multidrug resistance-associated protein 1 (*Pfmrp1*)

Pfmrp1 is an ABC transporter family of transmembrane protein that is located on chromosome 1 with a mass of 214.44KDa encoding about 1822 amino acids and it is predicted to have two nucleotide binding domains and two membrane-spanning domains with each having a 6-helical transmembrane domains [41]. It major function is to help in the transport of organic anionic substances such as: gluconate, oxidized glutathione, sulphate conjugates and also in transports of drug [41]. SNPs in *Pfmrp1* Y191H and A437S have previously been reported to be associated with *in vitro* sensitivity of

chloroquine and quinine [42]. SNP in 1466K of the *Pfmrp1* gene was associated with SP resistance in children that have uncomplicated *Plasmodium falciparum* malaria in Tanzania [43]. It was proposed that *Pfmrp1* might cause drug resistant by efflux of drugs and by managing drug-driven oxidative stress [42].

3.7 *Plasmodium falciparum* cg2 (*Pfcg2*)

Pfcg2 is a 320-330KDa gene that is found in the electron-dense patches at the periphery of the parasitophorous vacuole in the cytoplasm and in the digestive vacuole of the *Plasmodium falciparum* trophozoites and schizonts [17]. It is made up of four peptide repeats that include kappa (κ), gamma (γ), psi (ψ) and omega (ω), the kappa and the omega domain and some point mutations of cg2 have been linked to chloroquine resistance in some field isolates from Africa, thus suggesting them as molecular markers to monitor drug resistant parasite [21]. Also, a study done in Nigeria linked the cg2 Ala-281, Ddk2 repeats and *Pfmdr1* Tyr 86 genotype to chloroquine resistant parasite [44].

3.8 *Plasmodium falciparum* Ca²⁺–ATPase (*PfATP6*)

The *PfATP6* is an ATP-dependent calcium pump that is almost identical to the mammalian sarcoplasmic/endoplasmic reticulum type calcium ATPase (SERCA) pumps and acts as a receptor for artemisinin antimalarial drugs [33]. It was hypothesized that artemisinin interacts and selectively inhibits *PfATPase6* and the SNPs that have been linked to resistance of artemisinins are: S769N, A623E and E431K [45]. A report from Tanzania that involved assessing *PfATPase6* mutant S769N and A623E with 615 asymptomatic *Plasmodium falciparum* infections showed that there was no mutant genotype detected [45]. Also Krishna *et al*, 2006 reported that there have been no association between the mutant genotype S769N *PfATPase6* and antimalarial drugs like quinine and chloroquine [46].

3.9 Kelch 13 Gene (K13)

The *Plasmodium falciparum* K13 (PF3D7_1343700) is located on chromosome 13 of the parasite and it is a 1-exon gene that codes for the putative Kelch protein, which comprises of three domains: a *Plasmodium*-specific domain, a BTB/POZ and a C-terminal six-blade propeller [6]. The kelch motif containing gene known as Kelch 13 is the major molecular marker used in determining artemisinin resistance [8]. Arie *et al*, 2014 reported that mutations at Y493H, I543T, R539T and C580Y were associated with prolonged parasite survival *ex vivo* while mutations at Y493H, R539T and C580Y were associated with *in vivo* delayed parasite clearance and the M476I mutations was associated with artemisinin tolerance *in vitro* [47].

In an epidemiology study done in sub-Saharan Africa, none of the major five mutations previously described to cause resistance to artemisinins were found in the >1000 African parasite samples, although the P553L that was previously reported in Cambodia was found in the African samples [48]. This result is similar to another study that was conducted in Uganda where none of the 5 major mutations were detected in any of the samples [49]. While another reported that the 2 African non-synonymous mutations A578S and V566I SNPs were greater than 1% prevalent in at least 1 location among the 12 countries surveyed, and this is of interest because these two SNPs are found near to the C580Y that is the major genetic determinant for artemisinin resistance in Southeast Asia [50].

Table 2 Polymorphisms in molecular markers of antimalarial drug resistance and the various drugs of interest

Molecular Markers	Gene Polymorphism	Drugs (Decrease in susceptibility)
<i>Pfnhe-1</i>	Ms4760 DNNND and or NHDNHNDDDD	Quinine [14], [15] [16]
<i>Pfcrt</i>	76T	Chloroquine, Amodiaquine [22] [23]
<i>Pfdhfr</i>	51I, 59R, 108N, 164L	Pyrimethamine [5], [27], [28], [25]

<i>Pfdhps</i>	437G, 540E	Sulphadoxine [5], [27], [28], 25]
<i>Pfmdr1</i>	N86 86Y, 1246Y Increased copy number	Mefloquine, Quinine, lumefantrine [36], [10] Chloroquine, Amodiaquine [35] Mefloquine, Quinine, Lumefantrine, Halofantrine, Artemisinin [36], [10], [37]
Cytochrome b	268S	Atoquavone, Malarone [40], [35]
<i>Pfmrp1</i>	191H, 437S 1466K	Chloroquine, Quinine [42] Sulphadoxine/Pyrimethamine [43]
<i>Pfcg2</i>	Ala-281, Ddk2	Chloroquine [44]
<i>PfATP6</i>	769N, 623E, 431K	Artemisinin [45]
K13	476I, 439H, 543T, 539T, 580Y	Artemisinin [47]




Figure 1 the thread of antimalarial drug resistance in endemic areas in Africa

4 CONCLUSION

Plasmodium falciparum is the most virulent specie of *Plasmodium* and the most prevalent in sub-Saharan Africa, one of the most endemic regions of malaria in the world. Antimalarial drug resistant plays a major role in determining the therapy that will be used to eradicate this disease. For some time now, there have been records of antimalarial drug resistant to drugs like: quinine, chloroquine, mefloquine, and sulphadoxine-pyrimethamine. This is as a result of single nucleotide polymorphisms in some target genes of the parasite. From this review done, it was confirmed that most parasite isolates from this endemic regions are already resistant to most of the available drug except artemisinin. No paper reviewed documented any of the five key mutations (476I, 439H, 543T, 539T, 580Y) of the k13 gene responsible for artemisinin resistant. But almost all paper, reviewed confirmed resistance to ACTs partner drugs. Resistance to the partner drugs can lead to clinical failure for the treatment of this tropical disease. Therefore, more work has to be done to find a solution for this deadly disease before it becomes a sore in this endemic region.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

REFERENCES

1. Lee SJ, Seo E, Cho, Y. Proposal for a New Therapy for Drug-Resistant Malaria Using *Plasmodium* Synthetic Lethality Inference. *International Journal for Parasitology: Drugs and Drug Resistance* 2013;3:119–28.
2. Petersen I, Eastman R, Lanzer, M. Drug - resistant Malaria: Molecular Mechanisms and Implications for Public Health. *FEBS Letters* 2011;585(11):1551–62.
3. Le Bras J, Durand R. The Mechanisms of Resistance to Antimalarial Drugs in *Plasmodium falciparum*. *Fundamental & Clinical Pharmacology* 2003;17(2):147–53.
4. Müller IB, Hyde JE. Antimalarial Drugs: Modes of Action and Mechanisms of Parasite Resistance. *Future Microbiology* 2010;5(12):1857–73.
5. Zhong D, Afrane Y, Githeko A, Cui L, Menge DM, Yan G. Molecular Epidemiology of Drug-Resistant Malaria in Western Kenya Highlands. *BMC Infectious Diseases* 2008;8(1): 105.
6. Mishra N, Prajapati S, Kaitholia K, Bharti R, Srivastava B, Phookan S, et al. Surveillance of Artemisinin Resistance in *Plasmodium falciparum* in India Using the Kelch13 Molecular Marker. *Antimicrobial Agents and Chemotherapy* 2015;59(5):2548–53.
7. Dondorp AM, Nosten F, Yi P, Das D, Phyto AP, Tarning J, et al. Artemisinin Resistance in *Plasmodium falciparum* Malaria. *New England Journal of Medicine* 2009;361(5): 455–67.
8. Tun KM, Imwong M, Lwin KM, Win AA, Hlaing TM, Hlaing T, et al. Spread of Artemisinin-Resistant *Plasmodium falciparum* in Myanmar: A Cross-Sectional Survey of the K13 Molecular Marker. *The Lancet Infectious Diseases* 2015;15(4): 415–21.
9. Sinha S, Medhi B, Sehgal, R. Challenges of Drug-Resistant Malaria. *Parasite* 2014;21:61.
10. Apinjoh TO, Mugri RN, Miotto O, Chi HF, Tata RB, Anchang-Kimbi JK et al. Molecular Markers for Artemisinin and Partner Drug Resistance in Natural *Plasmodium falciparum* Populations Following Increased Insecticide Treated Net Coverage along the Slope of Mount Cameroon: Cross-Sectional Study. *Infectious Diseases of Poverty* 2017;6(1):136.
11. Bennett TN, Patel J, Ferdig MT, Roepe PD. *Plasmodium falciparum* Na⁺/H⁺ Exchanger Activity and Quinine Resistance. *Molecular and Biochemical Parasitology* 2007;153(1):48–58.
12. Ménard D, Andriantsoanirina V, Khim N, Ratsimbaoa A, Witkowski B, Benedet C, et al. Global Analysis of *Plasmodium falciparum* Na⁺/H⁺ Exchanger (*Pf*nhe-1) Allele Polymorphism and Its Usefulness as a Marker of *in vitro* Resistance to Quinine. *International Journal for Parasitology: Drugs and Drug Resistance* 2013;3: 8–19.
13. Andriantsoanirina V, Ménard D, Rabearimanana S, Hubert V, Bouchier C, Tichit M, et al. Association of Microsatellite Variations of *Plasmodium falciparum* Na⁺/H⁺ Exchanger (*Pf*nhe-1) Gene with Reduced *In vitro* Susceptibility to Quinine: Lack of Confirmation in Clinical Isolates from Africa. *The American Journal of Tropical Medicine and Hygiene* 2010;82(5): 782–87.
14. Briolant S, Pelleau S, Bogreau H, Hovette P, Zettor A, Castello J et al. *In vitro* Susceptibility to Quinine and Microsatellite Variations of the *Plasmodium falciparum* Na⁺/H⁺ Exchanger (*Pf*nhe-1) Gene: The Absence of Association in Clinical Isolates from the Republic of Congo.” *Malaria Journal* 2011;10(1):37.
15. Henry M, Briolant S, Zettor A, Pelleau S, Baragatti M, Baret E, et al. *Plasmodium falciparum* Na⁺/H⁺ Exchanger 1 Transporter is Involved in Reduced Susceptibility to Quinine. *Antimicrobial Agents and Chemotherapy* 2009;53:1926–30.
16. Sinou V, Quang LH, Pelleau S, Huong VN, Huong NT, Tai LM, et al. Polymorphism of *Plasmodium falciparum* Na⁺/H⁺ Exchanger Is Indicative of a Low *in vitro* Quinine Susceptibility in Isolates from Viet Nam. *Malaria Journal* 2011;10(1):164.
17. Cooper RA, Papakrivos J, Lane KD, Fujioka H, Lingelbach K, Wellem TE. *Pf*CG2, a *Plasmodium falciparum* Protein Peripherally Associated with the Parasitophorous Vacuolar

- Membrane, Is Expressed in the Period of Maximum Hemoglobin Uptake and Digestion by Trophozoites. *Molecular and Biochemical Parasitology* 2005;144(2):167–76.
18. Wellem's TE, Plowe CV. Chloroquine-Resistant Malaria. *The Journal of Infectious Diseases* 2001;184(6): 770–76.
19. Ndam NK, Basco LK, Ngane VF, Ayoub'a A, Ngolle EM, Deloron P, et al. Reemergence of Chloroquine-Sensitive Pfcrt K76 *Plasmodium falciparum* Genotype in Southeastern Cameroon. *Malaria Journal* 2017;16(1):130.
20. Alifrangis M, Lusingu JP, Mmbando B, Dalgaard MB, Vestergaard LS, Ishengoma D, et al. Five-Year Surveillance of Molecular Markers of *Plasmodium falciparum* Antimalarial Drug Resistance in Korogwe District, Tanzania: Accumulation of the 581G Mutation in the *P. falciparum* Dihydropteroate Synthase Gene. *The American Journal of Tropical Medicine and Hygiene* 2009;80(4): 523–27.
21. Sharma YD. Genetic Alteration in Drug Resistance Markers of *Plasmodium falciparum*. *Indian Journal of Medical Research* 2005;121(1):13.
22. Djimdé AA, Doumbo OK, Traore O, Guindo AB, Kayentao K, Diourte Y, et al. Clearance Of Drug-Resistant Parasites as a Model For Protective Immunity in *Plasmodium falciparum* Malaria. *The American Journal of Tropical Medicine and Hygiene* 2003;69(5):558–63.
23. Happi CT, Gbotosho GO, Folarin OA, Bolaji OM, Sowunmi A, Kyle DE, et al. Association between Mutations in *Plasmodium falciparum* Chloroquine Resistance Transporter and *P. falciparum* Multidrug Resistance 1 Genes And *in vivo* Amodiaquine Resistance In *P. falciparum* Malaria-Infected Children In Nigeria. *The American Journal of Tropical Medicine and Hygiene* 2006;75(1):155–61.
24. Meyer CG, May J, Arez AP, Gil JP, Do RosarioV. Review: Genetic Diversity of *Plasmodium falciparum*: Asexual Stages. *Tropical Medicine & International Health* 2002;7(5): 395–408.
25. Bertin G, Briand V, Bonaventure D, Carrieu A, Massoug'bodji A, Cot M, Deloron P. Molecular Markers of Resistance to Sulphadoxine-Pyrimethamine during Intermittent Preventive Treatment of Pregnant Women in Benin. *Malaria Journal* 2011;10(1):196.
26. Figueiredo P, Benchimol C, Lopes D, Bernardino L, do Rosário VE, Varandas L, et al. Prevalence of *Pfmdr1*, *Pfcrt*, *Pfdhfr* and *Pfdhps* Mutations Associated with Drug Resistance, in Luanda, Angola. *Malaria Journal* 2008;7(1):236.
27. Zeile I, Gahutu J-B, Shyirambere C, Steininger C, Musemakweri A, Sebahungu F, et al. Molecular Markers of *Plasmodium falciparum* Drug Resistance in Southern Highland Rwanda. *Acta Tropica* 2012;121(1):50–54.
28. Mbogo GW, Nankoberanyi S, Tukwasibwe S, Baliraine FN, Nsobya SL, Conrad MD, et al. Temporal Changes in Prevalence of Molecular Markers Mediating Antimalarial Drug Resistance in a High Malaria Transmission Setting in Uganda. *The American Journal of Tropical Medicine and Hygiene* 2014;91(1):54–61.
29. Mugittu K, Ndejemb'i M, Malisa A, Lemnge M, Premji Z, Mwit'a A, et al. Therapeutic Efficacy of Sulfadoxine-Pyrimethamine and Prevalence of Resistance Markers in Tanzania Prior to Revision of Malaria Treatment Policy: *Plasmodium falciparum* Dihydrofolate Reductase and Dihydropteroate Synthase Mutations In Monitoring *in vivo* Resistance. *The American Journal of Tropical Medicine and Hygiene* 2004;71(6):696–702.
30. Kublin JG, Dzinjalama'la FK, Kamwendo DD, Malkin EM, Cortese JF, Martino LM, et al. Molecular Markers for Failure of Sulfadoxine-Pyrimethamine and Chlorproguanil-Dapsone Treatment of *Plasmodium falciparum* Malaria. *The Journal of Infectious Diseases* 2002;185(3):380–88.
31. Amukoye E, Winstanley PA, Watkins WM, Snow RW, Hatcher J, Mosobo M, et al. Chlorproguanil-Dapsone: Effective Treatment for Uncomplicated *falciparum* Malaria. *Antimicrobial Agents and Chemotherapy* 1997;41(10):2261–64.
32. Mutabingwa T, Nzila A, Mberu E, Nduati E, Winstanley P, Hills E, et al. Chlorproguanil-Dapsone for Treatment of Drug-Resistant *falciparum* Malaria in Tanzania. *The Lancet* 2001;358(9289):1218–23.
33. Price RN, Uhlemann A-C, Brockman A, McGready R, Ashley E, Phaipun L et al. Mefloquine Resistance in *Plasmodium falciparum* and Increased *Pfmdr1* Gene Copy Number. *The Lancet* 2004;364(9432):438–47
34. Kaewpruk N, Tan-ariya P, Ward SA, Sitthichot N, Suwandittakul N, and Mungthin M. PFMDR1 Polymorphisms Influences on *in Vitro* Sensitivity of Thai *Plasmodium falciparum* Isolates to Primaquine, Sitamaquine and Tafenoquine. *Southeast Asian J. Trop. Med. Public Health* 2016;47:366–76.

35. Cui L, Mharakurwa S, Ndiaye D, Rathod PK, Rosenthal PJ. Antimalarial Drug Resistance: Literature Review and Activities and Findings of the ICEMR Network. *The American Journal of Tropical Medicine and Hygiene* 2015;93(3):57–68.
36. Spalding MD, Eyase FL, Akala HM, Bedno SA, Prigge ST, Coldren RL, et al. Increased Prevalence of the Pfdhfr/Phdhps Quintuple Mutant and Rapid Emergence of Pfdhps Resistance Mutations at Codons 581 and 613 in Kisumu, Kenya. *Malaria Journal* 2010;9(1):338.
37. Picot S, Olliaro P, de Monbrison F, Bienvenu A-L, Price RN, Ringwald P. “A Systematic Review and Meta-Analysis of Evidence for Correlation between Molecular Markers of Parasite Resistance and Treatment Outcome in *falciparum* Malaria. *Malaria Journal* 2009;8(1):89.
38. Nguetse CN, Adegnikaa AA, Agbenyega T, Ogutu BR, Krishna S, Kremsner PG, et al. Molecular Markers of Anti-Malarial Drug Resistance in Central, West and East African Children with Severe Malaria. *Malaria Journal* 2017;16(1):217.
39. Venkatesan M, Gadalla NB, Stepniowska K, Dahal P, Nsanzabana C, Moriera C, et al. Polymorphisms in *Plasmodium falciparum* Chloroquine Resistance Transporter and Multidrug Resistance 1 Genes: Parasite Risk Factors That Affect Treatment Outcomes for *P. falciparum* Malaria After Artemether-Lumefantrine and Artesunate-Amodiaquine. *The American Journal of Tropical Medicine and Hygiene* 2014;91(4):833–43.
40. Ekala MK, Khim N, Legrand E, Randrianarivelojosia M, Jambou R, Fandeur T, et al. Sequence Analysis of *Plasmodium falciparum* Cytochrome b in Multiple Geographic Sites. *Malaria Journal* 2007;6(1):164.
41. Antony HA, Parija SC. Antimalarial Drug Resistance: An Overview. *Tropical Parasitology* 2016;6(1):30–41.
42. Dahlström S, Veiga MI, Mårtensson A, Björkman A, Gil JP. Polymorphism in *PfMRP1* (*Plasmodium falciparum* Multidrug Resistance Protein 1) Amino Acid 1466 Associated with Resistance to Sulfadoxine-Pyrimethamine Treatment. *Antimicrobial Agents and Chemotherapy* 2009;53(6): 2553–2556.
43. Dahlström S, Ferreira PE, Veiga MI, Sedighi N, Wiklund L, Mårtensson A et al. *Plasmodium falciparum* Multidrug Resistance Protein 1 and Artemisinin-Based Combination Therapy in Africa. *The Journal of Infectious Diseases* 2009;200(9):1456–64.
44. Adagu IS, Warhurst DC. Association of Cg2 and Pfmdr1 Genotype with Chloroquine Resistance in Field Samples of *Plasmodium falciparum* from Nigeria. *Parasitology* 1999;119(4):343–48.
45. Mugittu K, Genton B, Mshinda H, Beck HP. Molecular Monitoring of *Plasmodium falciparum* Resistance to Artemisinin in Tanzania. *Malaria Journal* 2006;5(1):126.
46. Krishna S, Woodrow CJ, Staines HM, Haynes RK, Mercereau-Puijalon, O. Re-Evaluation of How Artemisinins Work in Light of Emerging Evidence of *in vitro* Resistance. *Trends in Molecular Medicine* 2006;12(5):200–205.
47. Arie F, Witkowski B, Amaratunga C, Beghain J, Langlois A-C, Khim N, et al. A Molecular Marker of Artemisinin-Resistant *Plasmodium falciparum* Malaria. *Nature* 2013;505: 50.
48. Taylor SM, Parobek CM, DeConti DK, Kayentao K, Coulibaly SO, Greenwood BM, et al. Absence of Putative Artemisinin Resistance Mutations Among *Plasmodium falciparum* in Sub-Saharan Africa: A Molecular Epidemiologic Study. *The Journal of Infectious Diseases* 2015;211(5):680–88.
49. Conrad MD, Bigira V, Kapisi J, Muhindo M, Kamya MR, Havlir DV et al. Polymorphisms in K13 and Falcipain-2 Associated with Artemisinin Resistance Are Not Prevalent in *Plasmodium falciparum* Isolated from Ugandan Children. *PLOS ONE* 2014;9(8):e105690.
50. Kamau E, Campino S, Amenga-Etego L, Drury E, Ishengoma D, Johnson K et al. K13-Propeller Polymorphisms in *Plasmodium falciparum* Parasites From Sub-Saharan Africa. *The Journal of Infectious Diseases* 2015;211(8):1352–55.