

1 Malaria related deaths among children with manifestations of fever
2 symptoms on admission in a secondary health care institution in
3 Western Region of Ghana - a retrospective study

4 Short title: Retrospective study of malaria deaths among children in Ghana.
5

6 ABSTRACT

7 Background: Malaria is a major contributor to deaths in children especially in sub-Saharan
8 Africa. Children less than five years of age are susceptible to malaria infection in endemic
9 regions leading to serious complications. Malaria causes death in children either directly
10 through Cerebral Malaria (CM) and Severe Malaria Anaemia (SMA) or indirectly through
11 co-morbidity with pneumonia or a sequela like hypoglycaemia.

12 Methods: This retrospective study examined malaria-related deaths among children at Effia
13 Nkwanta hospital within a study period of 3 years.

14
15 Results: A total of 1,416 medical records were reviewed, out of which 223 were medical
16 records of children with fatal outcomes. Deaths over the study period due to all causes were
17 15.7% (223/1416) and confirmed malaria was 13.7% (40/292). Deaths due to all causes and
18 to confirmed malaria decreased from 21.6% and 24.3% in 2010 to 11.1% and 4.4% in 2012,
19 respectively. Anti-malarial testing was done for 152 of the children with 40 positive and 112
20 negative results. Seventy-one children had no malaria testing done on them, with 23.4% in
21 2010 40.3% in 2011 and 35.5% in 2012. Anti-malarial treatment was administered to 83% of
22 children who tested negative and 80% of children without anti-malarial testing.

23 Conclusion: Deaths in the children declined from 2010 to 2012 in this study. Despite this
24 improvement, there was poor anti-malarial testing and improper use of anti-malarial

25 treatment. National malaria programs should ensure improvement in anti-malarial testing and
26 strict adherence to anti-malarial treatment protocol.

27 Key words: Malaria, Children, Mortality, Anemia, Death.

28 Introduction

29 Malaria remains a serious public health issue in malaria-endemic areas, with
30 residents at the risk of infection and higher risks are often seen in children and pregnant
31 women¹. Children less than five years are susceptible to malaria infection leading to serious
32 complications and death if not promptly and adequately treated with effective anti-malarial
33 drugs¹. *Plasmodium falciparum* is the predominant species in sub-Saharan Africa, the major
34 cause of severe disease leading to death¹. The exposure of children to sub-lethal *P.*
35 *falciparum* infection leads to the gradual acquisition of clinical immunity². However, some
36 children unfortunately never live to develop this immunity as their infection progresses to
37 severe disease leading to death before adulthood³.

38 Malaria infection due to *P. falciparum* can cause mortality directly by being the
39 underlying cause of death. Cerebral malaria and severe malaria anemia are very common
40 direct causes of childhood mortality⁴. Cerebral malaria defined as confirmed *P. falciparum*
41 infection with unrousable coma⁵ causes death in children due to hypo-perfusion of brain cells
42 as a result of sequestration of infected red cells within cerebral capillaries⁶. Another common
43 direct cause of death in *P. falciparum* infected children is anemia, referred to as severe
44 malaria anemia (SMA)⁴. World Health Organization (WHO) defines SMA as a hemoglobin
45 level (Hb) below 5.0 g/dL in the presence of malaria parasites⁷. Indirect causes of death in
46 malaria are cases in which malaria is only a contributing factor, such as in a child with
47 malaria-related anemia developing pneumonia or hypoglycemia and dying from it⁸.

48 There is a reported 18% global decline in malaria cases from 262 million in the year
49 2000 to about 214 million in 2015.⁹ It was estimated that 88% of the new cases in 2015
50 occurred in sub-Saharan Africa⁹. However, the region has recorded a tremendous progress in
51 terms of reduction in the number of malaria deaths among children under 5 years of age, with
52 the bulk of global decrease occurring in the region.⁹ The estimated number of malaria deaths
53 among children under 5 years of age fell from 694,000 in 2000 to 292,000 in 2015.⁹ These
54 findings suggest that malaria is no longer the leading but the fourth leading cause of death
55 among children, accounting for about 10% in 2015.⁹ This declining mortality in children from
56 sub-Saharan Africa is due to the positive outcome from **scale-up** of malaria intervention
57 programs deployed in the region.¹⁰ These control programs have facilitated the reduction of
58 childhood malaria morbidity and mortality in most parts of sub-Saharan Africa^{11, 12} These
59 control **programs** include **household** and community activities such as the use of insecticide
60 treated nets ITNs, indoor residual spraying IRS, larval control and health facilities activities
61 including adequate case management for malaria.¹³

62 Adequate case management of malaria is a veritable arm of the malaria control
63 program. It is primarily carried out in health facilities. Invariably, the quality of care in health
64 facilities can impact greatly on **malaria-related mortality** especially among children. The
65 **generally** dilapidated state of health care in many Africa countries negatively impacted on the
66 quality of management and treatment of many diseases especially malaria in children.¹⁴
67 However, there **has** been some improvement in malaria case management over the years with
68 the improvement of diagnosis and treatment by the provision of tools for on the spot
69 diagnosis like **the** rapid diagnostic test for malaria, effective drugs, and skilled health care
70 personnel.^{14,15} Despite this, quality of care in hospitals and clinics still need improvement
71 since avoidable malaria deaths have been reported.¹⁶

72 In Ghana, childhood mortality is a serious public health issue as diseases like
73 pneumonia; diarrhea, malnutrition, and malaria take their toll on children¹⁷. Malaria remains
74 a constant threat to the life of children in Ghana especially those who live in rural areas¹⁸. In
75 the first quarter of 2016, the country recorded a total of 2.2 million cases of malaria in all
76 health facilities nationwide. The number of malaria confirmed deaths recorded was 379
77 which was a 15.2% drop from the value in the first quarter of 2015. Of these, there were 152
78 malaria deaths in children under 5 years of age compared to 182 recorded in 2015 ¹⁹
79 However; this noted decline is not uniform within the country. It is very important to
80 constantly review mortality figures and pattern in different parts of the country to assess the
81 progress of malaria control programs and the dynamics of malaria infection. Hence, this
82 retrospective study evaluated the trend in malaria-related deaths among children who died on
83 admission for febrile illness in a referral hospital in Sekondi-Takoradi metropolis Ghana over
84 a three-year period.

85 Materials and Methods

86 Study Area

87 Effia-Nkwanta Regional Hospital is a secondary health-care institution and is the only
88 referral hospital for the whole western region of Ghana. Ghana is bordered by Togo on the
89 east and Ivory Coast on the west (figure 1). Effia Nkwanta hospital is located at the Regional
90 capital, Sekondi-Takoradi. It serves all other hospitals within the entire 22 districts of the
91 western region and sub-divisions of 13 major districts. Details of the study area have been
92 published previously ²⁰.




Figure 1: Map of Ghana showing location of Sekondi-Takoradi²¹

Data Collection

This is a retrospective study that was carried out at the Paediatric Unit of Effia-Nkwanta Regional Hospital (ENRH). Before this study, Paediatric emergencies were seen and where necessary, were admitted to the children's ward of the hospital. The patients admitted were recorded in the admissions register. The admissions register of the Accident and Emergency Unit of the hospital was used to collect **folder numbers** of children who were recruited into the retrospective study. Children admitted with fever as one of the presenting symptoms with subsequent fatal outcome over the study period of 2010, 2011 and 2012 were included in the study. Their medical records were searched and retrieved. Information on the demographics (i.e. age and sex), clinical and laboratory findings were collected. Clinical information retrieved included other symptoms accompanying fever, **the** weight of the child and temperature as well as information on sickle cell trait and blood transfusion during admission. Reported laboratory findings included malaria parasite diagnosis, white blood cell counts, sickling status and haemoglobin concentration. Information on treatment histories **was** also collected as well as the number of days the patients spent in **the** hospital. **Permission was granted for this work from the regional office Ghana Health Service and the medical director of the facility.**

Patients with malaria admitted in the hospital were classified according to the WHO criteria for mild and severe malaria⁷. Diagnosis of malaria parasite was done using *P. falciparum* specific rapid diagnostic test kits, containing membrane strip, precoated with a monoclonal antibody which is specific to the histidine-rich protein 2 of the *Plasmodium falciparum* (First Response, Premier Medical Corporation Ltd) and microscopy using Giemsa staining. The rapid response kit detects *P. falciparum* antigens and the presence of two lines in the test kit well indicate *P. falciparum* positive. Malaria parasites were confirmed with thick and thin peripheral blood smears stained with Giemsa and examined microscopically using 100 power fields under oil immersion. Two microscopists read the slides independently and a third microscopist was called upon to resolve discordant results. Haemoglobin estimation and white cell count were done as part of the complete blood count from an automated blood cells analyzer machine (Sysmex Haematology Analyser, Xuzhou Hengda, China). Certification and establishment of the cause of death were done mainly by using clinical judgment based on history, physical examination and laboratory findings. These were done by experienced medical officers or specialist paediatricians after proper review of patient's records.

Malaria was defined as the presence of any asexual blood stages of *P. falciparum* species in thin smear blood film while malaria negative samples were confirmed when 100 high power fields have been examined using x100 oil immersion objective lens. Anaemia was defined based on WHO criteria of haemoglobin (Hb) levels <11 g/dL^{22, 23}, and categorized as mild anaemia (Hb <11 g/dL), moderate anaemia (Hb <10 g/dL) and severe anaemia (Hb <7 g/dL). A febrile malarial episode in children was defined as a reported history of fever within the last 24–72 hrs of admission or a measured axillary temperature of 37.5°C or greater (or both), with a positive peripheral blood smear slide with asexual forms of *P. falciparum* at any level of parasite density at time of contact with ENRH clinic²⁴. A malaria death is defined as one death per parasitologically diagnosed clinical episode¹¹. We calculated malaria mortality

(expressed as a percentage) with the formula: malaria deaths per year / Number of children in the population at midyear. We calculated case fatality rate (express as a percentage) with the formula: malaria deaths per year / Number of confirmed malaria cases among admitted children per year²⁵.

Statistical analysis

Univariable analyses were performed and results were presented as proportions and mean $\pm$ standard deviation (SD). Statistical significance for cross-tabulations was performed using Pearson χ^2 test for categorical variables. One-way analysis of variance (ANOVA) was used for the comparison of the mean of continuous variables. All tests were two-tailed and $P \leq 0.05$ was considered statistically significant. All statistical analyses were performed using IBM SPSS software version 21.0 (IBM Corporation, Armonk, NY, USA).

Results

Characteristics of the study population

Table 1 describes the characteristics of the 223 children who died during the three years study period. Severe anaemia ($Hb < 7$ g/dL) was seen in 34.1% of the children while 18.8% of the children that died were not anaemic ($Hb > 11$ g/dL). Malaria diagnostic test was not performed in about 31.9% of the children that died. However, 17.9% of the children were laboratory confirmed malaria positives while 50.2% were malaria negatives. Ninety-seven children (43.5%) died within the first 24 hours of admission, 37.2% died within 72 hours and 19.3% after 72 hours of admission.

Figure 2 describes the percentage mortality of the children admitted into the children's ward. During the study period, a total of 1,416 children were admitted to the children unit of Effia-Nkwanta Regional Hospital. Out of this total number of admitted children, 1193 children were successfully treated and discharged home, while 223 children,

unfortunately, died over the same period, with a percentage mortality of 15.7%. There was a progressive decline in the percentage of mortality from 21.6% in 2010 to 11.1% in 2012 (Figure 2). Figure 3 describes malaria case fatality in the study population. During the three years study period, 40 children died out of the 292 admitted with supposedly malaria cases with a total case fatality rate of 13.7%. There was also a steady decline in the case fatality rate for laboratory-confirmed malaria cases from 24.3% in 2010 to 4.4% in 2012 (Figure 3).

Table 2 shows the characteristics of the study population stratified against malaria laboratory investigations. A significant proportion of malaria diagnosis was done without malaria testing and some were negative on malaria test. Antimalarial treatments were accurately given to all of the children with confirmed malaria test, while 83% and 80% of the children that tested negative and those without malaria test respectively were incorrectly prescribed anti-malarial drugs. Children without malaria testing significantly decreased as they stayed longer in the hospital (>24hrs, 63.4%; <72hrs, 28.2%; >72hrs, 8.4%).

Figure 4 shows that malaria testing rate worsened over time with 23.4% of children not tested in 2010, 40.3% in 2011 and 35.1% not tested in 2012.

Figure 5 summarizes the combination of the prescribed medications during the study period stratified by year. Apparently, patients without laboratory-confirmed malaria diagnosis or patients with negative malaria diagnosis were given a parenteral anti-malarial treatment of either quinine or artemether.

Other Non-Malaria Causes of Death

Figure 6 shows the distribution of other non-malaria causes of death. Among these children that died from non-malaria causes, 12.1% were diagnosed with sepsis, 9.4% were

185 diagnosed with bronchopneumonia, 8.5% were diagnosed with gastroenteritis, and 7.2% were
186 diagnosed with severe anaemia. Children who died without a specific diagnosis were 19.7%.

187 Discussion

188 This study evaluated the contribution of malaria to the mortality of children in the
189 children ward of Effia-Nkwanta Regional Hospital over a 3 year study period. There was a
190 steep decline in malaria deaths from 2010 to 2012. This is a drastic drop and an interesting
191 finding indeed. Firstly, there was high case fatality rate for malaria in 2010 compared to the
192 subsequent years, which might be due to both in the hospital and out of hospital factors.
193 These factors have contributed to poor and fatal treatment outcomes in children with
194 malaria²⁶. Late arrival to the hospital is one major out of hospital factor responsible for
195 malaria mortality in children in many countries in sub-Saharan Africa especially Ghana²⁷.
196 This is due to, poor health seeking behavior caused by poor understanding of the symptoms
197 of malaria, over reliance on home and herbal remedies and financial constraints^{27, 28}. In
198 hospital factors, especially in Ghana are hinged majorly on the poor quality of care for the
199 sick child, including poor and delayed diagnosis, lack of appropriate medications and
200 treatment and lack of well-trained health care personnel^{29,30}. Poor quality of care has been
201 reported in regional hospitals similar to the facility where the present study took place^{30,31}.

202 In 2011 there was a tremendous reduction in malaria confirmed deaths despite an increase in
203 admitted confirmed malaria cases. An improvement in the arrival time of the sick children to
204 the hospital and improvement in the quality of care in the hospital might have been
205 responsible for this. Parents and caregivers would have become better educated on, early
206 recognition of symptoms and signs of malaria, the dangers of herbal medication and the
207 importance of coming early to the hospital with their sick children. This is the likely
208 outcome of the various forms of health education carried out by the National Malaria Control

Program¹⁹. More so the improved enrolment of patients to National Health Insurance scheme would have overcome the barrier of financial constraint allowing more patients to assess health care in the hospital.^{29, 32} Also, increase in the confirmed malaria cases on admission, might be due to an improvement in malaria diagnosing capacity of the regional hospital based on the massive deployment of rapid diagnostic test kits as well as training of health care personnel on the use of the test kits^{19, 33}. This improvement in case identification and the diagnosis was invariably followed by prompt, adequate and successful treatment outcomes for the majority of the children that was admitted in 2011. This success in treatment was perhaps due to better training of the healthcare personnel as well as the availability of appropriate therapeutic drugs and effective therapy with strict adherence to standard treatment guidelines¹⁹. Ghana adopted the WHO artemisinin-based combination treatment (ACT) policy in 2004 with artesunate-amodiaquine as first-line treatment and a subsequent review in 2009 involving the inclusion of artemether-lumefantrine and dihydroartemisinin-piperaquine as alternative first-line drugs for children who cannot tolerate artesunate-amodiaquine combination therapy. Parenteral treatment includes artemether and quinine injection for severe malaria. This study showed that these anti-malarial drugs were basically used in the management of malaria in the children in this hospital, but the details of how they were used and administered were not available. Studies, however, showed that there were challenges in the practicality of this policy as it took some time for health workers through frequent training to fully acquire the necessary skills to use these treatments effectively^{31, 34, 35}.

In 2012, there was further improvement in the case fatality rate coupled with a decline in admitted malaria confirmed cases despite an overall increase in admitted children. The decline in the admitted malaria confirmed cases might be a reflection of the overall decline in the global and regional malaria burden noted in the country^{9, 19}, or a dividend of activities of

234 National Malaria Control Program involving continuous campaign on the use of ITNs and
235 IRS¹⁹. Another possible reason for the decline might be that most parents or caregivers have
236 well improved knowledge in the home management of uncomplicated malaria with an
237 affordable ACT, thereby preventing complication that will warrant admission in hospital^{36, 37}.
238 More so many primary health centers and clinics in the area are now well equipped with
239 RDTs and ACTs to handle malaria in children^{37, 38}. The lowest case fatality for malaria
240 recorded in 2012, was perhaps the result of progressive improvement in case management
241 and strict observance of the standard treatment guideline for malaria in the hospital³⁸.

242 It is important to highlight the ever-nagging problem of poor laboratory diagnosis of malaria
243 observed in this study; as 31.9% of children who died never had malaria test done on them.
244 This is a very common problem seen in many hospitals and clinics in developing countries
245 and it has been blamed on the incessant break down of the diagnostic machine and frequent
246 stock out of reagents^{39 40}. In all, the situation worsened between 2010 (23%) to 2012 (35%).
247 Another study was done among discharged children over the same period in the same hospital
248 also reported a significant proportion of children who were not tested for malaria^{20, 41}.
249 However, in this study, time spent on admission before death may have played a role in the
250 number of children who died without malaria test done on them, since the majority of
251 children who died within the first 24 hours never had malaria test done on them. The chances
252 of getting a malaria test done in the laboratory increased if the child survived beyond 72
253 hours before death. The presumptive diagnosis of malaria was woefully inaccurate in this
254 study as significant proportions of the case diagnosed were not confirmed by laboratory
255 investigation. This is really an important finding for clinicians to be reminded that in all
256 cases, their clinical judgment should always be supported by laboratory investigations.

257 As seen in most studies, children under five years of age remain the most vulnerable group
258 for fatal consequences of diseases and infections^{42, 43}. The higher proportion of under-five

children among those who died with malaria in this study further buttresses the age-specific severity of malaria morbidity and mortality as seen in malaria-endemic areas^{19,42,43}. Taking a closer look at the specific diagnosis of children who tested positive for malaria in this study, CM and SMA were more common. CM and SMA are the most common complications of malaria linked with pediatric mortality^{4, 44}.

Aside from malaria mortality, there was a progressive decline in overall mortality, despite the increased patient load over the study period. This decline is consistent with the declining trend in child mortality reported for sub-Saharan Africa over the years^{42, 43}. This can be attributed to the continuous improvement in the healthcare system and implementation of various preventative measures such as child survival programs and vaccinations as well as improvement in diagnostic methods and effective treatments that mitigate the impact of diseases responsible for child mortality in the region⁴⁵. Other diseases were also responsible for child mortality in this study as more of these children that had malaria laboratory investigation done were negative. Bacterial and viral diseases, malnutrition, congenital abnormalities, intrapartum events, and injuries have all been implicated in child mortality^{42, 43}⁴⁵. This study noted a rather high number of children with missing/unknown diagnoses. The probable clinical explanation for this would have been that some children die within a few hours of arrival to the ward or arrive at a period where the doctors and laboratory personnel are not readily available on duty especially during the weekends. Weekend admissions would have probably contributed to the mortality due to lack of quality care, which is a common finding in hospitals in similar settings⁴⁶.

Limitations

Despite the fact that in this study the anti-malarial oral ACT and parenteral drugs used were all documented, a detailed audit of how these treatments were used vis-à-vis dosages,

timelines, duration, ancillary treatment like infusions and correction of hypoglycemia, were not documented. Availability of this document **would have** revealed more information on the outcome of the treatment. There is a possibility that some discharged children died at home thereby making the mortality rate in this study a little conservative. Finally, it is possible that the children who had no malaria test done on them would have skewed the result in the comparison over time in this study.

Conclusion

This study noted a steep decline in **all-cause** mortality and malaria confirmed deaths despite an increase in admitted patients over a 3 years period in the pediatric department of Effia Nkwanta hospital. There was however poor anti-malarial testing among the children, a practice that **worsens** over the study period. There were incorrect diagnoses and treatments of malaria noted in this study with a significant proportion of children wrongly diagnosed with those negative for malaria and those not tested receiving anti-malarial treatment. Despite the commendable decline of deaths in children noted in this study, there is room for improvement vis-à-vis improvement in anti-malarial testing, and strict adherence to proper anti-malarial treatment protocol to ensure adequate case management which will go a long way in reducing malaria mortality in children.

References

1. Artavanis-Tsakonas K, Tongren JE, Riley EM. (2003). The war between the malaria parasite and the immune system: immunity, immunoregulation and immunopathology. Clin Exp Immunol, 133(2):145-52.
2. Bull PC, Marsh K. (2002). The role of antibodies to Plasmodium falciparum-infected-erythrocyte surface antigens in naturally acquired immunity to malaria. Trends Microbiol, 10(2):55-8.

3. Greenwood B, Marsh K, Snow R. (1991). Why do some African children develop severe malaria? *Parasitol Today*, 7(10):277-81.
4. Menendez C, Fleming AF, Alonso PL. (2000). Malaria-related anaemia. *Parasitol Today*, 16(11):469-76.
5. Molyneux ME, Taylor TE, Wirima JJ, Borgstein A. (1989). Clinical features and prognostic indicators in paediatric cerebral malaria: a study of 131 comatose Malawian children. *Q J Med*, 71(265):441-59.
6. Idro R, Jenkins NE, Newton CR. (2005). Pathogenesis, clinical features, and neurological outcome of cerebral malaria. *Lancet neurology*, 4(12):827-40.
7. WHO. Severe falciparum malaria. (2000). World Health Organization, Communicable Diseases Cluster. *Trans R Soc Trop Med Hyg*, 94 Suppl 1:S1-90.
8. Rowe, A.K., Rowe, S.Y., Snow, R.W., Korenromp, E.L., Schellenberg, J.R.A., Stein, C., Nahlen, B.L., Bryce, J., Black, R.E. and Steketee, R.W.,(2006). The burden of malaria mortality among African children in the year 2000. *Int J Epidemiol*, 35(3):691-704.
9. World Health Organization. (2016). World malaria report 2015. World Health Organization.
10. World Health Organization. (2015). World malaria report 2014. Available: http://www.who.int/malaria/publications/world_malaria_report_2014/en/.
11. Thwing J, Eisele TP, Steketee RW. (2011). Protective efficacy of malaria case management and intermittent preventive treatment for preventing malaria mortality in children: a systematic review for the Lives Saved Tool. *BMC Public Health*, 11:1-9.
12. O'Meara WP, Mangeni JN, Steketee R, Greenwood B. (2010) Changes in the burden of malaria in sub-Saharan Africa. *The Lancet infectious diseases*, 10(8):545-55.

332 13. Nyarango, P. M., Gebremeskel, T., Mebrahtu, G., Mufunda, J., Abdulummini, U.,
333 Ogbamariam, A., Kosia, A., Gebremichael, A., Gunawardena, D., Ghebrat, Y.,
334 Okbaldet, Y. (2006). A steep decline of malaria morbidity and mortality trends in
335 Eritrea between 2000 and 2004: the effect of combination of control methods. *Malaria*
336 *journal*, 5(1), 33.

337 14. Bryan, L., Conway, M., Keesmaat, T., McKenna, S., & Richardson, B. (2010).
338 Strengthening sub-Saharan Africa's health systems: a practical approach. *McKinsey*
339 *Quarterly*.

340 15. Obonyo, C. O., Vulule, J., Akhwale, W. S., & Grobbee, D. E. (2007). In-hospital
341 morbidity and mortality due to severe malarial anemia in western Kenya. *Am J Trop*
342 *Med Hyg*, 77: 23-28.

343 16. Biai, S., Rodrigues, A., Gomes, M., Ribeiro, I., Sodemann, M., Alves, F., & Aaby, P.
344 (2007). Reduced in-hospital mortality after improved management of children under 5
345 years admitted to hospital with malaria: randomised trial. *Bmj*, 335(7625), 862.

346 17. Caulfield LE, de Onis M, Blossner M, Black RE. Undernutrition as an underlying
347 cause of child deaths associated with diarrhea, pneumonia, malaria, and measles. *Am*
348 *J Clin Nutr*. 2004;80(1):193-8.

349 18. Binka FN, Indome F, Smith T. Impact of spatial distribution of permethrin-
350 impregnated bed nets on child mortality in rural northern Ghana. *Am J Trop Med*
351 *Hyg*. 1998;59(1):80-5.

352 19. Ghana Health Service. First Quarter Bulletin 2016. National Malaria Control
353 Surveillance, 2016
354 https://www.ghanahealthservice.org/downloads/1st%20Quarter%20Malaria%20Bulletin%20-%202016_14th%20July%202016.pdf. Site accessed Jan, 2018.

20. Orish, V. N., Ansong, J. Y., Anagi, I. B., Onyeabor, O. S., Sanyaolu, A. O., & Iriemenam, N. C. (2015). Malaria and associated co-morbidity in children admitted with fever manifestation in Western Ghana: A retrospective study. *The Journal of Infection in Developing Countries*, 9(11), 1257-1263.
21. Arias D, Taylor L, Ofori-Atta A, Bradley EH. Prayer camps and biomedical care in Ghana: Is collaboration in mental health care possible?. *PloS one*. 2016 Sep 12;11(9):e0162305.
22. WHO. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. <http://www.who.int/vmnis/indicators/haemoglobinpdf>. 2011.
23. DeMaeyer EM, Dallman P, Gurney JM, Hallberg L, Sood SK, Srikantia SG. Preventing and controlling iron deficiency anaemia through primary health care. http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/9241542497.pdf. [Published 1989. Updated 1990].
24. Massaga JJ, Kitua AY, Lemnge MM, Akida JA, Malle LN, Rønn AM, Theander TG, Bygbjerg IC. Effect of intermittent treatment with amodiaquine on anaemia and malarial fevers in infants in Tanzania: a randomised placebo-controlled trial. *The Lancet*. 2003 May 31;361(9372):1853-60.
25. Bonita R, Beaglehole R, Kjellström T. Basic epidemiology. *World Health Organization*; 2006.
26. Müller, O., Traoré, C., Becher, H., & Kouyaté, B. (2003). Malaria morbidity, treatment-seeking behaviour, and mortality in a cohort of young children in rural Burkina Faso. *Tropical Medicine & International Health*, 8(4), 290-296.
27. Hill, Z., Kendall, C., Arthur, P., Kirkwood, B., & Adjei, E. (2003). Recognizing childhood illnesses and their traditional explanations: exploring options for

380 care-seeking interventions in the context of the IMCI strategy in rural Ghana. *Tropical*
381 *Medicine & International Health*, 8(7), 668-676.

382 28. Malik, E. M., Hanafi, K., Ali, S. H., Ahmed, E. S., & Mohamed, K. A. (2006).
383 Treatment-seeking behaviour for malaria in children under five years of age:
384 implication for home management in rural areas with high seasonal transmission in
385 Sudan. *Malaria journal*, 5(1), 60.

386 29. Fenny, A. P., Hansen, K. S., Enemark, U., & Asante, F. A. (2014). Quality of
387 uncomplicated malaria case management in Ghana among insured and uninsured
388 patients. *International journal for equity in health*, 13(1), 63.

389 30. USAID/CDC/QHP, GHS/NMCP, WHO: Ghana – Nationwide Malaria Health Facility
390 Survey. Accra, Ghana: Quality Health Partners; 2008.

391 31. Zoyah, C. J., & Diedong, A. L. (2014). Health Professionals' Compliance with
392 National Treatment Guidelines for Treatment of Severe Malaria in Children in
393 Bolgatanga Hospital, Upper East, Ghana

394 32. Buabeng, K. O., Duwiewua, M., Dodoo, A. N., Matowe, L. K., & Enlund, H. (2007).
395 Self-reported use of anti-malarial drugs and health facility management of malaria in
396 Ghana. *Malaria journal*, 6(1), 85.

397 33. Chandler, C. I., Whitty, C. J., & Ansah, E. K. (2010). How can malaria rapid
398 diagnostic tests achieve their potential? A qualitative study of a trial at health facilities
399 in Ghana. *Malaria journal*, 9(1), 95.


400 34. Kwarteng, A., Asante, K. P., Abokyi, L., Gyaase, S., Febir, L. G., Mahama, E.,
401 Konadu, D.G, Tawiah, T, Adu-Gyasi, D, Dosoo, D, Amenga-Etego S, Ogutu, B,
402 Owusu-Agyei, S. (2015). Provider compliance to artemisinin-based combination
403 therapy at primary health care facilities in the middle belt of Ghana. *Malaria journal*,
404 14(1), 361.

35. National Malaria Control Programme. (2013) Annual Report. Accra, Ghana.
https://www.ghanahealthservice.org/downloads/ghana_malaria_programme_review_final_report_june_2013.pdf accessed Jan 2018
36. Adjei, A.A, Winch, P, Laar, A, Sullivan, D.J Jr, Sakyi, K.S, Stephens, J.K, Adjei, G.O, Boateng, I.A, Aubyn, V.N, Kubio C, Tuakli, J, Vanotoo L, Bortei, B.B, Amo-Addae, M, Sorvor, F, Coleman, N, Dalglish, S, Owusu, R, Gebreyesus, T, Essuman E, Greene R, Ankomah E, Houston K, Bart-Plange C, Salamat S, Addison EA, Quakyi, I. A. (2016). Insights into the Affordable Medicines Facility-malaria in Ghana: the role of caregivers and licensed chemical sellers in four regions. *Malaria journal*, 15(1), 263.
37. Ferrer, B.E., Webster, J., Bruce, J., Narh-Bana, S.A., Narh, C.T., Allotey, N.K., Glover, R., Bart-Plange, C., Sagoe-Moses, I., Malm, K. and Gyapong, M. (2016). Integrated community case management and community-based health planning and services: a cross sectional study on the effectiveness of the national implementation for the treatment of malaria, diarrhoea and pneumonia. *Malaria journal*, 15(1), 340.
38. Ameme, D. K., Afari, E. A., Nyarko, K. M., Malm, K. L., Sackey, S., & Wurapa, F. (2014). Direct observation of outpatient management of malaria in a rural Ghanaian district. *The Pan African medical journal*, 19.
39. Oladosu, O. O., & Oyibo, W. A. (2012). Overdiagnosis and overtreatment of malaria in children that presented with fever in Lagos, Nigeria. *ISRN Infectious Diseases*, 2013.
40. Okoro, C. I., Chukwuocha, U. M., Nwakwuo, G. C., & Ukaga, C. N. (2015). Presumptive Diagnosis and Treatment of Malaria in Febrile Children in Parts of South Eastern Nigeria. *Journal of Infectious Diseases & Therapy*. 3: 240

41. Orish VN, Ansong JY, Onyeabor OS, Sanyaolu AO, Oyibo WA, Iriemenam NC.(2016). Overdiagnosis and overtreatment of malaria in children in a secondary healthcare centre in Sekondi-Takoradi, Ghana. *Trop Doct*, 46(4):191-8.
42. Masquelier B, Reniers G, Pison G. (2014).Divergences in trends in child and adult mortality in sub-Saharan Africa: survey evidence on the survival of children and siblings. *Popul Stud (Camb)*, 68(2):161-77.
43. Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J. E., Cousens, S, Mathers, C, Black, R. E. (2015). Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *The Lancet*, 385(9966), 430-440.
44. Murphy, S.C, Breman, J.G. (2001). Gaps in the childhood malaria burden in Africa: cerebral malaria, neurological sequelae, anemia, respiratory distress, hypoglycemia, and complications of pregnancy. *Am J Trop Med Hyg*, 64(1-2 Suppl):57-67.
45. Tette, E. M., Neizer, M. L., Nyarko, M. Y., Sifah, E. K., Sagoe-Moses, I. A., & Nartey, E. T. (2016). Observations from Mortality Trends at The Children’s Hospital, Accra, 2003-2013. *PloS one*, 11(12), e0167947.
46. Berkley, J.A., Brent, A., Mwangi, I., English, M., Maitland, K., Marsh, K., Peshu, N. and Newton, C.R., (2004) Mortality among Kenyan children admitted to a rural district hospital on weekends as compared with weekdays. *Pediatrics*, 114(6):1737-8.

456


457 Figure Legends



458

459 Figure 2: Percentage mortality (all causes) from the year 2010 to 2012


460



461

462 Figure 3: Malaria case fatality rate

463



464

465 Figure 4 Malaria testing for year 2010 to 2012

466

List of prescribed medications	2010			2011			2012		
	Malaria positive (%)	Malaria negative (%)	Not done (%)	Malaria positive (%)	Malaria negative (%)	Not done (%)	Malaria positive (%)	Malaria negative (%)	Not done (%)
Artemether injection, Antibiotics and Antipyretics	2 (33.3)	3 (50)	1 (16.7)	1 (16.7)	3 (50)	2 (33.3)	3 (17.6)	10 (58.8)	4 (23.5)
Antibiotics	0 (0)	4 (66.7)	2 (33.3)	0 (0)	5 (62.5)	3 (37.5)	0 (0)	17 (65.4)	9 (34.6)
Antibiotics and Antipyretics	0 (0)	5 (62.5)	3 (37.5)	1 (8.3)	4 (33.3)	7 (58.3)	0 (0)	6 (85.7)	1 (14.3)
Artemether-lumefantrine, Antibiotics and Antipyretics	0 (0)	4 (66.7)	2 (33.3)	0 (0)	3 (100)	0 (0)	1 (6.6)	10 (66.7)	4 (26.7)
Artemether-lumefantrine and Antipyretics	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)
Artesunate-amodiaquine, Antibiotics and Antipyretics	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	2 (100)	0 (0)
Artesunate-amodiaquine and Antipyretics	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Artemether Injection, artemether-lumefantrine and Antipyretics	3 (33.3)	3 (33.3)	3 (33.3)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Quinine and Antipyretics	0 (0)	0 (0)	1 (100)	0 (0)	1 (50)	1 (50)	1 (12.5)	4 (50)	3 (37.5)
Antipyretics	0 (0)	0 (0)	0 (0)	0 (0)	1 (50)	1 (50)	0 (0)	2 (66.7)	1 (33.3)
Artemether-lumefantrine and Antibiotics	0 (0)	0 (0)	1 (100)	3 (33.3)	3 (33.3)	3 (33.3)	0 (0)	3 (60)	2 (40)
Artemether injection, artemether-lumefantrine and Antibiotics	0 (0)	3 (100)	0 (0)	0 (0)	2 (100)	0 (0)	0 (0)	8 (80)	2 (20)
Quinine, Antibiotics and Antipyretics	1 (6.7)	4 (26.7)	10 (66.7)	1 (10)	5 (50)	4 (40)	23 (37.7)	20 (32.8)	18 (29.5)
Artemether injection, Artesunate Amodiaquine and artemether-lumefantrine	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)
Artemether injection, artemether-lumefantrine, Quinine and Antibiotics	1 (100)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)	0 (0)	0 (0)	0 (0)
Artemether injection and Antipyretics	0 (0)	1 (50)	1 (50)	1 (25)	1 (25)	2 (50)	2 (28.6)	2 (28.6)	3 (42.9)
Artemether injection, Quinine, Antibiotics and Antipyretics	0 (0)	0 (0)	0 (0)	0 (0)	2 (50)	2 (50)	0 (0)	0 (0)	0 (0)
Artemether-lumefantrine, Quinine and Antipyretics	0 (0)	2 (100)	0 (0)	0 (0)	1 (100)	0 (0)	1 (12.5)	6 (75)	1 (12.5)
Artesunate-amodiaquine, Quinine, Antibiotics and Antipyretics	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (100)

467

468 Figure 5: List of combination of the prescribed medications during the study stratified by year

469




Figure 6: Other non-malaria causes of death.

Table 1: General characteristics of study population

Characteristics	Number (n=223)	Percentage (%)
Sex		
Male	122	54.7
Female	101	45.3
Age (Years)		
<1	110	49.3
2-5	63	28.3
>5	50	22.4
Haemoglobin		
<7 g/dL	76	34.1
<10 g/dL	75	33.6
<11 g/dL	30	13.5
≥11 g/dL	42	18.8
Sickle cell*		
HB (AS)	21	15.8
HB (SS)	6	4.5
HB (AA)	106	79.7
Malaria parasite	40	17.9
MP Seen	112	50.2
No MP Seen	71	31.9
Not done		
Death on admission in hospital (Hours)		
<24	97	43.5
<72	83	37.2
>72	43	19.3

*Sickle cell (n=133). MP = Malaria parasites

Table 2: Patients characteristics stratified by malaria laboratory diagnosis

Clinical and demographic characteristics	Malaria positive n = 40 (%)	Malaria Negative N=112(%)	Not done N=71(%)	P-VALUE
Age (Mean ± SD)	3.1 ± 3.1	2.9 ± 3.2	2.5± 3.5	0.6
Age (Years)				
<1	15(37.5)	57(50.9)	38(53.5)	0.01
2-5	20(50)	28 (25.0)	15(21.1)	
6-11	5(12.5)	27 (24.1)	18(25.4)	
Sex				
Male	24 (60)	66 (59)	32(45.1)	0.23
female	16 (40)	46 (41)	39(54.9)	
Hemoglobin				
				0.18
<7g/dL	22 (55)	36 (32.1)	18(25.4)	
<10g/dL	10 (25)	43 (38.4)	22(31.0)	
<11g/dL	4 (10)	12 (10.7)	14(19.7)	
≥11g/dL	4 (10)	21 (18.8)	17(23.9)	
Death on admission in hospital (Hours)				
<24	12 (30)	40 (35.7)	45(63.4)	0.001
24-72	25 (62.5)	38 (33.9)	20(28.2)	
>72	3 (7.5)	34 (30.4)	6 (8.4)	
Blood transfusion				
Yes	22 (55)	36 (32.1)	17(23.9)	0.02
No	18 (45)	76 (67.9)	54(76.1)	
Presumptive diagnosis for malaria				
Severe malaria anaemia	12 (30)	30 (26.8)	10(14.08)	0.004
Cerebral malaria	16 (40)	29 (25.9)	14(19.7)	
Severe malaria	10 (25)	19 (16.9)	7(9.9)	
Bronchopneumonia +malaria	2 (5)	17 (15.2)	8(11.3)	
Epilepsy + malaria	0 (0)	17 (15.2)	3(4.2)	
Antimalaria treatment given				
Yes	40(100)	93(83.0)	57(80.3)	0.05
No	0(0)	19(17)	9(12.7)	

*Significant P-Value= (P<0.05); SD=Standard Deviation