

Short Research Article

Synthesis and Crystal Structure of an Air Stable Primary Phosphine-Manganese (+1) Complex; *fac*-Tricarbonyl-tris(*o*-florophenylphosphine)-Manganese(+1)-hexafluorophosphate.

ABSTRACT

The synthesis of an air-stable manganese(+I)-primary phosphine complex from *fac*-[Mn(CO)₃(CH₃CN)₃]PF₆ and (*o*-FC₆H₄)PH₂ is described. The X-ray single crystal structure of the manganese(I) complex is reported.

Keywords: [((*o*-FC₆H₄)PH₂)₃Mn(CO)₃]PF₆, crystal structure, synthesis, complex, X-ray.

1. INTRODUCTION

In comparison with other types of organophosphine metal derivatives, the number of known primary phosphine transition metal complexes is relatively small. This is partially due to some of the chemical and physical properties of primary phosphines. Generally speaking, primary phosphines are easily oxidized by air (in some cases they can ignite spontaneously if not handled under an atmosphere of oxygen free inert gas), and have high toxicity. They are, however, versatile starting materials due to the possibility of functionalising the hydrogen bonds. For this reason, there is increasing interest in the synthesis and coordination chemistry of substituted primary phosphines containing ortho-substituted aryl group [such as (*o*-F-C₆H₄)PH₂], particularly for applications in the synthesis of macrocyclic ligands containing three or more phosphorous atoms using the synthetic procedure called a metal template [1].

In this synthetic method, one of the acidic hydrogen atoms of the metal coordinated-PH₂ group and the ortho-flouro group can easily be activated and then eliminated as a small molecule; HF. As a result, new P-C bonds between the already metal-coordinated phosphine moieties can be formed. This synthetic technique leads to the formation of a metal complex containing a macrocyclic phosphine ligand. For example, a series of triphosphine 9 and 12 membered macrocycles containing a variety of alkyl and aryl phosphine groups have been produced using the metal-template procedure [2].

Generally, the macrocycles-metal complexes produced are kinetically stable. This may explain why the liberation of free macrocycles containing phosphorous (III) is not always achieved [3]. Edwards et al [4-5], showed that the liberation of free phosphorous (III) containing macrocycles can be obtained if the centre of the precursor macrocyclic complex is the ion Mn(+1). For this reason the title complex, [((*o*-FC₆H₄)PH₂)₃Mn(CO)₃]PF₆, was prepared and characterized by a variety of techniques.

2. EXPERIMENTAL DETAILS

2.1 Synthesis

To a stirred dry methanoic solution (25 ml, 0.001 mmol., 0.4gm.) of [Mn(CO)₃(CH₃CN)₃]PF₆ (prepared as described in [6-7] kept under a positive pressure of nitrogen gas, a dry methanoic solution of *o*-flourophenylphosphine (25 ml, 0.004 mmol.; 0.52gm) was slowly added and then stirred continuously for two days at room temperature. The resulting orange yellow slurry was filtered and the yellow sticky solid

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

washed twice with diethyl ether (15 ml each). To the solid obtained, 60 ml of dry methanol was added. The reaction mixture was heated at 50 °C with stirring until all solid dissolved and a clear orange yellow solution obtained. The solution was allowed to cool extremely slowly down to room temperature. The very pale yellow cuboid-like crystals obtained were then collected by filtration. A second batch of crystals can be obtained when the filtrate was kept at -25 °C for two days. The total mass of the pale yellow crystals was 0.6 gm (yield ≈ 90%), M.P. = 142°C.

Elemental analysis for $[(o\text{-FC}_6\text{H}_4)\text{PH}_2]_3\text{Mn}(\text{CO})_3\text{PF}_6$ calculated M.W.: 668.062; C: 37.76, and H: 2.69. Found: C:37.80, and H: 2.63. The electrospray mass spectrum of the complex shows peaks at m/z = 522.9816 (calc.=522.9801), which corresponds to the ion $(M\text{-PF}_6)^+$.

^{31}P NMR (none solvent, 300 MHz, ppm): = -37.8356 (assigned to coordinated PH_2 group).

IR (KBr pellet, cm^{-1}): weak, 2367.68; strong sharp, 2051.89; strong sharp, 1995.96; strong sharp, 1978.12; shoulder, 1941; strong medium 1603.04; medium sharp, 1570.25; strong sharp, 1476.24; strong sharp 1448.28; strong sharp, 1269.9; strong sharp 1219.76; small sharp, 1162.38; medium sharp, 1128.74; medium sharp, 1086.69; medium sharp, 1074.64; small sharp, 1028.84; shoulder, 947.84; strong sharp 893.84; shoulder, 874.07; strong broad, 836.47; shoulder, 813.33; strong sharp, 773.797; strong sharp 765.6; strong sharp, 759.8; shoulder, 740.04; medium sharp, 707.26; weak sharp 675.9; strong sharp, 654.7; strong sharp, 621.448; strong sharp, 556.8; shoulder, 543.3; weak sharp, 509.6; weak sharp, 503; weak sharp, 480.2; weak sharp, weak sharp, 456.1; weak sharp, 444.5; weak sharp, 413.4. A single crystal structure analyses confirmed the molecular formula. The complex is air stable, moderately soluble in methanol and ethanol but very soluble in acetonitrile, THF, and CH_2Cl_2 .

2.2 Crystal structure determination

Single-crystal XRD data were collected on an Agilent SuperNova Dual Atlas diffractometer with a mirror monochromator using Cu (λ = 1.5418 Å) radiation and equipped with an Oxford Cryosystems cooling apparatus. Crystal structures were solved and refined using SHELX.[8]. Non-hydrogen atoms were refined with anisotropic displacement parameters. Hydrogen atoms were inserted in idealized positions, and a riding model was used with U_{iso} set at 1.2 times the value of U_{eq} for the atom to which they are bonded. Structure determination and refinement data are shown in Table 1.

Table 1. Crystal data and structure refinement data for $[(o\text{-FC}_6\text{H}_4)\text{PH}_2]_3\text{Mn}(\text{CO})_3\text{PF}_6$.

Empirical formula	$\text{C}_{21}\text{H}_{18}\text{F}_9\text{MnO}_3\text{P}_4$
Formula weight	668.17
Temperature	150(2) K
λ	1.54184 Å
Crystal system	Triclinic
Space group	P -1
Unit cell dimensions	$a = 10.4907(3)$ Å, $b = 10.9299(5)$ Å $c = 12.2565(5)$ Å, $\alpha = 71.670(4)^\circ$ $\beta = 89.807(3)^\circ$, $\gamma = 85.696(3)^\circ$.
Volume	$1329.99(9)$ Å ³
Z	2
Density (calculated)	1.668 Mg/m^3
Absorption coefficient	7.130 mm^{-1}
F(000)	668
Crystal size	$0.260 \times 0.230 \times 0.160 \text{ mm}^3$
Theta range for data collection	4.227 to 73.411°
Index ranges	$-12 \leq h \leq 10$, $-12 \leq k \leq 13$, $-15 \leq l \leq 15$
Reflections collected	9171
Independent reflections, R_{int}	5166, 0.0166

Formatted: Highlight

Formatted: Highlight

Completeness to theta = 67.684°	99.5 %
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	1.00000 and 0.75518
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	5166 / 0 / 343
Goodness-of-fit on F2	1.074
Final R indices [I>2sigma(I)]	R ₁ = 0.0403, wR ₂ = 0.1138
R indices (all data)	R ₁ = 0.0418, wR ₂ = 0.1153

3. RESULTS AND DISCUSSION

The cationic complex of manganese (I), *fac*-tricarbonyl-tris(*o*-florophenylphosphine)-manganese(I), was prepared by a single full displacement reaction of the kinetically liable acetonitrile ligands from the *fac*-[Mn(CO)₃(CH₃CN)₃] PF₆ by three groups of (*o*-FC₆H₄)PH₂. To make sure that all the three acetonitrile ligands were displaced, a 4:1 mole ratio of the phosphine ligand : manganese precursor was used. The insolubility of the new manganese(I) complex in diethyl ether helped to remove any unreacted phosphine, then the yellow solid was extract with hot methanol. The resulting hot yellow filtrate was allowed to cool down slowly to room temperature producing air stable yellow crystals, suitable for single-crystal X-ray structure analysis, in good yield. Elemental microanalysis and the sharpness of the measured melting point confirmed the high purity of the crystalline materials. The observation of the [(*o*-FC₆H₄)PH₂]₃Mn(CO)₃⁺ ion in the mass spectrum as well as the clear melt obtained when the complex was heated in open air indicate the thermodynamic stability of the complex in both the solid and liquid states.

In the IR spectrum of the complex, the weak absorption at about 2368 cm⁻¹ is assigned to the stretching of the -PH₂ group, whereas absorptions at 2052; 1996 ;1978; 1941 cm⁻¹ are attributed to the various modes of carbon monoxide stretching[4].

Atomic coordinates and displacement parameters for [(*o*-FC₆H₄)PH₂]₃Mn(CO)₃ PF₆ obtained from structure refinement are shown in Table 2.

Table 2. Atomic coordinates and displacement parameters for [(*o*-FC₆H₄)PH₂]₃Mn(CO)₃]PF₆. Displacement parameters are in the form: -2π²[h²a²U¹¹ + ... + 2 h k a* b* U¹²].

	x	y	z	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
C(1)	0.2587(2)	0.9883(2)	0.3198(2)	0.034(1)	0.022(1)	0.026(1)	-0.007(1)	-0.005(1)	0.002(1)
C(2)	0.1886(3)	0.9801(3)	0.4167(2)	0.038(1)	0.037(1)	0.039(1)	-0.015(1)	0.004(1)	-0.005(1)
C(3)	0.2203(3)	1.0351(3)	0.4985(2)	0.062(2)	0.042(2)	0.032(1)	-0.016(1)	0.008(1)	-0.004(1)
C(4)	0.3293(3)	1.1012(3)	0.4830(2)	0.072(2)	0.029(1)	0.036(1)	-0.014(1)	-0.007(1)	-0.008(1)
C(5)	0.4024(3)	1.1120(3)	0.3879(3)	0.052(2)	0.037(1)	0.049(2)	-0.016(1)	-0.001(1)	-0.017(1)
C(6)	0.3683(3)	1.0566(2)	0.3061(2)	0.044(1)	0.030(1)	0.035(1)	-0.010(1)	0.004(1)	-0.007(1)
C(7)	0.1263(2)	0.5150(2)	0.1190(2)	0.023(1)	0.029(1)	0.027(1)	-0.008(1)	-0.007(1)	-0.002(1)
C(8)	0.0852(2)	0.4059(2)	0.1992(2)	0.024(1)	0.033(1)	0.029(1)	-0.008(1)	-0.005(1)	0.000(1)
C(9)	0.0855(3)	0.2879(3)	0.1814(2)	0.047(2)	0.026(1)	0.043(1)	-0.004(1)	-0.014(1)	0.001(1)
C(10)	0.1330(3)	0.2773(3)	0.0789(3)	0.053(2)	0.039(2)	0.055(2)	-0.026(1)	-0.025(1)	0.019(1)
C(11)	0.1781(3)	0.3826(3)	-0.0018(3)	0.035(1)	0.071(2)	0.040(2)	-0.033(2)	-0.008(1)	0.010(1)
C(12)	0.1739(2)	0.5011(3)	0.0169(2)	0.031(1)	0.054(2)	0.027(1)	-0.012(1)	-0.005(1)	-0.007(1)
C(13)	0.5573(2)	0.5683(3)	0.2220(2)	0.032(1)	0.037(1)	0.035(1)	-0.003(1)	0.001(1)	0.004(1)
C(14)	0.5580(3)	0.4490(3)	0.2056(3)	0.035(1)	0.045(2)	0.049(2)	-0.009(1)	0.004(1)	0.001(1)
C(15)	0.6544(3)	0.3520(3)	0.2492(3)	0.050(2)	0.037(2)	0.063(2)	-0.004(1)	0.020(2)	0.003(1)
C(16)	0.7519(3)	0.3759(3)	0.3119(3)	0.034(1)	0.055(2)	0.052(2)	0.014(1)	0.008(1)	0.010(1)
C(17)	0.7548(3)	0.4936(3)	0.3314(3)	0.030(1)	0.062(2)	0.041(2)	0.006(1)	-0.001(1)	0.002(1)
C(18)	0.6580(3)	0.5904(3)	0.2871(2)	0.031(1)	0.049(2)	0.037(1)	-0.001(1)	0.000(1)	-0.003(1)

C(19)	0.1174(3)	0.6694(2)	0.3831(2)	0.037(1)	0.030(1)	0.034(1)	-0.002(1)	-0.013(1)	0.000(1)
C(20)	0.2917(2)	0.5067(3)	0.3493(2)	0.031(1)	0.032(1)	0.041(1)	-0.008(1)	-0.013(1)	-0.001(1)
C(21)	0.3682(2)	0.7008(2)	0.3979(2)	0.032(1)	0.029(1)	0.031(1)	-0.006(1)	-0.007(1)	0.000(1)
O(1)	0.0302(2)	0.6546(2)	0.4390(2)	0.041(1)	0.053(1)	0.045(1)	-0.002(1)	0.002(1)	-0.004(1)
O(2)	0.3136(2)	0.3990(2)	0.3907(2)	0.051(1)	0.027(1)	0.067(1)	0.000(1)	-0.019(1)	0.004(1)
O(3)	0.4383(2)	0.7119(2)	0.4639(2)	0.043(1)	0.052(1)	0.037(1)	-0.013(1)	-0.017(1)	-0.005(1)
F(1)	0.0837(2)	0.9131(2)	0.4322(2)	0.052(1)	0.091(2)	0.078(1)	-0.051(1)	0.026(1)	-0.032(1)
F(2)	0.0419(2)	0.4182(2)	0.2996(1)	0.042(1)	0.051(1)	0.033(1)	-0.012(1)	0.008(1)	-0.008(1)
F(3)	0.4610(2)	0.4273(2)	0.1445(2)	0.056(1)	0.055(1)	0.079(1)	-0.031(1)	-0.009(1)	0.000(1)
F(4)	0.1776(4)	0.9263(3)	-0.2107(2)	0.203(4)	0.076(2)	0.067(2)	-0.035(1)	-0.027(2)	-0.036(2)
F(5)	0.1464(2)	1.1348(2)	-0.2414(2)	0.075(1)	0.068(1)	0.066(1)	0.014(1)	-0.014(1)	0.020(1)
F(6)	0.2786(3)	1.1182(3)	-0.0909(2)	0.124(2)	0.084(2)	0.065(1)	-0.036(1)	0.010(1)	-0.048(2)
F(7)	0.3108(3)	0.9067(3)	-0.0616(2)	0.102(2)	0.078(2)	0.058(1)	-0.001(1)	-0.022(1)	0.042(1)
F(8)	0.1166(2)	0.9948(3)	-0.0639(2)	0.059(1)	0.111(2)	0.055(1)	-0.010(1)	0.018(1)	-0.016(1)
F(9)	0.3433(2)	1.0532(3)	-0.2377(2)	0.061(1)	0.160(3)	0.049(1)	-0.021(2)	0.021(1)	0.013(2)
P(1)	0.2190(1)	0.9027(1)	0.2218(1)	0.037(1)	0.024(1)	0.026(1)	-0.007(1)	-0.010(1)	0.003(1)
P(2)	0.1221(1)	0.6660(1)	0.1501(1)	0.041(1)	0.025(1)	0.038(1)	-0.008(1)	-0.020(1)	0.000(1)
P(3)	0.4261(1)	0.6903(1)	0.1685(1)	0.035(1)	0.032(1)	0.033(1)	-0.005(1)	-0.006(1)	0.005(1)
P(4)	0.2296(1)	1.0219(1)	-0.1522(1)	0.040(1)	0.034(1)	0.024(1)	-0.006(1)	0.000(1)	0.004(1)
Mn(1)	0.2576(1)	0.6817(1)	0.2905(1)	0.027(1)	0.022(1)	0.027(1)	-0.006(1)	-0.011(1)	0.001(1)

106
107 In the crystal structure of *fac*-tricarbonyl-tris(*o*-florophenylphosphine)-manganese(I) hexafluorophosphate
108 (Fig 1), the manganese coordination is distorted octahedral with mean Mn-P distances in the range
109 2.298(1) Å -2.299(1)Å and Mn-C distances in the range 1.827(3) Å - 1.842(1) Å and angles in the range
110 87.17(8)°– 94.41(11)°. Selected bond lengths and angles are in Table 3.

111 The cation is structurally related to *fac*-tricarbonyl-tris(phenylphosphine)-molybdenum(0) (but with a
112 different metal, oxidation state and contains fluoro substituents)[9]. Similar cation coordination is
113 observed in *fac*-tris(1-methylethenylphosphine)-tricarbonyl-chromium(0) and *fac*-tris(1-
114 methylethenylphosphine)-tricarbonyl-molybdenum(0) [10].




Fig. 1. The asymmetric unit showing 50% thermal ellipsoids for non-hydrogen atoms in $[(o\text{-FC}_6\text{H}_4)\text{PH}_2]_3\text{Mn}(\text{CO})_3\text{PF}_6$.

Table 3. Selected bond lengths [Å] and angles [°] for $[(o\text{-FC}_6\text{H}_4)\text{PH}_2]_3\text{Mn}(\text{CO})_3\text{PF}_6$.

C(1)-P(1)	1.806(2)	C(2)-F(1)	1.347(3)
C(7)-P(2)	1.804(2)	C(13)-P(3)	1.813(3)
C(14)-F(3)	1.344(4)	C(19)-O(1)	1.132(4)
C(19)-Mn(1)	1.842(3)	C(20)-O(2)	1.131(3)
C(20)-Mn(1)	1.827(3)	C(21)-O(3)	1.136(3)
C(21)-Mn(1)	1.828(2)	P(1)-Mn(1)	2.2985(7)
P(2)-Mn(1)	2.2912(7)	P(3)-Mn(1)	2.2991(8)
C(2)-C(1)-P(1)	121.42(19)	C(6)-C(1)-P(1)	121.48(19)
C(14)-C(13)-P(3)	121.2(2)	C(18)-C(13)-P(3)	120.6(2)
C(20)-Mn(1)-C(21)	88.89(11)	C(20)-Mn(1)-C(19)	89.27(12)
C(21)-Mn(1)-C(19)	94.41(11)	C(20)-Mn(1)-P(2)	93.17(8)
C(21)-Mn(1)-P(2)	177.42(8)	C(19)-Mn(1)-P(2)	87.17(8)
C(20)-Mn(1)-P(1)	177.93(8)	C(21)-Mn(1)-P(1)	89.05(8)
C(19)-Mn(1)-P(3)	176.42(8)	P(2)-Mn(1)-P(3)	89.59(3)

In the crystal, o-fluorophenylphosphine groups of neighbouring cations are involved in both edge-to-face and π - π interactions. Two edge-to-face interactions with a centroid-to-centroid distance of 4.87(1) Å link pairs of molecules as shown in Fig. 2. A similar arrangement is observed in the crystal structure of fac-tricarbonyl-tris(phenylphosphine)-molybdenum(0).[9] For each cation in the Mn complex (I), one of the o-fluorophenyl phosphine groups already involved in edge-to-face contact also participates in π - π contact

(centroid-to-centroid contact of 3.77(1) Å) with an inversion related ring from a neighbouring molecule leading to the formation of ribbons parallel to the *a*-axis.




Fig. 2. The crystal structure of $[(o\text{-FC}_6\text{H}_4)\text{PH}_2]_3\text{Mn}(\text{CO})_3\text{PF}_6$ showing edge-to-face and $\pi\text{-}\pi$ contacts as dotted lines. The PF_6^- counterions and hydrogen atoms have been omitted for clarity.

4. CONCLUSION

The kinetic **trans** effect of carbon monoxide ligands has facilitated the full displacement of all three acetonitrile groups in one reaction. As a result, the entering ligands (namely, *o*-fluorophenylphosphine) have taken a facial arrangement which is an essential requirement for any successful template cyclisation attempt of the coordinated *o*-fluorophenylphosphine ligands.

REFERENCES

1. Atherton MJ, Fawcett J, Holloway JH, Hope EG, Karacar A, Russell DR, et al. Carbon-fluorine and -hydrogen bond activation and carbon-carbon bond formation in η^5 -pentamethylcyclopentadienyl-rhodium and -iridium phosphine complexes; crystal structures of $[\text{M}(\eta^5\text{-C}_5\text{Me}_5)\text{Cl}][(\text{C}_6\text{F}_5)_2\text{PCH}_2\text{CH}_2\text{P}(\text{C}_6\text{F}_5)_2]^+\text{BF}_4^-$ ($\text{M} = \text{Rh}$ or Ir). *J Chem Soc, Dalton Trans.* **1996**(15):3215-20.
2. Battle AR, Edwards PG, Haigh R, Hibbs DE, Li D, Liddiard SM, et al. Synthesis and characterization of iron(II) complexes of 10- and 11-membered triphosphamacrocycles. *organometallics.* **2007**;26(2):377-86.
3. Edwards PG, Haigh R, Li D, Newman PD. Template synthesis of 1,4,7-triphosphacyclononanes. *J Am Chem. Soc.* 2006;128(11):3818-30.
4. Edwards PG, Mahon MF, Newman PD, Reixach E, Zhang W. Manganese(I) templates for the construction of benzannulated triphosphamacrocycles. *Dalton Trans.* 2014;43(41):15646-55.

Formatted: Subscript

Formatted: Subscript

Formatted: Highlight

Formatted: Highlight

155 5. Edwards PG, Kariuki BM, Newman PD, Zhang W. Peripheral methyl activation in η^4 -1,2,3,4-
156 tetramethylcyclobutadienecobalt complexes: template synthesis and subsequent reactivity of
157 triphosphamacrocycles. *Organometallics*. 2014;33(19):5440-7.
158 6. Reimann RH, Singleton E. Simple route to new cationic carbonyl compounds of manganese and
159 rhenium. *J Organometal Chem*. 1973;59:C24-C6.
160 7. Edwards DA, Marshalsea J. The synthesis and spectral features of some cationic manganese(I) and
161 rhenium(I) tricarbonyl complexes. *J Organomet Chem*. 1977;131(1):73-91.
162 8. Sheldrick GM. A short history of SHELX. *Acta Crystallogr, Sect A: Found Crystallogr*. 2008;64(1):112-
163 22.
164 9. Campbell T, Gibson AM, Hart R, Orchard SD, Pope SJA, Reid G. Synthesis, spectroscopic and
165 structural characterisation of chromium(0), molybdenum(0) and tungsten(0) complexes involving primary
166 and secondary phosphines. *J Organomet Chem*. 1999;592(2):296-305.
167 10. Edwards PG, Fleming JS, Liyanage SS, Coles SJ, Hursthouse MB. Primary alkenyl phosphine
168 complexes of chromium and molybdenum; synthesis and characterization of tricarbonyl(1,5,9-
169 triphosphacyclododecane)chromium(0). *J Chem Soc, Dalton Trans*. 1996(9):1801-7.