

Synthesis, Characterization and Antibacterial studies of some Isoniazid-derived Schiff bases

ABSTRACT

Aims: To synthesize some isoniazid-derived imines by the condensation of isoniazid (INH) with some substituted benzaldehydes as well as pyridine carboxaldehydes and to determine their structures by spectroscopic methods.

Methodology: Isoniazid was mixed with equimolar amounts of some substituted benzaldehydes as well as pyridine carboxaldehydes and refluxed for 4-7 hours. The solid products were filtered by gravity and dried over anhydrous CaCl_2 in a desiccator. The ligands were screened for the biological activities against *Salmonella typhi*, *Staphylococcus aureus*, *Escherichia coli*, *Shigella flexneri*, and *Salmonella enterica* by paper disc diffusion method.

Results: Nine Schiff bases were synthesized and characterized by elemental analyses, ^1H NMR, and ^{13}C NMR, and single crystal X-ray crystallography. Their melting temperatures were determined and their purity ascertained by HPLC.

Conclusion: Nine isoniazid-derived imines were synthesized and characterized, two of which were new (compound 6 and 8). The antibacterial activities of some of the imines were comparable to the activity of isoniazid against *E. coli*, whereas the other Schiff bases were more active than isoniazid against the other tested pathogens. Compound 7 showed good antibacterial activity compared to isoniazid against all the tested pathogens and compound 4 was inactive against all pathogens.

Keywords: isoniazid, hydrazones, Schiff bases, antibacterial

1. INTRODUCTION

Isoniazid is a pro-drug with established therapeutic applications and has been in clinical use for almost half a century as a front-line drug for prophylaxis and treatment of tuberculosis [1,2]. It is a nicotinamide analog made up of a pyridine ring and a hydrazide group [3]. The free $-\text{NH}_2$ group in the hydrazine moiety can be condensed with various carbonyl compounds (aldehydes and ketones) to form hydrazones (Schiff bases), containing the characteristic $\text{H}-\text{C}=\text{N}-\text{NH}-$ group. The presence of a carbonyl group adjacent to the $-\text{NH}-$ group in hydrazine provides a possibility of tautomerism in some cases [4]. It is well established that the presence of OH or NH groups in an appropriate position relative to the $-\text{C}=\text{N}$ group provides greater

chances for increased intramolecular as well as intermolecular interactions [5]. This implies that hydrazones derived from isoniazid may show greater biological activities due to improved interactions with the binding sites.

Hydrazones have been reported to possess antitumoral, antimicrobial, antitubercular, anticonvulsant, analgesic antifungal, antiviral and antibacterial activities [6]. Mainsah *et al.* [7] recently reported, for the first time, the anti-onchocercal activity of some isoniazid-derived Schiff bases and their copper(II) and zinc(II) complexes with the copper(II) complexes showing significant activity against both micro- and macrofilariae.

Antimycobacterial activities of Schiff bases derived from isoniazid or possess a pyridine moiety have also been reported [8,9]. Numerous studies have pointed out the importance of developing novel isoniazid hydrazones as promising anti-tubercular agents [10]. Chemical modification of the hydrazine portion of isoniazid with a functional group that blocks acetylation, while maintaining strong biological action has the potential to improve clinical outcomes and reduce the emergence of acquired isoniazid resistance in patients. Thus, hydrazones, which contain the -HN=N=CH- moiety from isoniazid [11], constitute an important class of compounds for drug development.

Schiff bases derived from isoniazid have been synthesized by classical methods in which the aldehydes and ketones are refluxed with isoniazid in organic solvents [12,13], in a catalytic amount of acetic acid. Other environmentally friendly and less costly synthetic methods have been attempted including green synthesis of hydrazones using natural fruit-derived organic acids as catalysts [14, 15]

In continuation of our research on the isoniazid-derived hydrazones, the present study describes the synthesis, characterization of some imines and their antibacterial activity on selected pathogens.

2. EXPERIMENTAL

2.1 Materials

All chemicals used were analytical grade (Sigma-Aldrich), and used as purchased without further purification.


2.2 Physical Measurements

^1H -NMR and ^{13}C -NMR spectra were recorded with a 600 MHz Bruker Advanced II spectrophotometer in dimethylsulfoxide (DMSO-d_6) as solvent and were reported relative totetramethylsilane (TMS) as internal standard. CHN contents of the hydrazones were performed using the FLASH 2000 Organic Elemental Analyzer, CHNS-O analyzer (Thermo Scientific), melting temperatures were determined using a Mel-Temp II laboratory device, while infrared (IR)spectroscopy was performed on a Perkin-Elmer FT-IR 100 spectrometer. The spectra were recorded from 4000 to 600 cm^{-1} in attenuated total reflectance (ATR) mode.HPLC chromatograms of the compounds were obtained on the HEWLETT PACKARD, Agilent 1100 series, using the OpenLABChemStation, 2010-2014 software, in acetonitrile/water (1:9) solvent.

2.3 Synthesis of the Schiff bases

The Schiff bases were obtained by refluxing equimolar proportions of isoniazid and the various aldehydes in methanol as solvent at 70°C with continuous stirring for 4-7 hours as shown in scheme 1. For example, isoniazid (0.274 g, 2.0 mmol) was dissolved in 5.0 mL of warm methanol and to this solution was added to 0.258 g (2.0 mmol) of 4-(dimethylamino)benzaldehyde, dissolved in 5.0 mL of warm methanol, as well (scheme 1). The mixture was refluxed at 70°C, with continuous stirring for 4 hours in a water bath, resulting in a yellow solution. The progress of the reaction was monitored by TLC using a 40% isopropyl alcohol/60% hexane solvent system. The solution was allowed to cool as the solvent evaporated and yellow crystals were obtained after two days. [4,16-18]. The product was filtered, washed with cold methanol, and dried in a desiccator over anhydrous CaCl_2 . The same compound was synthesized from the isonicotinic acid hydrazide salt in an ethanol/water solvent system and refluxed for 24 hours [16].The

83 purity of the compounds was ascertained by HPLC analysis and melting temperature determination
84 (Table 2).





85


86 Scheme 1: General method of synthesis of Schiff Bases

87 The aldehydes used for the synthesis of the Schiff bases are shown in Table 1.

88 Table 1. Aromatic aldehydes used for synthesis of Schiff Bases


Compound	Structural formula and name of ArCHO
1	 4-dimethylaminobenzaldehyde
2	 2-methoxybenzaldehyde

3




3-methoxybenzaldehyde

4




4-methoxybenzaldehyde

5




2,3-dimethoxybenzaldehyde

6




2,3,5-trimethoxybenzaldehyde

7




pyridine-2-carboxaldehyde

8



9



89

90 Table 2 Physical properties of the Schiff bases

Compound (molwt)	Color	mp (°C)	Yield (%)	Elemental Analysis					
				Calculated			Experimental		
				%C	%H	%N	%C	%H	%N
1 (286.3)	yellow	205	72.99	62.94	5.59	19.58	63.54	5.56	19.92
2 (255.2)	colorless	190	83.35	65.62	5.13	16.46	65.87	5.04	16.45
3 (255.2)	colorless	195	78.47	65.87	5.13	16.46	65.91	4.98	16.50
4 (255.2)	colorless	160	82.00	61.54	5.49	15.38	62.99	4.18	15.92
5 (285)	colorless	138	93.78	63.16	5.26	14.74	63.47	5.32	14.57
6 (315)	yellow	193	99.28	60.95	5.40	13.33	60.91	5.42	13.28
7 (226.2)	white	175	66.86	63.71	4.46	24.77	63.46	4.31	24.80
8 (226.2)	white	238	79.09	63.71	4.46	24.77	63.77	4.27	24.80
9 (226.2)	white	235	88.27	63.71	4.46	24.77	62.55	4.37	24.68

91

92

93

2.3.1 Characterization Data of the Schiff Bases

N'-(4-(dimethylamino)benzylidene)isonicotinoylhydrazone monohydrate (1): ^1H NMR (δ ppm): 11.76 (s, 1H, NH); 8.76 (d, 2H, pyr); 8.32 (s, 1H, =C-H), 7.80 (d, 2H, pyr); 7.56 (d, 2H, benz); 6.74 (d, 2H, benz); 3.01 (s, 6H, $\text{N}(\text{CH}_3)_2$). ^{13}C NMR (δ ppm): 161.5 (C-7); 152.2 (C-14); 150.7 (C-2, C-6); 150.3 (C-10); 141.3 (C-4); 129.1 (C-12, C-16); 121.9 (C-3, C-5); 121.7 (C-11); 112.3 (C-13, C-15); 40.2 (C-18, C-19). IR (cm^{-1}): 3385 (O-H); 3190 (N-H, str); 2939 (=C-H, str); 1738 (s, amide II) 1662 (C=O); 1603 (C=N); 1588 (C=C).

N'-(2-methoxybenzylidene)isonicotinoylhydrazone (2): ^1H NMR (δ ppm): 12.06 (s, 1H, NH); 8.84 (s, 1H, =C-H); 8.78 (d, 2H, pyr); 7.90 (d, 1H, benz), 7.85 (d, 2H, pyr); 7.43 (t, 1H, benz); 7.11 (d, 1H, benz) 7.04 (t, 1H, benz); 3.86 (s, 3H, OCH_3). ^{13}C NMR (δ ppm): 161.4 (C-7); 157.9 (C-16); 150.3 (C-2, C-6); 144.5 (C-10); 140.4 (C-4); 131.9 (C-14); 125.6 (C-12); 122.0 (C-13); 121.5 (C-3, C-5); 120.8 (C-11); 111.9 (C-15); 55.7 (C-17). IR (cm^{-1}): 3195 (N-H, str); 1651 (C=O); 1598 (C=C); 1550 (C=N); 1252 (C-O-).

N'-(3-methoxybenzylidene)isonicotinoylhydrazone (3): ^1H NMR (δ ppm): 12.09 (s, 1H, N-H); 8.78 (d, 2H, pyr); 8.45 (s, 1H, N=C-H); 7.83 (d, 2H, pyr); 7.38 (t, 1H, benz); 7.31 (d, 2H, benz); 7.02 (dd, 1H, benz); 3.80 (s, 3H, OCH_3). ^{13}C NMR (δ ppm): 162.2 (C-7); 160.0 (C-13); 150.8 (C-2, C-6); 149.5 (C-10); 140.9 (C-4); 135.9 (C-11); 130.5 (C-15); 122.0 (C-3, C-5); 120.7 (C-16). 117.0 (C-14); 111.9 (C-12); 55.6 (C-17). IR (cm^{-1}): 3188 (N-H, str); 3011 (=C-H, str); 1677 (C=O); 1582 (C=N); 1276 (C-O-).

N'-(4-methoxybenzylidene)isonicotinoylhydrazone (4): ^1H NMR (δ ppm): 11.95 (s, 1H, N-H); 8.78 (dd, 2H, pyr); 8.41 (s, 1H, N=C-H); 7.82 (dd, 2H, pyr); 7.70 (dd, 2H, benz); 7.02 (dd, 2H, benz); 3.80 (s, 3H, OCH_3). ^{13}C NMR (δ ppm): 161.5 (C-7); 161.1 (C-14); 150.3 (C-2, C-6); 149.0 (C-10); 140.6 (C-4); 129.0 (C-12, C-16); 126.6 (C-11); 121.5 (C-3, C-5); 114.4 (C-13, C-15); 55.3 (C-17). IR (cm^{-1}): 3435 (O-H, str); 3139 (=C-H, str); 1653 (C=O); 1597 (C=N); 1253 (C-O-).

N'-(2,3-dimethoxybenzylidene)isonicotinoylhydrazone (5): ^1H NMR (δ ppm): 12.12 (s, 1H, NH); 8.79 (d, 2H, pyr); 8.78 (s, 1H, =C-H); 7.86 (d, 2H, pyr); 7.51 (dd, 1H, benz); 7.13 (m, 2H, benz); 3.83 (s, 3H, OCH_3); 3.80 (s, 3H, OCH_3). ^{13}C -NMR (δ ppm): 162.0 (C-7); 153.1 (C-16); 150.8 (C-2, C-6); 148.7 (C-15);

119 145.2 (C-10); 140.8 (C-4); 127.9 (C-13); 122.0 (C-3, C-5); 117.6 (C-11); 115.0 (C-14); 61.7 (C-17); 56.2
120 (C-18). IR(cm^{-1}): 3455(N-H, str); 3147 (=C-H, str); 1659 (C=O); 1557 (C=N); 1262 (C-O-)

121 ***N'-(2,3,5-trimethoxybenzylidene)isonicotinoylhydrazone (6)***: ^1H NMR (δ ppm): 11.96 (s, 1H, NH);
122 8.77 (m, 2H, pyr); 8.76 (s, 1H, =C-H); 7.84 (d, 2H, pyr); 7.40 (s, 1H, benz); 6.74 (s, 1H, benz); 3.86 (d, 6H,
123 OCH_3); 3.76 (s, 3H, OCH_3). ^{13}C -NMR (δ ppm): 161.7 (C-7); 154.1 (C-13); 152.8 (C-15); 150.7 (C-2, C-6);
124 145.2 (C-16); 143.7 (C-10); 141.0 (C-4); 121.9 (C-3, C-5); 113.7 (C-11); 108.2 (C-12); 98.2 (C-14); 56.9
125 (C-19); 56.3 (C-17); 56.2 (C-18). IR (cm^{-1}): 3568 (O-H, str); 3197 (=C-H); 1663 (C=O); 1595 (C=C); 1554
126 (C=N); 1276, 1222, 1206 (3C-O-).

127 ***N'-(pyridine-2-carboxaldehyde)isonicotinoylhydrazone (7)***: ^1H NMR (δ ppm): 12.24 (s, 1H, -NH); 8.80
128 (d, 2H, INH); 8.63 (d, 1H, pyr); 8.48 (s, 1H, -N=C-H); 7.99 (d, 1H, pyr); 7.90 (td, 1H, pyr); 7.83(d, 2H,
129 INH); 7.44 (ddd, 1H, pyr). ^{13}C -NMR (δ ppm): 161.9 (C-7); 152.9 (C-12); 150.4 (C-2, C-6); 149.6 (C-14);
130 149.2 (C-11); 140.2 (C-4); 137.0(C-15); 124.7 (C-16); 121.5 (C-3, C-5); 120.1 (C-17). IR (cm^{-1}): 3296 (N-
131 H, str); 1672 (C=O); 1584 (C=N); 1546 (C=C).

132 ***N'-(pyridine-3-carboxaldehyde)isonicotinoylhydrazone (8)***: ^1H NMR (δ ppm): 12.20 (s, 1H, -NH); 8.88
133 (s, 1H, pyr); 8.78 (d, 2H, INH); 8.63 (dd, 1H, pyr); 8.51 (s, 1H, -N=C-H); 8.17 (d, 1H, pyr); 7.83 (d, 2H,
134 INH); 7.51 (dd, 1H, pyr). ^{13}C -NMR (δ ppm): 161.9 (C-7); 151.0 (C-15); 150.3 (C-2, C-6); 148.9 (C-13);
135 146.3 (C-11); 140.3 (C-4); 133.6 (C-17); 129.9 (C-12); 124.0 (C-16); 121.5 (C-3, C-5). IR (cm^{-1}): 3180 (N-
136 H, str); 1673 (C=O); 1590 (C=N); 1547 (C=C).

137 ***N'-(pyridine-4-carboxaldehyde)isonicotinoylhydrazone (9)***: ^1H NMR (δ ppm): 12.32 (s, 1H, -NH); 8.80
138 (d, 2H, INH); 8.67 (d, 2H, pyr); 8.46 (s, 1H, -N=C-H); 7.84 (d, 1H, INH); 7.68 (d, 2H, pyr); ^{13}C -NMR(δ
139 ppm): 162.0 (C-7); 150.4 (C-2,C-6); 150.3 (C-14,C-16); 146.7 (C-11); 140.2 (C-12); 121.6 (C-3,C-5);
140 120.8 (C-13, C-17). IR (cm^{-1}): 3181 (N-H, str); 1690 (C=O); 1572 (C=N); 1546 (C=C).

141 2.4 Biological activity

In vitro antibacterial activities of the ligands were evaluated by the disc–diffusion method. Mueller – Hinton agar was employed as microbial growth medium. The antibacterial diffusion tests were carried out as described by Berghe and Vlientink [19] using a cell suspension of about 1.5×10^6 CFU/mL obtained from the McFarland Turbidity Standard N° 0.5. Antibacterial activity was evaluated by measuring the diameter of the inhibition zone (IZ) around the discs. Three replicates were made for each sample and mean values of the growth inhibition zone were measured. Compounds were considered active only when inhibition zone was greater than 6 mm as indicated in Table 3.

Table 3: Antibacterial screening of isoniazid derivatives by disc diffusion method

Compound	<i>E. coli</i> BL-21	<i>S. enterica</i> NR-4311	<i>Salmonellatyphi</i>	<i>S. flexneri</i>	<i>S. aureus</i> NR-46003
1	10	-	-	-	-
2	10	-	10	-	-
3	9	-	9	-	-
4	-	-	-	-	-
5	10	-	-	-	9
6	-	-	-	-	-
7	11	10	11	7	10
8		-	10	-	-
9	11	-	10	-	-
INH	10	-	-	-	-
DMSO	-	-	-	-	-
gentamicin	24	*		*	*
chloramphenicol	*	25	*	*	*

*Positive control not tested

3. RESULTS AND DISCUSSION

3.1 Elemental analysis

The physical properties of the isoniazid-derived Schiff bases including molecular weight, color, melting point and elemental analysis are tabulated in Table 2. The experimental and theoretical values of elemental analysis agree very closely. The compounds had sharp melting points which differ from those of the starting materials confirming that new compounds were formed. These high melting temperatures indicate that the molecules are held together in a lattice by strong intermolecular hydrogen bonds.

3.2 Infrared

The average stretching frequency of N-H band of the compounds ranges from 3139 to 3296 cm^{-1} and the characteristic $\text{C}=\text{N}$ (imine) peak varies between 1603 and 1550 cm^{-1} . The band due to NH-stretching in 6 occurs at 3197 cm^{-1} which is lower than that of its isomer, N'-(3,4,5-trimethoxybenzylidene)isonicotinoylhydrazone whose NH stretching frequency falls in the 3350 to 3230 cm^{-1} region [20]. The NH band of mono-substituted methoxyaldehydes shows a slight increase due to resonance effects with the *p*-substituted Schiff base having the highest frequency and the *m*-substituted isomer the lowest. Some strong bands are clustered around the 1500 cm^{-1} range in all the compounds and this can be assigned to the $\text{C}=\text{N}$ and $\text{C}=\text{C}$ bonds of the pyridine/aromatic rings. The $\text{O}-\text{H}$ bands which appear in compounds 1, 4, 5 and 6 can be attributed to water molecules trapped in the crystal. It has been reported that compound 5 traps chloroform when it is slowly crystallized in this solvent as indicated by X-ray diffraction analysis [21].

3.3 ^1H and ^{13}C NMR

A sharp singlet centered on 3.80 ppm in Schiff bases formed from methoxy-substituted aldehydes is assigned to methyl protons. The dimethylaminobenzaldehyde Schiff base also gives a singlet at δ 3.01 attributed to methyl protons. The aromatic protons for both the pyridine and phenyl ring range from 6.74 to 8.88 ppm in the hydrazones. The azomethine proton, which confirms the formation of the product, appears as a sharp singlet in the range 8.32 – 8.78 ppm [20, 22] while the N-H proton can be assigned to the signals which appear at 11.76 to 12.32 ppm range. The ^{13}C NMR confirms the formation of the hydrazones with the azomethine carbon at δ 143.7- 150.3. The chemical shift of the carbonyl carbon varies very little from 160.0 – 162.0 ppm in the Schiff bases. The presence of the O-H protons in

compounds **1**, **4**, **5** and **6** is confirmed by the chemical shifts in the 3.38 – 3.49 ppm range contrary to similar compounds [20].

3.3 Antibacterial Activity

From the data, the zones of inhibition for the Schiff bases are relatively low (7-11 mm) when compared to those of the standard antibiotics, gentamicin and chloramphenicol. This suggests weak antibacterial activity. However, the activities of these Schiff bases are comparable to those of the parent compound, isoniazid, which in itself is an antibiotic of proven therapeutic applications. *E. coli* and *S. typhi* were the most susceptible the synthesized ligands. Compound **7** was the most active and showed activity against all the bacterial strains used and compound **4** was inactive against all strains. Compound **7** proved to be better than the parent isoniazid against all the bacterial strains. The Minimum Inhibitory Concentration (MIC) results for these compounds that were active against at least two bacterial strains were very high, that is 1000 µg/mL. The MIC of INH was greater than 1000 µg/mL, implying that the synthesized compounds had better antibacterial properties than the parent compound although the bacterial species tested are not the primary target of INH.

4.0 CONCLUSION

Nine Schiff bases derived from isoniazid have been synthesized in good yield and analytical purity. They were characterized by elemental analyses, ¹H and ¹³C NMR spectroscopy. Compounds **6** and **8** are reported for the first time. Compound **7** was the most biologically active and showed activity against all the bacterial strains used and compound **4** was inactive against all strains. Compound **7** proved to be a better antibacterial than the parent compound, isoniazid against all the bacterial strains. The biological activities of the Schiff bases showed that some of the compounds synthesized have the potential for the development of good therapeutic agents against a few selected bacterial strains.

REFERENCES

1. Goodman LS, Gilman A. The Pharmacological basis of therapeutics. 3rded. New York: Macmillan Company; 1965.



- 205 2. Kolyva AS, Karakousis PC. Old and New TB drugs: mechanisms of action and resistance,
206 understanding tuberculosis - New approaches to fighting against drug resistance. Pere-Joan
207 Cardona (Ed.), ISBN: 978-953-307-948-6. Intech DOI: 10.5772/2477; 2012.
- 208 3. Rodrigues FAR, Oliveira ACA, Cavalcanti BC, Pessoa C, Pinheir AC, De Souza MVN. Biological
209 evaluation of isoniazid derivatives as an anticancer class. Scientia Pharmaceutica, 2014; 82:21–
210 28.
- 211 4. Mainsah EN, Ndifon PT, Nfor EN, Njapba JN. Synthesis, characterization and antibacterial
212 properties of some transition metal complexes of (1H-pyrrol-2-yl)isonicotinoylhydrazone. Bull.
213 Chem. Soc. Ethio.2013; 27(3):395-404.
- 214 5. BerkesiO, KörtvélyesiT, HetényiC, Németh T, Pálinkól. Hydrogen bonding interaction of
215 benzylidene type Schiff bases studied by vibrational spectroscopic and computational methods.
216 Chemical Phys. 2003; 5:2009-2014.
- 217 6. Elumalai K, Ali MA, Elumalai M, Eluri K, Srinivasan S. Novel isoniazid cyclocondensed 1,2,3,4-
218 tetrahydropyrimidine derivatives for treating infectious diseases: a synthesis and *in vitro* biological
219 evaluation. J. Acute Dis. 2013; 2013:316-321
- 220 7. Mainsah EN, Ntun S-JE, Samje M, Cho-Ngwa F, Ndifon PT, Yong JN. Synthesis and Anti-
221 onchocercal Activity of Isonicotinoylhydrazones and their Copper(II) and Zinc(II) Complexes. Anti-
222 Infective Agents. 2016;14(1):47-52
- 223 8. Ramadevi P, Singh R, Prajapati A, Gupta S., Chakraborty D. Cu(II) complexes of isoniazid Schiff
224 bases: DNA/BSA binding and cytotoxicity studies on A549 cell Line. Adv. Chem.2014, ID 276598:
225 [http:// dx. doi.org /10.1155/2014/276598](http://dx.doi.org/10.1155/2014/276598).
- 226 9. Ahmed RM, Yousif EI, Al-Jeboori MJ. Co(II), Cd(II) complexes derived from heterocyclic Schiff
227 bases: synthesis, structural characterisation, and biological activity. The Sci. World J.2013: ID
228 754868: [http:// dx.doi. org 10.1155/2013/754868](http://dx.doi.org/10.1155/2013/754868).
- 229 10. Judge V. Narasimhan B, Ahuja M, Sriram D, Yogeewari P, Clercq ED, Pannecouque C,
230 Balzarini J. Synthesis, antimycobacterial, antiviral, antimicrobial activities, and QSAR studies of

- isonicotinic acid-1-(substituted phenyl)-ethylidene/cycloheptylidenehydrazides. Med. Chem. Res. 2012; 21:1935–1952.
11. Mahapatra S, Woolhiser LK, Lenaerts AJ, Johnson JL, Eisenach KD, Joloba ML, Boom WH, Belisle JT. A novel metabolite of antituberculosis therapy demonstrates host activation of isoniazid and formation of the isoniazid-NAD⁺ adduct. Antimicrobial Agents and Chemo. 2012; 56(1):28-35
12. Hearn MJ, Cynamon MH. Design and synthesis of antituberculars: preparation and evaluation against *Mycobacterium tuberculosis* of an isoniazid Schiff base. J. Antimicrobial Chemo. 2004; 53:185–191.
13. Eldehna WM, Fares M, Abdel-Aziz MM, Abdel-Aziz HA. Design, synthesis and antitubercular activity of certain nicotinic acid hydrazides. Molecules. 2015; 20:8800-8815.
14. Yadav G, Mani JV. Green Synthesis of Schiff Bases by Natural Acid Catalysts. Int. J. Sci. Res., 2015;4(2):121-127.
15. Voguri HB, Venkateswararao A, Ashok M. Synthesis, Characterization and Biological properties of Schiff Base and Transition Metal Complexes. Int. J. Chem. Pharm Sci., 2015, 3(1):1453-1457
16. Sahebalzamani H, Khsligh N, Ghammamy S, Salimi F and Mehrani K. Crystal structure and density functional theory study on structural properties and energies of an isonicotinohydrazide compound. Molecules. 2011; 16:7715-7724.
17. Kumar PP, Rani BL. Synthesis and Characterization of New Schiff bases containing pyridine moiety and their derivatives as antioxidant agents. Int. J. ChemTech Res., 2011, 3(1):155-160.
18. Parekh J., Inamdhar P., Nair R., Baluja S., Chanda S. Synthesis and antibacterial activity of some Schiff bases derived from 4-aminobenzoic acid. J. Serb. Chem Soc. 2005, 70(10):1155-1161.
19. Berghe VA and Vlietinck AJ. Screening Methods for Antibacterial and Antiviral Agents from Higher Plants, Methods for Biochemistry, 1991;6:47-68.
20. Agarwal RK Singh L, Sharma DK, Singh R. Synthesis, Spectral and Thermal Investigations of Some Oxovanadium(IV) Complexes of Hydrazones of Isonicotinic Acid Hydrazide. Turk. J. Chem. 2005; 29:309 – 316.

- 258 21. Peralta MA, de Souza MNV, Wardell SMSV, Wardell JL, Low JN, Glidewell C. 2,3-
259 Dimethoxybenzaldehyde isonicotinoylhydrazone chloroform monosolvate, and the mono- and
260 dihydrates of 3,4,5-trimethoxy-benzaldehyde isonicotinoyl-hydrazone: hydrogen-bonded
261 supramolecular structures in one, two and three dimensions. Acta Cryst. (2007). C63, o68 - o72.
262 DOI: 10.1107/S0108270106052486
- 263 22. Hussain Z, Yousif E, Ahmed A, Altaie A. Synthesis and Characterization of Schiff bases of
264 Sulfamethoxazole. Org. Med. Chem. Let. 2004;4(1) DOI: 10.11862/2191-2858-4-1



Structural formula of Aldehyde

Structural formula and Name of Isoniazid-derived product





N'-(4-dimethylaminobenzylidene)isonicotinoylhydrazone monohydrate

(1), CAS Registry Number 13059-77-3




N'-(2-methoxybenzylidene)isonicotinoylhydrazone (2)


CAS Registry Number 6342-48-9



N'-(3-methoxybenzylidene)isonicotinoylhydrazone (3)





CAS Registry Number 303066-20-8





N'-(4-methoxybenzylidene)isonicotinoylhydrazone (**4**)

CAS Registry Number 893-42-5





N'-(2,3-dimethoxybenzylidene)isonicotinoylhydrazone (**5**)

CAS Registry Number 71579-89-0





N'-(2,3,5-trimethoxybenzylidene)isonicotinoylhydrazone (6)





N'-(pyridine-2-carboxyldehyde)isonicotinoylhydrazone (7)

CAS Registry Number 15017-32-0



N'-(pyridine-3-carboxyldehyde)isonicotinoylhydrazone (8)

CAS Registry Number 15017-31-9



N'-(pyridine-4-carboxaldehyde)isonicotinoylhydrazone (**9**)

CAS Registry Number 13025-99-5