

Original Research Article**Relationships among phenotypic Traits of Giant African Land Snails in Western region of Nigeria.****ABSTRACT**

The relatedness among three populations (*A. achatina*, *A. fulica* and *A. marginata*) of giant African land snails, a highly relished source of protein in West Africa was studied. Three hundred (300) matured snails consisting of one hundred (100) each of *A. achatina*, *A. fulica* and *A. marginata* collected randomly from three locations in Western region of Nigeria were used to evaluate the relationships among phenotypic traits. The results of the mean body weight and shell traits (body weight, shell length, shell width, shell 'mouth' length and shell 'mouth' width) studied showed highly significant differences ($P < 0.05$). Phenotypic correlations among the pairs of traits showed positive, strong and highly significant ($P < 0.01$) correlation coefficients (r_p). Similarly, the clustering of the phenotypic traits also showed that the three populations were divided into two major groups with *A. marginata* being more closely related to *A. achatina* than *A. fulica*. The body traits studied was grouped into one principal component with a total variability of 98.75%. Snail populations used in this study showed very wide phenotypic relationship, suggesting that they may come from a common ancestor.

Key words: Phenotypic, relationship, snail, trait, African Land Snails.

INTRODUCTION

The intake of animal protein in the diet of an average Nigerian has been reducing from year to year (Nworgu and Fapohunda, 2002). In Africa, Nigeria is still among the least consumer of animal protein, consuming on the average about 6g to 8.4g by an individual per day (Egbunike, 1994). This may be attributed to non-availability and high cost of the product which has impoverished Nigerians (Nworgu and Fapohunda, 2002). Owing to these, a common man finds it difficult to buy animal products from conventional animals. Besides, the cost of production inputs of conventional animal protein sources like fish, chicken and beef have become highly expensive for most Nigerians (Hodasi, 1983). Efforts are being made towards producing animal protein from unconventional animals like snails and grass-cutters by domesticating them because of the shortage of meat protein from conventional animal sources (Okon *et al.*, 2009). Nigeria is richly endowed with various species of snails and it is the cheapest animal protein source during the rainy season, especially for rural dwellers in the rainforest zone of Nigeria (Ojating, 2001; Okon *et al.*, 2012a). In the past, snails were cheap because of their enormous supply and small numbers of individuals were interested in their meat (Udedibie, 1989). In the quest to meet daily protein intake, the consumption of snail meat has increased tremendously in Nigeria over the years (Ojating and Ogar, 2002). Giant African land snails have unique traits and characteristics such as adaptation to extreme climatic conditions by aestivation process that could prove fundamental to the food security of future generations.

Upon all these great benefits derived from snails, they are still being gathered from the wild indiscriminately and excessively. This, together with deforestation and bush burning, has

drastically reduced giant African land snails population, thus subjecting the species to the threat of extinction (Ojating and Ogar, 2002; Okon *et al.*, 2008; Okon and Ibom, 2012).

Giant African land snails consist of the following breeds; *Archachatina marginata*, *Achatina fulica*, *Achatina achatina*, *Limicolaria species* and *Thapsia species* in Nigeria and other African countries as stated in Okon *et al.* (2012a) and they belong to the family Achatinidae, consisting of about thirteen genera. These snail breeds could be black or white with respect to their skin colouration [Omole *et al.*, (2007) and Ibom (2009)]. This difference in the colour of the skin could be genetic in origin, and capable of causing remarkable differences in a population (Ibom *et al.*, 2011).

Nisbet (1974) and Plummer (1975) stated that *Achatina achatina* is the largest land gastropod. However, Okon *et al.* (2012 b) reported that *Archachatina marginata* is the largest known land snails in Nigeria in the Guinness Book of records. Differences exist among the populations of snails in Nigeria which could be chromosomal, genetic and/ or phenotypic and animal breeders and geneticists can exploit it for the species improvement. The performance of animals including snails depends on the environment where they are found and their genetic make-up (Ibom *et al.*, 2014). The Western states in Nigeria have different environmental conditions (relative humidity, vegetation, daylight hours and temperature and rainfall pattern). These differences in environmental conditions could give rise to diversity in gene pool of snails, which will eventually influence the body components and growth pattern of the organisms including the snail species that inhabit the area. The knowledge of phenotypic relationship among giant African land snails is so important for the understanding of economic traits of the snail species. This study therefore seeks to evaluate the relationships among phenotypic traits of giant African land snails in western region of Nigeria (Osun, Ondo and Oyo States).

MATERIALS AND METHODS

Sample collection and snail management

The experiment was conducted at the Botanical garden, University of Calabar, Calabar. A total of three hundred (300) mature giant African land snails consisting of 100 each of *Achatina fulica* (Plate 1), *Achatina achatina* (Plate 2) and *Archachatina marginata* (Plate 3) respectively, were procured from three different locations (Ile – Ife forest in Osun State, 7⁰.55¹N, 4⁰.43¹E, Oluwa forest reserve in Ondo State, 6⁰.82¹N, 4⁰.67¹E and Amusan’s Snailery in Oyo State, 7⁰.38¹N, 3⁰.95¹E) of western region of Nigeria for phenotypic traits studies.

The snails were kept alive and maintained under moist conditions to ensure their survival. They were fed daily with fresh paw-paw leaves and fruits, and succulent plant parts before the commencement of the experiment. The snails were managed intensively in condemn drums kept under shade to provide suitable environment for the snails. Loamy soil was collected, made un-compacted by sun-drying, heated in an oven at a temperature of 65⁰C to kill all inhabited organisms. The loamy soil had a pH range of 7.00 to 8.00 as measured by pH meter (p1367 SIGMA), as snails show preference for neutral to slightly alkaline pH soils (Ibom, 2009). The soil in the drums was turned regularly with care to enhance burrowing activities of the snails. Proper sanitation was duly carried out daily.



Plate 1: *A. fulica*



Plate 2: *A. achatina*



Plate 3: *A. marginata*

PLATES 1-3: The three species of giant African land snails

Phenotypic measurement of snails

Parameters that were measured as phenotypic traits included; Body weight (BDW), Shell length (SHL), Shell width (SHW), Shell 'mouth' length (SML) and Shell 'mouth' width (SMW). Shell length and width, and Shell 'mouth' length and width were measured using Vernier caliper (MITUTOYO 500-752-10, Spain), while body weight was measured using Pro-ScoutST electronic scale with sensitivity of 0.01g.

Statistical analysis

The mean phenotypic traits comparison among/within the three species of giant African land snails was carried out using 3 × 3 factorial under a Complete Randomized Design (CRD) in statistical package for Social Science (SPSS) software version 20.0. Data collected were transformed using the log transformation method in order to reduce possible biases in the measurements. Significant means were separated using Least Significant Difference (LSD) at

0.05 probability level. The data were further subjected to correlation, principal component analysis (PCA) and clustering analysis.

RESULTS AND DISCUSSION

The weight of the snails used ranged from 128g to 221g (*Achatina achatina*), 36g to 57g (*Achatina fulica*) and 165g to 390g (*Archachatina marginata*). The results of mean phenotypic traits of giant African land snails obtained from different populations in this study are presented on Table 1. The results revealed that there were significant differences ($P < 0.05$) in all the parameters investigated. It however, showed that the *A. marginata* obtained from Ondo State recorded the highest body weight (328.33g), which was not significantly different from the ones from Oyo state (291.00g). It was also observed that all other phenotypic traits were highest in *A. marginata* from all the locations (Table 1). Mean body weights reported in this study for the three species; *Achatina achatina* (181.67g, 176.67g and 169g), *Achatina fulica* (45.67g, 41g, and 40.67g), *Archachatina marginata* (328.33g, 291g and 258.33g) are significantly higher than the mean body weights of 120.90g for *A. marginata*, 93.70g for *A. achatina* and 79.80g for *A. fulica* respectively reported by Ibom *et al.* (2014). These differences in mean body weights may be due to differences in the locations, types of diets fed, size, physiological stage and age of the snails used. Snails used by Ibom *et al.* (2014) were growers and were gathered from central agro-ecological zone of Delta State, while the snails used in this study were adults and were gathered from three locations (Osun, Ondo and Oyo) States in the western part of Nigeria. Results of phenotypic traits for the three snail breeds in this study were not in line with the views of Nisbet (1974) and Plummer (1975) that the largest land gastropod is *Achatina achatina*. The results, however, agreed with that of Omole (1998), Olawoyi and Ogogo (2006), Okon *et al.* (2012b) and Anon (2013), who stated that *Archachatina marginata* is the largest known land snail in Nigeria.

The results of correlation analysis revealed that all the phenotypic traits (body weight, shell length, shell widths, shell 'mouth' lengths and shell 'mouth' widths) measured were significant ($p < 0.01$) and positively/highly correlated. The highest correlation value was recorded between shell length (SHL) and body shell width (SHW) ($r = 0.992$) (Table 2). The phenotypic correlations recorded in this study for all the pairs of traits were positively and closely correlated. The observed correlations and positive values among these pairs of traits could suggest that the traits are linked and that improvement in one trait might lead to improvement in the other (Etukudo *et al.*, 2015). The correlation results recorded in this study is in agreement with the works of Ibom (2009) and Ibom *et al.* (2014) who reported positive and close correlations among snail traits. The results also agreed with the view of Ehiobu and Kyado (2000), that correlation could be high or low and positive or negative between traits.

TABLE 1

 Mean phenotypic traits of giant African land snails obtained from different populations

Traits	Osun State			Ondo State			Oyo State		
	<i>Achatina Achatina</i>	<i>Achatina Fulica</i>	<i>Archachatina marginata</i>	<i>Achatina achatina</i>	<i>Achatina fulica</i>	<i>Archachatina marginata</i>	<i>Achatina achatina</i>	<i>Achatina fulica</i>	<i>Archachatina Marginata</i>
Body weight (g)	169±18.02 ^b	45.67±5.55 ^c	258.33±38.69 ^a	176.67±23.14 ^b	40.67±0.88 ^c	328.33±35.96 ^a	181.67±14.80 ^b	41±3.51 ^c	291±66.34 ^a
Body shell length (cm)	12.77±0.67 ^b	8.17±0.22 ^c	14.43±0.99 ^a	12.70±0.80 ^b	7.93±0.69 ^c	16.07±0.84 ^a	13.13±0.59 ^b	7.83±0.24 ^c	15.97±0.74 ^a
Body shell width (cm)	6.67±0.24 ^b	4.40±0.15 ^c	8.07±0.58 ^a	6.87±0.69 ^b	4.30±0.12 ^c	8.63±0.45 ^a	6.93±0.16 ^b	4.73±0.23 ^c	8.83±0.48 ^a
Mouth shell length (cm)	6.90±0.21 ^b	4.93±0.07 ^c	8.97±0.29 ^a	7.13±0.18 ^b	4.97±0.09 ^c	9.0±0.43 ^a	6.93±0.05 ^b	4.83±0.39 ^c	8.87±0.34 ^a
Mouth shell width (cm)	4.70±0.25 ^{ab}	2.87±0.18 ^c	5.50±0.45 ^a	4.47±0.29 ^b	2.57±0.09 ^c	5.47±0.32 ^a	4.37±0.11 ^b	2.73±0.13 ^c	5.70±0.30 ^a


^{abc} Means bearing different superscripts along the same row are significantly different ($p < 0.05$) 

Table 2: The results of Pooled correlation between body weight and other phenotypic traits of the snails' species evaluated.

	BDW	SHL	SHW	SML	SMW
BDW	1.000	0.988**	0.989**	0.987**	0.973**
SHL		1.000	0.992**	0.975**	0.986**
SHW			1.000	0.985**	0.986**
SML				1.000	0.982**
SMW					1.000

** = Highly significant level ($P < 0.01$)

BDW = Body weight, SHL = Shell length, SHW = Shell width, SML = Shell 'mouth' length, MSW = Shell 'mouth' width.

Results of the principal component analysis (PCA) showed that only one principal component was extracted where all the phenotypic traits contributed 98.75% to the total variability (Table 3). The highest component matrix was obtained from shell width (0.997), while the lowest was from shell 'mouth' length and width (0.992). The communality ranged from 0.983 to 0.989 (Table 3). The variability obtained for the snails was explained by the contribution of the body traits from PCA analysis. All the body traits contributed highly to the total variability of 98.75% recorded in this study. According to Udeh and Ogbu (2011), communality values of principal component analysis indicate the fit of PCA in handling a morphometric dataset. It estimates the total amount of variance that an original variable shares with all other variables included in the analysis. The high communality values obtained in this study might suggest strong contribution of these traits to the variation observed in the three species.

Table 3: The results of Principal component analysis of evaluated traits

Phenotypic traits	Communality	PC ₁
Eigen value	-	4.94
Proportion of variance	-	98.75
Cumulative variance	-	98.75
Body weight	0.987	0.994
Shell length	0.989	0.995
Shell width	0.994	0.997
Shell 'mouth' length	0.984	0.992
Shell' mouth' width	0.983	0.992

The results of the cluster analysis for phenotypic data revealed two major clusters. Cluster 2 was divided into two sub-clusters consisting of *A. achatina* and *A. marginata*. The closeness of *A. marginata* to *A. achatina* phenotypically might be due to the closeness in sizes of the two species of snails as reported by Ibom *et al.* (2014) that *A. marginata* and *A. achatina* are closely related in terms of weight and their sizes.




Fig. 1: The results of cluster analysis of the phenotypic traits evaluated for the three species of giant African land snails.

Key:

A.f-D = *Achatina fulica* from Ondo State; A.f-I = *Achatina fulica* from Oyo State; A.f-O = *Achatina fulica* from Osun State; A.a-D = *Achatina achatina* from Ondo State; A.a-I = *Achatina achatina* from Oyo State; A.a-O = *Achatina achatina* from Osun State; A.m-D = *Archachatina marginata* from Ondo State; A.m-I = *Archachatina marginata* from Oyo State; A.m-O = *Archachatina marginata* from Osun State.

CONCLUSION

This study indicated ^{was} wide morphological variability among the species of snails investigated. The populations notwithstanding, showed that *A. marginata* and *A. achatina* were more closely related morphologically than *A. fulica*. Snails ^{from} other locations should also be used to assess the relationships among phenotypic traits of giant African land snails in Nigeria. Also, molecular markers should be employed to confirm these relationships.

REFERENCES

Anon (2013). Contribution of discussants on snail types and sizes in Africa. *Proceedings of the 2nd International Conference of Giant African Land Snails*. Abeokuta, Nigeria. Pp. 30-32.

Egbunike, G. N. (1994). What is Animal Science? And can Nigeria get out of malnutrition. In: Iyayi, E. A., Adeshehinwa, A. O. K. and Bamgbose, A. M. (eds.). *Proceedings of 2nd*

- Annual Conference, Animal Science Association of Nigeria (ASAN), Ikeja, Nigeria. Pp.1-12.*
- Ehiobu, N. G. and Kyado, J. A. (2000). Heritability, repeatability and genetic correlation of Swine. *Proceedings of the 25th Annual Conference of Nigerian Society for Animal Production (NSAP)*. Umudike, Nigeria. Pp.260-261.
- Etukudo, O. M., BIna-ibor, O. and Ekerette, E.E. (2015). Comparative study of egg quality traits of black-skinned and white-skinned ectotypes of snails (*Archachatina marginata*) based on four whorls in Calabar, Nigeria. *Journal of Biological Sciences*, 15 (2): 98-101
- Hodasi, J. K. M. (1983). The potentials of snail farming in West Africa. *Paper Presented at the 8th Annual Conference of Nigerian Society for Animal Production (NSAP)*. Port-Harcourt, Nigeria. Pp.9-13.
- Ibom, L. A. (2009). Variation in reproductive and growth performance traits of white-skinned Black-skinned African giant snail hatchlings (*A. marginata* (Swainson)) in Obubra, Nigeria. *Ph.D. Thesis*. Department of Animal Science, University of Calabar, Calabar, Nigeria.
- Ibom, L. A., Okon, B. and Adinya, I. B. (2011). Evaluation of growth and body traits of snaillets Obtained from the cross breeding of black-skinned × white-skinned snails (*A. marginata* (S)) in the Niger Delta area of Nigeria. *African Journal of Agricultural Research*, 6(21):4968-4972.
- Ibom, L. A., Okon, B. and Essien, A. (2008). Morphometric analysis of eggs laid by two ectotypes of Snails, *Archachatina marginata* (Swainson) raised in captivity. *Global Journal of Agricultural Sciences*, 7(2):119-121
- Ibom, L. A., Okon, B., Okon, F. I. and Nwambam, I. U. (2014). Morphological characterization of giant African land snails from three agro-ecological zones of Delta State, Nigeria. *Proceedings of the 3rd International Conference/Workshop on Giant African Land Snail (NetGALS)*. Abeokuta, Nigeria. Pp. 38-44.
- Nisbet, R. N. (1974). The life of Achatinidae in London. *Proceedings of the Malacological Society of London*, 41:1171.
- Nworgu, F. C. and Fapohunda, J. B. (2002). Performance of broiler chicks fed mimosa (*Mimosa invisa*) Leaf meal supplements. *Proceedings of the 27th Annual Conference of Nigerian Society for Animal Production (NSAP)*. Akure, Nigeria. Pp. 128-131.
- Ojating, I. (2001). Rearing, food predilection and reproductive performance of African land snail, *Marginata*, in kitchen farm at Okuku and Umudike. *Journal of Sustainable Agriculture and Environment*, 3(1):135-140.

- Ojating, I. and Ogar, G. E. (2002). Effect of feed types on the growth performance of African giant snail Fattened at room temperature. *Journal of Sustainable Agriculture and Environment*, 4(2):303-308.
- Okon, B., Ibom, L. A., Etuk, N. E. and Akpan, E. W. (2008). Variations in growth patterns and Conformation of snails: Influence of strain and location on isometry of growth in Cross River State of Nigeria. *Journal of Agriculture, Forestry and the Social Sciences (JOAFSS)*, 6(2):218-227.
- Okon, B., Ibom, L. A., Ekpo, I. A. and Ewa, E. C. (2009). Evaluation of reproductive and some egg Quality parameters of Albino snails (*A. marginata saturalis*(Swainson)). *Journal of Applied Sciences (JAS)*, 12(1):8234-8241.
- Okon, B., Ibom, L. A., Williams, M. E. and Etukudo, O. M. (2010a). Effects of parity on breeding and Morphometric traits of eggs and hatchlings of F₁ crossbred of snails. *Journal of Agriculture, Biotechnology and Ecology*, 3(1):36-43.
- Okon, B., Ibom, L. A., Williams, M. E. and Akwa, N. T. (2010b). Parity effects on breeding and morpho-metric traits of eggs and hatchlings of purebred albino snails (*A. marginata*(Swainson)). *Journal of Agriculture, Biotechnology and Ecology*, 3(1):44-54.
- Okon, B. and Ibom, L. A. (2011). Phenotypic correlation and body weights prediction using morpho-Metric traits of snails in Calabar, Nigeria. *Ethiopian Journal of Environmental Studies and Management*, 4(3):7-11.
- Okon, B. and Ibom, L. A. (2012). *Snails Breeding and Snailery Management*. Calabar:Freshdew Publication. P.90.
- Okon, B., Ibom, L. A., Williams, M. E. and Ekong, N. B. (2012a). Effects of breed on reproductive Efficiency of two most popular snails (*A. marginata* (S) and *A. achatia*) in Nigeria. *Journal of Agricultural Science*, 4(8):236-245.
- Okon, B., Ibom, L. A., Ettah, H. E. and Udoh, U. H. (2012d). Comparative differentiation of morphometric traits and body weight prediction of giant African land snails with four whorls in Niger Delta region of Nigeria. *Journal of Agricultural Science*, 4(10):9756-9760.
- Olawoyin, O. O. and Ogogo, A. U. (2006). Prediction of optimum stocking density in growing African giant land snails. *Tropical Journal of Animal Science*, 9(2): 75-84.
- Omole, A. J. (1998). The utilization of different energy supplements on performance characteristics of grower edible giant land snails (*A. marginata*). M.Sc. Thesis, Department of Animal Science, University of Ibadan, Ibadan, Nigeria.
- Omole, A. J. K., Taiwo, A. A. and Amusan, J. A. (2007). *Guide Technical /Bulletin, Practical Snail Farming*. Ibadan, Nigeria. Institute of Agriculture Research and Training, Moor Plantation. P.26.

- Plummer, J. M. (1975). Observations on the reproductive, growth and longevity of laboratory colony of *Archachatina (Calachatina) marginata*. *Proceedings of the Malacological Society*. London. 141:395-413.
- Udedibe, A. B. I. (1989). The giant land snails and prospects of snails farming in Nigeria. *A paper presented at the 25th Annual Conference, Agricultural Society of Nigeria*. Owerri, Nigeria. P.14.
- Udeh, I. and Ogbu, C.C. (2011). Principal component analysis of body measurement in three strains of broiler chicken. *Animal Science*, 6(2): 1597-6343.