

Original Research Article

Survival, biomass production and growth response of seedlings to seed size variation of *Albizia lebbbeck* (L.) Benth.

Comment [M1]: This is also growth and can be deleted. Consider, otherwise title is greatly improved

ABSTRACT

Aims: To investigate the influence of seed size variation on seedling growth height, root collar diameter, root length, seedling length, root shoot ratio, number of leafs, leaf area, dry weight and survival of *Albizia lebbbeck* in the nursery.

Place and Duration of Study: The experiment was conducted in the nursery at Malawi College of Forestry and Wildlife between March and September 2014.

Methodology: Seeds were categorized into three groups in regard to their length and weight: large (≥ 0.80 cm) and (≥ 0.20 g); medium (0.51-0.79cm) and (0.11-0.19g); and small (≤ 0.50 cm) and (≤ 0.10 g). The treatments were completely randomized into four replications. A total of 600 seeds were used with each treatment having 50 seeds. To improve germination, seeds were mechanically nicked with secateurs.

Comment [M2]: Be sure that you have used the means with $\pm$ se throughout

Results: Seedling height, root collar diameter, root length, seedling length, root shoot ratio, number of leafs, leaf area, dry weight and percentage survival were significantly ($P < 0.001$) affected by seed size variation. Large seeds produced higher seedling height (58.4cm), root collar diameter (5.97mm), root length (29.7cm), seedling length (88.1cm), root shoot ratio (0.51), number of leafs (8.21), leaf area (405.0 cm² plant⁻¹), dry weight (11.3 g plant⁻¹) and percentage survival differences (89.6%) compared to seedling height (43.7cm), root collar diameter (4.03mm), root length (20.2cm), seedling length (63.8cm), root shoot ratio (0.46), number of leafs (5.93), leaf area (207.3 cm² plant⁻¹), dry weight (7.8 g plant⁻¹) and percentage survival differences (65.2%) of medium seeds and seedling height (29.3cm), root collar diameter (2.01mm), root length (12.6cm), seedling length (41.9cm), root shoot ratio (0.43), number of leafs (3.66), leaf area (136.8 cm² plant⁻¹), dry weight (5.1 g plant⁻¹) and percentage survival differences (46.1%) of small seeds. There was a strong positive relationship significant ($P = .05$, $r > .80$) relationship ($r > .80$) between seed

Comment [M3]: Revise this statement as suggested : There significant correlation between seed size and biomass production (P., r..), height (P., R..) etc

size and seedling growth, biomass production and percentage survival. This indicateThis indicates that seed size could be used as parameter for predicting seedling growth rates in the nursery.

Conclusion: The present study has demonstrated the merits of grading seeds and that use of large size seeds for production of high quality transplants of *Albizialebbeck* in the nursery is recommended.

Key words: shoot length; root collar diameter;root length;leaf area; number of leaf; dry weight;survival.

1. INTRODUCTION

Seed plays an important role in transfer of genetic characters and improvement of qualitative and quantitative traits of production. One of the most important aspects in maximizing crop yield is planting high quality seed [1]. Seed size is an important parameter of seed quality that influences the germination, growth and biomass of the nursery seedlings and that trend correlates to the future crop [2].

The influence of seed size on seedling emergence and growth in various tree species have been studied by many researchers [1, 2, 3 – 9], but there is no consistent trend between seed size and seedling growth and survival. For example, large seeds produced more vigorous seedlings than small seeds in *Mammeasuriga* and in *Afzeliaquanzensis* [2, 9], because of higher food reserves stored in large seeds [1]. On contrast, there were no significant differences in seedling growth of *Bauhinia thonningii* due to seed size [8].

*Albizialebbeck*L. Benth.is an important multipurpose tree species belonging to the family of *Leguminosae*[10]. The tree species is well distributed in all regions of Malawi and it has been prioritized as one of the species for conservation in Malawi to enhance its contribution to livelihood of communities. The morphological description, its uses and application in Malawi has been well explained by [11].Seeds sizes of *A. lebbeck* vary widely [12], however no attempt has been made to investigate the effect of seed size on seedling growth and survival of *A. lebbeck*. It is considered necessary to know the effect of seed size on seedling growth and survival so as to get healthy seedlings of plant species for a purpose of transportation in the field [13].To fill the knowledge gap, this study was conducted to investigate the influence of seed size variation on seedling growth, biomass productionand survival of *A. lebbeck*in the nursery.

59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88

2. MATERIAL AND METHODS

2.1 Seed Acquisition and Study site

Albizia lebbek seed from Salima (13°53'S, 34°26'E, and about 540m above sea level), was supplied by Forestry Research Institute of Malawi (FRIM) in March 2014. This study was conducted at Malawi College of Forestry and Wildlife (MCFW) nursery (14°19'S, 34°17'E, and 1591 m above sea level) which is located in Malawi near the tropical savannah region in Southern Africa. Mean annual rainfall at MCFW is receives 1200 mm to 1800 mm rainfall per annum, with annual temperature ranging from -7 °C to 25 °C. It is situated about 85 km south east of Lilongwe the capital.

2.2 Experimental design and treatments

Seeds were categorised into three treatments: large, medium and small according to both seed length and weight (Table1). The treatments were completely randomized in four replicates. Each treatment had 50 seeds making a total of 600 seeds that were used for the whole experiment. All experimental seeds were mechanically nicked with secateurs to improve germination as the technique has been effective in other studies [8, 11, 14].

Seed were sown on 31st March 2014 in 10 cm black polythene tubes filled with soil collected from a natural *Brachystegia* woodland stand as which contain adequate nutrients for seedling growth [9]. One seed was sown per tube at a depth of 2cm, as previously recommended[15]. All pots were watered regularly to maintain adequate moisture necessary for germination and seedling growth.

Table 1: Mean seed length and weight of large, medium and small seeds used in growth and survival experiments. Standard error of the mean (e) is shown in Seed size categories in terms of length and weight with standard errors in parenthesis

Seed size category	Range of length (cm)	Average length (cm)	Range of seed weight (g)	Average weight (g) of 100 seeds
Small	≤ 0.50	0.32 ± 0.05	≤ 0.10	0.09 ± 0.01
Medium	0.51 – 0.79	0.64 ± 0.04	0.11 – 0.19	0.13 ± 0.02
Large	≥ 0.80	0.98 ± 0.05	≥ 0.20	0.22 ± 0.02

Comment [M4]: You have a choice to use the range and the mean or mean alone. I think the range is not important statistically

Formatted: Highlight

Formatted: Highlight

Formatted: Font: Not Bold

Formatted: Highlight

Formatted: Highlight

2.3 Data collection and analysis

Seeds germination was recorded daily and the experiment concluded after 30 days when no more germination occurred. A seed was considered germinated when there was emerged by a visible protrusion of split seed coat with the cotyledons, hypocotyls, and epicotyl on the soil surface. Daily germination percentages were summed up to obtain cumulative germination for each treatment. After the completion of seed germination experiment, shoot height, root collar diameter and survival of seedlings were monitored for 150 days. All the seedlings were measured for total shoot height and root collar diameter once for every at 30 days interval and seedling survival recorded simultaneously. of the seedlings was also assessed at the same interval. Total shoot height was measured by using a 30 cm ruler and collar diameter by using a microcalliper to the nearest 0.01 mm. The measurements were taken just below the cotyledons. The number of seedlings that survived was also counted in each and every treatment unit. At the termination of the experiment root length, number of leaves, leaf area and dry weight of the seedlings were determined.

Comment [M5]: Is a repetition from line 96.

Germination percentage was calculated by dividing the total number of seeds that germinated in each treatment by the number of seeds sown (excluding dead seeds) and multiplied by 100. Survival percentage of seedlings that survived was also calculated based on the seeds that germinated. Data on seedling length and root shoot ratio were obtained by adding root length to shoot length and dividing root length by shoot length for each seedling respectively. Germination percentage, seedling height, seedling length, root shoot ration, root collar diameter, survival percentage, root length, number of leaves, leaf area and dry weight data were tested for normality and homogeneity with Kolmogorov-Smirnov D and normal probability plot tests using Statistical Analysis of Systems software version 9.1.3[16]. After the two criteria were met the data were subjected to analysis of variance (ANOVA) using the same Statistical Analysis of Systems software and means were separated with using Fischer's least significant difference (LSD) at the 0.05 level. Pearson's correlation was used to determine the relationship between seed size and the parameters studied.

Comment [M6]: Delete as dead seeds could not be determined before sowing. Assume all planted seeds were live and death/decay is just a natural occurrence that you could not control. Moreover, its random across seeds sizes. You are free to accept or reject the advice

Comment [M7]: In most studies, root :shoot ratio is for dry weights. In your case, you may consider specifying the fact in the table's caption.

Comment [M8]: See comment in the regression table results

119 3. RESULTS AND DISCUSSION

120 3.1 Germination

121 There was no significant ($P>.05$) differences in cumulative germination percentage among the three seed
122 size variation (Table 2). ~~The present results~~ Results of present study are consistent with the previous
123 studies [8, 9, 11]. This could mean that all the three seed size categories can germinate provided that high
124 physiological quality seeds are used [9]. According to [17] some of the desirable seed physiological
125 qualities include plumpness, high purity, freedom from pests and diseases, and being dried to a right
126 moisture content.

127

128 3.2 Seedling growth and biomass production

129 There were significant ($P<.001$) differences in seedling height, root collar diameter, root length, seedling
130 length, root shoot ratio, number of leaves, leaf area and dry weight among the three seed sizes studied
131 (Tables 2 and 3). Large seeds produced seedlings with higher height, root collar diameter, root length, root
132 shoot ratio, number of leaves, leaf area and dry weight than medium and small seeds, while medium seeds
133 produced seedlings with medium height, root collar diameter, root length, seedling length, root shoot ratio,
134 number of leaves, leaf area and dry weight than small seeds. The findings of the present study are in
135 agreement with previous studies [1, 2, 18, 19]. Production of superior seedlings by large seeds has been
136 attributed to growth enhancement by higher carbohydrates reserves in their endosperm when compared to
137 medium and small seeds [2, 20]. On the other hand, [8, 21, 22] reported no significant influence of seed size
138 on seedling growth in a few tropical tree species, while [2] reported that medium sized seeds produced
139 better seedling growth than those of smaller and larger ones. These differences may be due to the fact that
140 seedling growth is highly controlled by many intrinsic and extrinsic factors and is species specific
141 [23]. Although seedling growth is not always correlated to seed size [21, 22], the present study has shown
142 that large seeds should be used for production of superior seedling in terms of seedling vigour, in *A.*
143 *lebbeck*.

144

145

146

147

Comment [M9]: Conform to this style i.e.
 $P<0.0..$ instead of $P<.0..$ throughout the paper.

148

149 **Table 2** Effect of seed size variation on germination percentage, root length, shoot length, seedling length
150 and **root shoot ratio** with standard errors in parenthesis of *Albizia lebeck* at nursery

Seed size	Germination (%)	Root length (cm)	Shoot length (cm)	Seedling length (cm)	Root shoot ratio
Large	100.0(0.0) ^a	29.7(1.5) ^a	58.4(1.4) ^a	88.1(1.5) ^a	0.51(0.01) ^a
Medium	96.3(1.5) ^a	20.2(1.3) ^b	43.7(2.7) ^b	63.8(1.9) ^b	0.46(0.03) ^b
Small	95.1(1.6) ^a	12.6(1.3) ^c	29.3(1.3) ^c	41.9(1.3) ^c	0.43(0.03) ^c
CV%	5.2	4.6	7.3	8.9	6.3

Comment [M10]: I prefer use of ± instead of parenthesis. Unless it a journal requirement, consider changing as it reduces wordiness.

151 | Note: ^{a,b,c} means with different subscript within a column significantly differ ($P < 0.001$)

152

153 **Table 3** Effect of seed size variation on root collar diameter, number of leaves, leaf area, dry weight and
154 survival percentage with standard errors in parenthesis of *Albizia lebeck* at nursery

Seed size	Root collar diameter (mm)	Number of leaves	Leaf area (cm ² plant)	Dry weight (g plant)	Survival (%)
Large	5.97(0.01) ^a	8.21(0.04) ^a	405.0(2.4) ^a	11.3(0.6) ^a	89.6(2.5) ^a
Medium	4.03(0.02) ^b	5.93(0.03) ^b	207.3(3.2) ^b	7.8(0.3) ^b	65.2(2.3) ^b
Small	2.01(0.02) ^c	3.66(0.03) ^c	136.8(2.2) ^c	5.1(0.3) ^c	46.1(2.4) ^c
CV%	5.3	4.6	6.8	8.9	7.2


155 | Note: ^{a,b,c} means with different subscript within a column significantly differ ($P < 0.001$)

156 | Shoot length and root collar growth trend of *Albizia lebeck* for the period of 150 days are presented in
157 Figures 1 and 2 respectively. The shoot length and root collar growth of *Albizia lebeck* was significantly
158 ($P < 0.001$) higher in large seeds than in other seed size variations. Large seeds gave the highest seedling
159 height growth which increased rapidly between 30 and 120 days, then increased at the decreasing rate
160 between 120 and 150 days. A similar trend was also observed for the other seed size variations and for
161 the root collar diameter growth (Figure 2). According to [6], the higher growth rate of large seeds could be
162 due to the reason that large-seeded species are predicted to be better adapted to the catastrophic events
163 encountered by seedlings because they can compensate for damage using seed reserves. The growth
164 increased at an increasing rate between 60 and 120 days because the seedlings were using the
165 photosynthesized food for growth, as the food reserves were being invested in the tap root of the plants.
166 This was indicated by the plants changing of cotyledons colour from brown yellowish to right green in the
167 nursery.

Comment [M11]: Consider abbreviating throughout the paper after the first full mention of species, unless at the beginning of a sentence.

Formatted: Left


168



169 **Figure 1** Seedling height growth of *Albizia lebeck* from different seed size categories

170

171



172
173 **Figure 2** Seedling root collar growth of *Albizia lebeck* from different seed size categories

174

175

3.3 Seedling survival

A log survival percentage against time was constructed (Figure 3). Survivorship curves of seedlings from all the three seed size variation were close to Deevy Type II curve with more or less constant mortality rate. However, survivorship curves for medium and small seeds were steeper than that for large seeds. The results are in conformity with other researchers [24, 25]. Survival of seedlings is dependent on the amount of the reserves at the time of re-growth, such that those with greater amount of food reserves are able to regenerate new shoots [25].




Figure 3 Survival trend of *Albizia lebeck* seedlings from different seed size categories

3.4 Correlation among seed size, seedling growth, biomass production and survival

Correlation and regressions equations results among seed size, shoot length, root length, root collar diameter, number of leafs, leaf area, dry weight and percentage survival are presented in Table 4. The results shows that there was a strong positive significant ($P=0.05$) relationship ($r>0.80$) between seed size and shoot length, root length, root collar diameter, number of leafs, leaf area, dry weight and percentage survival. This implies that seed size could be used as parameter for predicting seedling growth rates in the nursery. The present results are in agreement with the results of [9] who reported a strong relationship between seed size and seedling growth height and root collar diameter growth of *Azadirachta indica*.

198 **Table 4** Regression equations (above diagonal) and correlations (below diagonal) among seed size, shoot
 199 length, root length, root collar diameter, number of leaves, leaf area, dry weight and percentage survival of
 200 *Albizia lebbeck*
 201

Parameter	Seed size	Shoot length (cm)	Root length (cm)	RCD (mm)	Number of leaves	Leaf area (cm ² plant)	Dry weight (g plant)	Survival (%)
Seed size	-	Y=198X+13	Y=118X+2.4	Y=28X-0.22	Y=32X+1.1	Y=1892X-52	Y=40X+1.4	Y=283X+20
Shoot length (cm)	r=0.92*	-	Y=0.59X-5.0	Y=0.13X-1.8	Y=0.15X-0.7	Y=9.37X-166	Y=0.22X-2.0	Y=1.55X-3.1
Root length (cm)	r=.93*	r=.99**	-	Y=0.23X-0.6	Y=0.26X+0.6	Y=16.1X-91	Y=0.38X-0.2	Y=2.64X+9.9
RCD (mm)	r=.95*	r=.99**	r=.99**	-	Y=1.15X+1.3	Y=67.7X-34	Y=1.57X+1.2	Y=11X+19.4
Number of leaves	r=.95*	r=.99**	r=.99**	r=.99**	-	Y=59.4X-114	Y=1.38X-0.6	Y=9.67X+6.4
Leaf area (cm ² plant)	r=.90*	r=.96*	r=.98*	r=.94*	r=.95*	-	Y=0.02X+2.1	Y=0.16X+26
Dry weight (g plant)	r=.81*	r=.96*	r=.97*	r=.92*	r=.93*	r=.97*	-	Y=6.93X+11
Survival (%)	r=.82*	r=.97*	r=.97*	r=.93*	r=.94*	r=.97*	r=.99**	-

Comment [M12]: Not sure if r in the table is from regression or Pearsons correlation analysis?? If from Pearsons, then we should have two tables as the methods are distinct and their results should be separated and adequately described in results section.

202 **Note:****Highly significant ($P<0.001$), *significant at ($P<0.05$), RCD: Root collar diameter
 203
 204

205 4. CONCLUSION

206 The current study has revealed that seed size has a clear effect on seedling growth, biomass production
 207 and survival of *A. lebbeck* in the nursery. Larger seeds produced high quality seedlings and that had a
 208 higher survival rates than medium and small seeds. This confirms merits of grading seed to enhance
 209 transplant quality. There was a strong positive significant relationship between seed size and seedling
 210 growth, biomass production and survival. This indicates that seed size could be used as parameter for
 211 predicting seedling growth rates in the nursery. Therefore, it is recommended to use large sized seeds for
 212 production of high quality transplants of *A. lebbeck* in the nursery.

214 REFERENCES

- 215 1. Zareian A, Hamidi A, Sadeghi H, Jazaeri MR. Effect of Seed Size on Some Germination
 216 Characteristics, Seedling Emergence Percentage and Yield of Three Wheat (*Triticum aestivum* L.)
 217 Cultivars in Laboratory and Field. Middle-East Journal of Scientific Research. 2013; 13(8):1126 –
 218 1131.
- 219 2. Gunaga RP, Doddabasava D, Vasudeva R. Influence of seed size on germination and seedling
 220 growth in *Mammea suriga*. Karnataka Journal of Agriculture and Science. 2011; 24(3):415 – 416.

3. Dar FA, Gera M, Gera N. Effect of seed grading on germination pattern of some multi-purpose tree species of Jammu region. *Indian Forester*. 2002; 128:509 – 512.
4. Larsen SU, Andreassen C. Light and heavy seeds differ in germination percentage and mean germination thermal time. *Crop Science*. 2004; 44:1710 – 1720.
5. Willenborg CJ, Wildeman JC, Miller AK, Rossmann BG, Shirliffe SJ. Oat Germination Characteristics Differ among Genotypes, Seed Sizes and Osmotic Potentials. *Crop Science*. 2005; 45:2023 – 2029.
6. Baraloto C, Forget PM. Seed size, seedling morphology, and response to deep shade and damage in Neotropical rain forest trees. *The American Journal of Botany*. 2007; 94(6):901–911.
7. Kaydan D, Yamur M. Germination, seedling growth and relative water content of shoot in different seed sizes of triticale under osmotic stress of water and NaCl. *African Journal of Biotechnology*. 2008; 7:2862 – 2868.
8. Mwase WF, Mvula T. Effect of seed size and pre-treatment methods of *Bauhinia thonningii* Schum. on germination and seedling growth. *African Journal of Biotechnology*. 2011; 10(13): 5143 – 5148.
9. Mtambalika K, Munthali C, Gondwe D, Missanjo E. Effect of Seed Size of *Azelaquanzensis* on Germination and Seedling Growth. *International Journal of Forestry Research*. 2014; doi.org/10.1155/2014/384565.
10. Bhat GS, Chauhan PS. Provenance variation in seed and seedling traits of *Albizia lebbek* Benth. *Journal of Tree Science*. 2002; 21:52–57.
11. Missanjo E, Maya C, Kapira D, Banda H, Kamanga-Thole G. Effect of Seed Size and Pre-treatment Methods on Germination of *Albizia lebbek*. *ISRN Botany*. 2013; doi:10.1155/2013/969026.
12. Uddin MB, Mukul SA, Khan MASA, Hossain MK. Effects of phosphorous fertilizer on seedlings growth and nodulation capabilities of some popular agroforestry tree species of Bangladesh. *Journal of Forestry Research*. 2007; 18: 283 – 286.
13. Ahrwar JR. Effect of seed size and weight on seed germination of *Alangium lamarckii*, Akola, India. *Research Journal of Recent Sciences*. 2012; 1(ISC-2011): 320 – 322.
14. Likoswe MG, Njoloma JP, Mwase WF, Chilima CZ. Effect of seed collection times and pre-treatment methods on germination of *Terminalia sericea* Burch. *African Journal of Biotechnology*. 2008; 7(16):2840 – 2846.

15. Abideen MZ, Gopikumar K, Jamaludheen V. Effect of Seed Character and its Nutrient content on vigour of Seedlings in *Pongamiapinnata* and *Tamarindasindica*. My Forest Journal. 1993; 29:225 – 230.
16. SAS9.1.3. Qualification Tools User's Guide. SAS Institute Inc., Cary, NC, USA, 2004.
17. Schimdt L. Dormancy and Pre-Treatment: Guide to Handling of Tropical and Subtropical Forest Seeds. Danida Forest Seed Center, Denmark. 2000.
18. Manonmani V, Vanangamudi K, Rai RSV. Effect of seed size on seed germination and vigour in *Pongamiapinnata*. Journal of Tropical Forestry and Science. 1996; 9:1 – 5.
19. Ebofin AO, Agboola DA, Ayodele M, Aduradola AM. Effect of seed sizes and seedling growth of some Savanna Tree legumes. ASSET Journal. 2003; 3(2):109–113.
20. Hewitt N. Seed-size and shade tolerance: a comparative analysis of North American temperate trees. Oecologia. 1998; 114:432 – 440.
21. Agboola DA. The effect of seed size on germination and seedling growth of three tropical tree species. Journal of Tropical Forestry and Science. 1996; 9:44 – 51.
22. Malavasi M, Malavasi UC. Effect of seed size on seedling growth of a shade-tolerant tropical tree (*Hymaneastilbocarpa* Haynes). Tree Planting Notices. 1996; 46:130 – 133.
23. Murali KS. Patterns of seed size, germination and seed viability of tropical tree species in Southern India. Biological Tropical. 1997; 29:271 – 279.
24. Khurana E, Singh JS. Germination and seedling growth of five tree species from tropical dry forest in relation to water stress: impact of seed size. Journal of Tropical Ecology. 2004; 20:385 – 396.
25. Mwitwa JP, Munthali CRY, van Wyk G. Half-sib family Variations in shoot and root Traits of Seedlings of *Pterocarpus angolensis* (family: Fabaceae, syn. papilionaceae). Southern Hemisphere Forestry Journal. 2007; 69:91 – 94.