

Concentrations of Cadmium, Copper, Lead and Zinc in Soils and Vegetable Organs from Periurban Agriculture Areas of Abidjan in Cote d'Ivoire

ABSTRACT

Production quality in periurban agriculture is question mark regarding to soil potential contamination affecting yields. The level of contaminations of soils and vegetables by copper (Cu), zinc (Cu), cadmium (Cd) and lead (Pb) around Abidjan city were assessed. Survey was conducted in 2013 within cultivated areas of sweet potato and hibiscus locally named "Dah" as encountered in three locations of Abidjan district (Port-Bouët, Yopougon and Bingerville) according to the intensities of industrial and commercial activities of which, Bingerville was the control site with lowest activities. Soil (0 – 20 cm) samples associated to that of plants (leaf, stem and root) were taken randomly for laboratory analysis. Toxic levels ($> 8 \text{ mgkg}^{-1}$) of Pb were significantly ($p < .0001$) determined in plant organs from Port-Bouët site indifferently to crops while, lower soil content of Pb (35.5 mgkg^{-1}) than that of Yopougon (39.8 mgkg^{-1}) was observed, however. Except the synergisms observed between leave concentration of Pb and soil contents of Cd, Cu and Zn, non of soil parameters were relevant for this while, the proximity of inland waters was suspected. The partitioning of Pb in plant organs pointed out phytoremediation potential of hibiscus with lowest risk of toxicity ($2.92 - 9.72 \text{ mgkg}^{-1}$) in edible leave against an average of 8.08 mgPbkg^{-1} in the tuber of sweet potato. For strengthening consistence of knowledge, studies of Pb and Zn interaction as well as Pb translocation in tuber plants of tropical ecosystems were suggested.

Key words: *Metallic Trace Elements, vegetable, lead, interaction, phytoremediation, soil and plant contamination*

1. INTRODUCTION

In the humid forest zone of West Africa, the urban population is most important than that of rural zone (1) characterized by poverty and lack of job opportunities: about 2/3 of the population will be urban in the foreseeable future while agriculture practice remains most important in rural zones. This trend is of challenge for food security and supplying (2). In turn, informal economies can emerge from this exodus (3) including periurban agriculture. In fact, precarious lifestyle of a part of urban population imposes upon him reconnection with agriculture around the cities for subsistence and household income (4). This agriculture is supplying about 30% of world food (5; 6). However, the concerning agricultural production is more prone to pollution risk from urbanization (7) including metallic trace element accumulations in soils and crop yields. Many studies reported the harmfulness of such accumulation for humans and animals (8 - 14) as well as for plants in certain extend (15). Nowadays, there is increasing of urbanization coupled with periurban agriculture adoption in many african countries including Côte d'Ivoire. In Abidjan areas (economic capital of Côte d'Ivoire), leafy vegetable production and local market supplying of fresh products are increasing (16). Well, this agriculture is practiced in free spaces around manufactures sometime and somewhere with household refuse stocks which are auspicious to pollution (16 - 19) while, yet, this pollution effect is unknown in local crops of mass-market products as vegetable, especially, as sweet potato and hibiscus (Dah). Current study is volunteer to identify the level of pollution in soils and plants as induced by cadmium (Cd), copper (Cu), lead (Pb) and zinc (Zn) as pollutants in periurban agro-ecosystems (20; 21), especially for mass-market products of vegetables including hibiscus and sweet potato characterized by edible leaves locally. The aims were, i) to determine these metal concentrations in soils and plants in relation with other soil characteristics, ii) identify the most prone to contamination risk among the studied vegetable species as local mass market products, and iii) underline toxicity risk and phytoremediation potential of studied crop according to international standard values of Pb. Roughly, the study will point out soil characteristics most relevant to contamination of studied vegetable species for promotion of periurban agriculture carefully to consumer health.

2. MATERIAL AND METHODS

2.1 Description of studied sites

Abidjan is the economic capital of Côte d'Ivoire (West Africa), on the shoreline of Guinea Golf within latitudes 5°00 - 5°30 N and longitudes 3°50 - 4°10 W. It accounts for 10 districts and the major industrial activities of the country (22). It is characterized by subequatorial climate with annual average rainfall amount fluctuating between 1637 mm and 2048 mm irregularly distributed in time and space scales as bimodal rainfall in pattern (Two rainy seasons alternating with two dry seasons). Annual averages air temperature and hygroscopic measurement are recorded between 24°C - 30°C and 75% - 88% respectively according to Société d'Exploitation et de Développement Aéroportuaire, Aéronautique et Météorologique (SODEXAM). Open forest characterized the southern coastal zone neighboring a rainforest toward the northern zone. They are all degraded consecutive to anthropic activities including industrialization. The geology of the studied area belongs to the coastal sedimentary basin of Côte d'Ivoire, named "Continental terminal" and restricted to 2.5% of the country area (22). The soil is sand clayed Ferralsols somewhere Acrisols developed on tertiary and quaternary sand deposits. Actual survey was conducted in vegetable cropping areas of Port-Bouët (5°25N - 3°94W) and Yopougon (5°35N - 4°04 W) characterized by higher industrial and commercial activities contrasting with Bingerville (5°31 N - 3°87 W) as control site with lower activities (Fig.1).

2.2 Plants studied

Two vegetable species characterized by edible leaf as local dietary habits (16) were concerned: Hibiscus (*Hibiscus sabdaroufa*) locally named "Dah" and sweet potato (*Ipomoea batatas*). While hibiscus is matured about 2 -3 months of cropping duration sweet potato does so in a longer period of 3 - 6 months depending to the cultivars and the roots are also edible as tubers. Both are characterized by shallow rhizosphere (0 - 30 cm) receiving manual daily irrigation using perched ground water of well (2 - 3 m in depth).



Fig.1. Localization of studied sites (red marks)

2.3 Soil and plant sampling

Multi-sites (Bingerville, Yopougon and Port-Bouët) survey was conducted in 2013 in the district of Abidjan (5°18' N; 4°00' W; 10 m asl): The localities of Port-Bouët and Yopougon are characterized by higher intensity of industrial and commercial activities while prevailing agricultural activity accounts for Bingerville.

In 600 m² of vegetables cultivated areas in each of these locations, 12 soil composite samples were randomly taken in 0 – 20 cm using hand augur (5 subsamples for about 50 m²). Soil sampling was coupled with that of plant selecting one plant between 4 positions of soil sampling (50 m²). About 5 g sample of matured plant organs as fresh leaf, stem and root were taken from hibiscus (*Dah*) and sweet potato respectively. Soil and plant samples were kept in plastic package and transported in icebox for laboratory analysis.

2.4 Laboratory analysis

Soil samples were dried in room condition before being ground and sieved (2mm). Soil particle size and pH were determined using the methods of Robinson pipette (23) and glass electrode in 1/2.5 ratio of soil/solution (water or KCl) respectively. Soil electric conductivity (EC) measurement was coupled with that of soil pH. The soil contents of organic-C and total-N were determined as described by Walkley and Black (24) and Bremner (25) respectively and a factor of 1.72 was applied to the content of organic-C to determine that of soil organic matter. Furthermore, 1N H₄OAc (pH 7.0) (extracting) and Atomic Absorption Spectrometry (apparatus) were used for soil Cation Exchange Capacity (CEC) determination.

Soil chemical extraction of metal (Cd, Cu, Pb and Zn) was also done using a ground soil sample (0.3 g) of 150 µm in size and aqua regia [3 mL de HNO₃ (65%; v/v) + 1 mL de HCl (37%; v/v)] as described by (26; 27). Three measurements (Atomic Absorption Spectrometry) were done for each analysis and the average was reported and compared to the standard values (Table 1) of Alloway (28).

Plant material was washed in tap water to remove adhered soil particles and subsequently shredded, oven dried (60°C), ground (1mm) for the use of 0.5 g of sample. Plant samples (leaf, stem and root) were separately digested in 6 ml of each of H₂O₂ and HNO₃ during 3 hours at constant temperature of 95°C in a Digi PREP and repeated for hibiscus and sweet potato respectively. Flame atomic absorption spectrometry was used for the determination of plant concentrations of Cd, Cu, Pb and Zn.

Table 1. Standard ranges (mgkg⁻¹) of soil content of metallic trace elements (Alloway (28))

Metals	Moderate	High	Average worldwide
Cu	20-62	65-160	20-30
Zn	100-250	250-426	50
Cd	0.2-1.0	0.70-2.00	2.0-46.3
Pb	10-30	60-90	100-180

2.5 Statistical analysis

By descriptive analysis, mean values of different studied metal contents in soil and their ratios were determined for a given site. Analyze of variance was also done to determine the mean values of Cd, Cu, Pb and Zn concentrations in a specific plant (hibiscus and sweet potato) organs (leaf, stem and root) according to studied sites (Bingerville, Port-Bouët and Yopougon). Moreover, Pearson correlation analysis were performed between soil (0 – 20 cm) contents of Cd, Cu, Pb, Zn and their concentrations in the leave of *Dah* (hibiscus) and sweet potato respectively. SAS (version 8) was used for statistical analysis in the confidence intervals ranging from 95% (ANOVA) to 90% (correlation).

3. RESULTS

3.1 Soil characteristics

Table 2 shows the values of physicochemical properties of soil for different studied sites respectively. According to the pH values, the soil is acidic at Bingerville (pH = 5.6) and neutral (pH = 7) at Port-Bouët, contrasting with the alkalinity observed for the soil of Yopougon (pH= 8.7). Major difference is observed in soil ΔpH which is 3 – 4 times higher at Yopougon than that of the other sites. The values of soil CEC are ranging between 1.4 and 11.8 cmolkg⁻¹ wherever, hence characterizing extensive low-clay activity of soils. In spite of the contrast in soil acidity, there is low values of EC at Bingerville (0.08 mScm⁻¹) and Yopougon (0.45 mScm⁻¹) where the soils are richer in fine particles (clay + silt) coupled with outstanding amounts of organic matter contents. In turn, EC value is relatively higher in the soil of Port-Bouët (0.65 mScm⁻¹) ranging in acceptable level for the normal growth of most of the plants. The ratios of soil contents of Cd, Cu, Pb and Zn as measured in 0 – 20 cm are presented for each of the studied sites (Bingerville, Port-Bouët and Yopougon) in the Fig 2. Port-Bouët site is characterized by the highest proportions of Cd (44%), Cu (49%) and Zn (75%) measurements while, 36% of Pb is noticed in there next to that (49%) of Yopougon site. Lowest values of measurement proportions (< 30%) are for Bingerville as the control site with 26% and 9% accounting for Cu and Zn respectively: soil contents of Cd (0.5 mg kg⁻¹), Cu (18.5 mgkg⁻¹) and Zn (35.6 mgkg⁻¹) are low in the topsoil at Bingerville while that of Pb (21 mgkg⁻¹) is moderate (Fig.3) according to international standard values (Table 1). In turn, highest values of soil contents of Cd (0.70 – 2 mgkg⁻¹) and Pb (60 – 90 mgkg⁻¹) are noticed at Port-Bouët and Yopougon respectively. Soil content of Zn is also high in Port-Bouët while that of Cu is low (< 20 mgkg⁻¹) or moderate (20 – 62 mgkg⁻¹).


3.2 Plant analysis

Mean values of Cd, Cu, Pb and Zn concentrations in plant organs of sweet potato according to studied sites are presented in Table 3. Roughly, there is significant high grand mean concentrations of Cd (0.72 mgkg⁻¹) and Zn (57.93 mgkg⁻¹) in the root as observed for Cu (25.8 mgkg⁻¹) in leaf. In turn, highest concentration of Pb accounts for root (9.60 mgkg⁻¹) and stem (10.50 mgkg⁻¹). This trend of partitioning is confirmed for Zn in each of the studied sites while, highest concentration of Cd is significantly determined in the leaf at Port-Bouët also outstanding with highest concentration of Cu in the root contrasting with the corresponding grand mean value for the three sites. Lowest concentrations of Cd, Cu and Pb are observed in the leaf collected at Yopougon as observed for Cu at Port-Bouët.


150 **Table 2:** Physicochemical characteristics in topsoil (0 - 20 cm) at Port-Bouët, Bingerville and
 151 Yopougon sites

Soil characteristics	Sites		
	Bingerville	Port-Bouët	Yopougon
pH_{water}	5.6 $\pm$ 0.50	7 $\pm$ 0.75	8.7 $\pm$ 0.55
pH_{KCl}	4.6 $\pm$ 0.76	6.8 $\pm$ 0.98	8.4 $\pm$ 0.53
ΔpH	1 $\pm$ 0.05	0.2 $\pm$ 0.02	0.3 $\pm$ 0.04
MO (gkg ⁻¹)	19.26 $\pm$ 1.35	21.19 $\pm$ 1.42	13.21 $\pm$ 2.34
C (gkg ⁻¹)	11.2 $\pm$ 1.05	12.32 $\pm$ 1.56	7.68 $\pm$ 2.25
N (gkg ⁻¹)	1.1 $\pm$ 0.23	1.8 $\pm$ 0.36	0.8 $\pm$ 0.16
CEC (cmol kg ⁻¹)	1.47 $\pm$ 0.88	1.53 $\pm$ 0.56	11.8 $\pm$ 0.35
Electric conductivity (EC) (mScm ⁻¹)	0.08 $\pm$ 0.01	0.65 $\pm$ 0.06	0.45 $\pm$ 0.03
Sand (%)	81 $\pm$ 9.32	84 $\pm$ 7.75	50 $\pm$ 6.26
Clay (%)	4 $\pm$ 1.45	1.50 $\pm$ 0.53	11 $\pm$ 2.02
Silt (%)	15 $\pm$ 1.56	14.50 $\pm$ 1.32	39 $\pm$ 1.75

152 $\pm$ is referring to standard deviation



153 **Fig.2.** Proportion (%) of metal contents in topsoil (0 – 20 cm) as determined in vegetable growing
 154 areas at Bingerville, Port-Bouët and Yopougon.
 155



156

157

158

Fig.3: Mean values of soil contents of Cu, Zn, Pb and Cd in topsoil (0 – 20 cm) according the three sites studied (Moderate levels are pointed respectively)

159 **Table 3:** Concentrations of Zn, Cu, Cd and Pb in leaf, stem and root of sweet potato at Bingerville, Port-Bouët and Yopougon sites

	Concentration of Zn (mgkg ⁻¹)				Concentration of Cu (mgkg ⁻¹)			
	Bingerville	Port-Bouet	Yopougon	Average	Bingerville	Port-Bouet	Yopougon	Average
Leaf	52.63 bA	40.74 cC	42.33 cB	45.24c±6.81	25.80 aA	21.73 bB	12.60 aC	20.03a±1.11
Stem	52.19 bA	41.21 bB	50.91 bC	48.11b±7.53	20.58 cB	21.67 bA	7.84 bC	16.68c±6.57
Root	57.93 aA	43.74 aC	53.01 aB	51.56a±8.48	21.31 bB	24.08 aA	6.7 cC	17.36b±7.90
P>F	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<0.001
	Concentration of Cd (mgkg ⁻¹)				Concentration of Pb (mgkg ⁻¹)			
	Bingerville	Port-Bouet	Yopougon	Average	Bingerville	Port-Bouet	Yopougon	Average
Leaf	0.52 cB	0.81 aA	0.57 bC	0.63b±0.14	6.17 cB	9.72 bA	2.92 bC	6.27b±3.23
Stem	0.64 bB	0.72 bA	0.41 cC	0.59b±0.15	6.41 bB	10.50 aA	5.01 aC	7.31a±2.99
Root	0.72 aB	0.65 cC	0.94 aA	0.77a±0.13	7.80 aB	9.60 bA	5.81 aC	7.73a±2.45
P>F	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<0.001

160 a, b and c are indicating mean values with significant difference in column for α=0.05; A, B and C are indicating mean values with significant difference in line
 161 for α=0.05 (± is referring to standard deviation)

162

163

164

165

166

167

168 **Table 4:** Concentrations of Zn, Cu, Cd and Pb in leaf, steam and root of *Dah* (hibiscus) at Bingerville, Port-Bouët and Yopougon sites

Concentration of Zn (mgkg ⁻¹)					Concentration of Cu (mgkg ⁻¹)			
	Bingerville	Port-Bouët	Yopougon	Average	Bingerville	Port-Bouët	Yopougon	Average
Leaf	50.39bA	42.31bB	41.72bC	44.81b±4.65	22.44bA	18.70 bB	12.93aC	18.02a±4.37
Stem	50.52bA	47.3aC	52.52aB	50.13a±2.80	20.60bA	19.62bB	9.3bC	16.51b±5.34
Root	58.32aA	42.46bC	51.06aB	50.62a±8.38	23.40aA	22.71aB	7.23bC	17.78a±7.74
P>F	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001

Concentration of Cd (mgkg ⁻¹)				Concentration of Pb (mgkg ⁻¹)				
Leaf	0.46cC	0.64aA	0.48bB	0.48b±0.14	6.91bB	8.81bA	3.10cC	6.28c±2.51
Stem	0.53bB	0.60aA	0.47bC	0.47b±0.09	6.17bB	9.42aA	5.52bC	7.04b±2.17
Root	0.61aC	0.66aB	0.77aA	0.77a±0.11	8.42aB	9.71aA	6.12aC	8.08a±1.93
P>F	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001	P<.001

169 a, b and c are indicating mean values with significant difference in column for α=0.05; A, B and C are indicating mean values with significant difference in line
170 for α=0.05 (± is referring to standard deviation)

Similar results are noticed for the grand mean of Cd, Cu, Pb and Zn concentrations in the different organs of hibiscus (Table 4): Highest concentrations are determined in the root. However, highest significant concentration of Zn is similarly observed in the stem as previously observed for Pb concentration in both root and stem of sweet potato. Nevertheless, there is variance of these results at Port-Bouët and Yopougon sites characterized by highest concentrations of Zn (47.3 mgkg^{-1}) and Cu (12.93 mgkg^{-1}) in the stem and leaf respectively. Furthermore, highest concentrations of Zn (50.39 mgkg^{-1}) and Cu (22.4 mgkg^{-1}) are determined in the leave collected at Bingerville while, highest concentrations of Cd (0.64 mgkg^{-1}) and Pb (8.81 mgkg^{-1}) account for the leave sampled at Port-Bouët. The concentrations of studied metals (Cd, Cu, Pb and Zn) are low or moderate in the leaf of Dah (hibiscus) at Yopougon area.

3.3 Soil and plant relationship

Table 5 shows Pearson correlation coefficients (R) as observed between soil contents of Cd, Cu, Pb and Zn and their corresponding concentrations in the leave of studied plants (sweet potato and hibiscus) indifferently to the sites. Significant correlation ($R = 0.75$) is limited soil contents of Cu and Zn among soil chemical parameters. But, there is significant and positive correlations between soil content of Cu and the concentrations of Pb in the leave of sweet potato and Dah (hibiscus) in similar range of 0.61 – 0.63 respectively. However, there are different relations between soil content of Cu and leaf concentration in Cd according to crops: In fact, significant and positive R (0.75) value is observed for sweet potato while no significant correlation accounts for hibiscus. Significant and positive correlations are observed for the concentrations of Pb (0.76) and Cd (0.91) in the leave of sweet potato regarding to soil content of Zn while, only occurring similarly for Pb (0.71) in the leaf of hibiscus as the most pollutant influenced by studied soil chemical elements. Moreover, there is no significant correlation for soil contents of Cd and Pb regarding to studied metal concentrations in both soil and plant leave indifferently to the sites.

Table 5 : Pearson correlation coefficients (R) between soil contents of Cd, Cu, Pb and Zn and corresponding concentrations in leaf, stem and root indifferently to studied sites

Coefficient of correlation (R) for sweet potato				
	Cu (soil)	Zn (soil)	Pb (soil)	Cd (soil)
Cu (soil)	1.00			
Zn (soil)	0.75***	1.00		
Pb (soil)	-0.01	0.01	1.00	
Cd (soil)	0.07	0.05	0.27	1.00
Cu (leaf)	0.16	0.19	-0.47	-0.09
Zn (leaf)	-0.41	-0.50	-0.35	-0.19
Pb (leaf)	0.63***	0.76***	-0.17	0.06
Cd (leaf)	0.77***	0.91***	0.15	0.16
Coefficient of correlation (R) for Dah (hibiscus)				

Cu (soil)	1.00			
Zn (soil)	0.75***	1.00		
Pb (soil)	-0.01	0.01	1.00	
Cd (soil)	0.07	0.05	0.27	1.00
Cu (leaf)	0.07	0.09	-0.48	-0.03
Zn (leaf)	-0.36	-0.41	-0.39	-0.25
Pb (leaf)	0.61***	0.71***	-0.29	-0.11
Cd (leaf)	0.48	0.58	0.10	-0.06

*** significant for $P < .001$; () : origin

4. DISCUSSION

4.1: Soil potential as source of pollutant

All the studied soils are derived from sedimentary material named locally “*continental terminal*” (30) and their difference could be accounted in serial variances of this sedimentation: Yopougon site in Northern of lagoon fault line was differing to the other sites by ferruginous sand stone as bed rock with the most fine particle size (clay + silt) of soil. However, difference in serial sedimentations should not be able to induce significant difference in soil content of metallic trace elements regarding to the extending homogeneous geologic and climatic conditions as basic characteristics of their native sources (31). In contrast with this homogeneity, soil contents of Cd and Pb were higher for Yopougon and Port-Bouët sites compared to that of Bingerville and were likely contaminated according to the standard values of Alloway (28). Hence, there was clearness of soil pollution as anthropic consequence (e.g. agriculture and industry) against the native homogeneity. High contribution of surface water dynamic as runoff and deep infiltration of lagoon water affecting ground water quality as early reported by Coulibaly *et al.* (32) and Soro *et al.* (33) in the studied area could be relevant for the pollution process since, this ground water was daily used for crop irrigation.

Thus, soil polluted potential as observed for Cd and Pb was likely relative to the distance from the local lagoon (*Lagune Ebrié*), as a driving factor early stigmatized as by Koné *et al.* (34) in the pollution hazard of coastal environment of Côte d'Ivoire; Bingerville site was most distant from the lagoon and less impacted than the other two sites.

Yet, of differences in studied soil acidities and soil contents of organic matter as well as the proportion of fine particle sizes in the soil, non accounted for obviousness in effective crop pollution by Cd, Cu, Pb and Zn contrasting with the arguments of Schwartz *et al.* (35) relating to highest mobility of these metals from a soil into plant with moderate acidity (pH ~6.5) and high content of organic matter. This contrast was further illustrated by the case of lowest soil content of Zn ($35.6 \text{ mgkg}^{-1} = 9\%$ of measurements) at Bingerville characterized by highest concentrations ($> 50 \text{ mgkg}^{-1}$) of Zn indifferently to plant organs (root, stem and leaf) and crops (sweet potato and hibiscus). Moreover, highest concentrations of Pb in plant organs were observed at Port-Bouët with 36% of measurements (33.6 mgkg^{-1} in soil) against 42% (39.8 mgkg^{-1} in soil) at Yopougon. The consideration of metal speciation which is

moreover widely variable (36), and their interactions (37) could have contributed to this: Bio-availability of organic-Pb²⁺ and Pb⁺ is depending to ecological conditions and some forms (polysaccharide) of Cd are not extractable. As consequence, soil total contents of these metals have limited accuracy in presuming soil potential of plant contamination. In turn, it is reported synergistic relation between Cd and Pb (not significant in current study) beside the effects of Cu/Cd and Cd/Zn ratios controlling the availability of soil Cd for plant uptake (38). Consequently, synergism can occurred between Cu and Pb as documented in Table 5 of current study. In contrast, antagonism was suspected between Cd and Zn (39) while highest concentrations of Cd and Pb were determined in the leave and stem collected at Port-Bouët in spite of noticeable amounts of Cu (35 mgkg⁻¹) and Zn (320.7 mgkg⁻¹) in the soil. With that in mind, positive correlations were observed between soil content of Zn and the concentrations of Cd and Pb in the leaf and steam of sweet potato while the concentration of Cd was not concerned as much for hibiscus. Hence, there is insight of minimizing atmospheric (including traffic) deposit of Pb effect in the studied environment while this process was accounting for 95% of Pb-source in plant pollution elsewhere (40) and ecological variance of Pb-Zn interaction is pointed out as further challenge of new coming study.

In the light of this analysis, plant pollution by Pb is most likely concerned in the periurban agro-ecosystem of Abidjan characterized by higher concentration (> 8 mgkg⁻¹) in plant tissues than the standard values defined by Godin (41). The conditions observed at Port-Bouët (neutral pH, sandy, high contents of organic matter and Zn coupled with moderate content of Cu and high EC) could be concerned with bio-availability of Pb, inducing plant contamination around the lagoon affected by wastewater as observed by Hassan et al. (42) elsewhere.

4.2: Phytoremediation of lead polluted soil

In spite of noticeable soil content of Pb (21 mgkg⁻¹) at Bingerville, lowest synergism with the other metals (Cd, Cu and Zn) observed in low range of soil content respectively can be suspected as condition of riskless observed for plant pollution in Pb. Similarly, the low soil contents of Cu (18.2 mgkg⁻¹) and Zn (69.6 mgkg⁻¹) at Yopougon could have inhibited bio-availability of Pb (39.8 mgkg⁻¹) which was characterized by highest soil content of Pb however. Coupling these contrasts with plant contamination as observed at Port-Bouët allows the proscription of Zn and Cu fertilizer applications in agro-ecosystem prone to Pb pollution in periurban agriculture. This finding reinforces the knowledge of fertilizer practice as previously limited to Ca, P and S potentially reducing the bio-availability of Pb (43; 39). However, there is no improving of soil health and quality by these practices beside its high cost while, the soil still remained harmfulness for microbial activity which is required in sustaining agricultural ecosystem services (44; 45). Therefore, polluted soil phytoremediation is a required efficient strategy in restoring soil health: soil metal bioextraction (bioharvesting) is estimated by high accumulation in growing plant (45), depleting soil content of concerning pollutant metal.

Lead (Pb) concentrations in vegetal organs of sweet potato were roughly lower in leaf than the stem and root. But at Port-Bouët site where Pb concentration was over the critical toxicity level (> 8 mgkg⁻¹), highest concentration (10.5 mgkg⁻¹) accounted for the stem contrasting with the argument of low translocation of Pb in plants. Thereby, an average yield of 10 kgha⁻¹ will export about 105 mg Pb ha⁻¹ from soil according to soil particle size density in Africa (46). This amount of Pb is a noticeable removal as compared with the normal level of 100 mg Pb kg⁻¹ in soil (41). This aptitude can be considered as a potential of soil remediation so far, only the root and leave of sweet potato are edible.

280 This property is better with hibiscus (*Dah*) characterized by non edible root and steam
 281 accounting for about 20 mgkg⁻¹ for Pb concentration corresponding to twice level determined
 282 for the steam of sweet potato. Moreover, the concentration of Pb in edible leaf of hibiscus did
 283 not exceed the critical level of 8 mg Pb kg⁻¹ at Port-Bouët as the most polluted site and a low
 284 grand mean value of 6.3 mg kg⁻¹ observed in there was riskless for human consumption.
 285 Hence, hibiscus can be recommended in periurban agriculture in area prone to industrial and
 286 commercial pollution as Abidjan city for consumption and Pb-polluted soil remediation in
 287 some extend when compared with sweet potato. This tuber crop (sweet potato) has highest
 288 concentration of Pb in both root and steam at Port-Bouët and tuber consumption will be risky
 289 for human health (47) meanwhile, tuber crops are supposed to have low potential of Pb
 290 accumulation comparing with leafy vegetables (46). Genotypes and ecological differences
 291 could have contributed to this contrast that should be explored with variance of scenarios in
 292 tropical ecosystems by further study.

293

294 5. CONCLUSION

295 The practice of periurban agriculture is of effective risk of Pb toxicity around Abidjan district,
 296 especially for tuber crop as sweet potato, and the proximity of surface water is likely more
 297 contributing to this compared with soil properties in extended similar geologic and climatic
 298 conditions. However, moderate soil contents of Cu and Zn are required at least for high Pb
 299 accumulation in plant tissues independently to soil total content of metal which may have
 300 effective effects relevant to the speciations. Cropping of vegetable as hibiscus (*Dah*) with
 301 edible leaf, was recommended for remediation of Pb-polluted soil because of high
 302 accumulation of Pb in both root and steam.

303 However, pending matters are underlined by the study in relation with genotype and
 304 ecological effects in plant nutrition of Pb, especially, the interaction with Zn and that of Pb
 305 accumulation in tuber crop are concerned.

306

307 REFERENCES

- 308 1. Drechsel, P., Quansah, C., Penning de Vries, F., 1999. Urban and periurban
 309 agriculture in West Africa: Characteristics, Challenges and need for action. Smith,
 310 O.B. (Ed), *Urbanagriculture in West Africa: Contributing to food security andUrban*
 311 *Sanitation*, CRDI, CTA, Ottawa (Canada). 19-40
- 312 2. Maxwell D. Sécurité alimentaire dans les centres urbains d'Afrique subsaharienne.
 313 Armer les villes contre la faim. Systèmes alimentaires urbains durables. CRDI, 2000,
 314 260 p.
- 315 3. Nations Unies, DAES. World population prospects: The 2005 Revision(2006).New
 316 York,
- 317 4. FAO. Agriculture urbaine et sécurité alimentaire. FAO, 2008.Salle de Presse.
 318 Rome : FAO
- 319 5. Smit J, Ratta A, Nassr J. Urban Agriculture: Food, Jobs and Sustainable Cities,
 320 1996. PNUD, New York, 302 p.

- 321 6. N'Dienor M. Fertilité et gestion de la fertilisation dans les systèmes maraichers
322 périurbains des pays en développement : Intérêts et limites de la valorisation
323 agricole des déchets urbains dans ces systèmes, cas de l'agglomération
324 d'Antananarivo (Madagascar). Thèse de Doctorat Institut National Agronomique
325 Paris-Grignon, 2006. 242 p.
- 326 7. Sposito T. Agriculture urbaine et périurbaine en Afrique de l'ouest : le cas de micro
327 jardin dans la municipalité de Dakar. Thèse de Doctorat, 2010. Université de Milan.
328 232p.
- 329 8. Melsted. Possibilités d'épandages des eaux usées urbaines. Etude bibliographique,
330 Agence du bassin Rhône-Méditerranée-Corse, I.N.R.A., 1979- S.C.P.A.R.P., 371 p.
- 331 9. Godin P. "Les sources de pollution des sols: essai de quantification des risques dus
332 aux éléments traces", Sciences du sol, 1983. p73-87.
- 333 10. Grandjean P, Anderson O, Berlin A, Grant L, Sors A, Tarkowski S, Victory W. Trace
334 elements in human health and disease. Report on second Nordic symposium, 17-21
335 August 1987, Odense. World Health Organization, Commission of European
336 Communities, US Environmental Protection Agency, 1987. 134p.
- 337 11. OMS. L'utilisation des eaux usées en agriculture et en aquaculture :
338 Recommandations à visées sanitaires. Genève, Suisse, OMS, Séries de rapports
339 techniques, 1989. n° 778, 83 p.
- 340 12. Walker C, Hopkin S, Sibly R & Peakall D. Principles of ecotoxicology. Taylor &
341 Francis. 1996, 321p.
- 342 13. Theresa A. F., Trace Element Monitoring and Therapy for Adult Patients Receiving
343 Long-term Total Parenteral Nutrition. Nutrition Issues in Gastroenterology, 2005,
344 25 : 44 – 51.
- 345 14. Suttle, N. F. Mineral nutrition of livestock. 4th Edition. Massachusset 2010: CAB
- 346 15. Baize D. Guide des analyses en pédologie. Ed. INRA, 2000. Paris, 257 p
- 347 16. Kouakou K. J. Etude des métaux traces (Cd, Cu, Pb, Zn, Ni) dans les sols et les
348 produits maraîchers de deux sites d'agriculture dans la ville d'Abidjan (Côte
349 d'Ivoire), Thèse de Doctorat Unique de l'université d'Abobo-Adjame, Option:
350 Biodiversité et gestion durable des écosystèmes, spécialité : Biogéochimie, Mémoire
351 soutenu publiquement le 04 juillet 2009, 145p
- 352 17. Kouamé K.I, Gone D.L, Savane I, Kouassi E.A, Koffi K, Goula B.T, Diallo M. Mobilité
353 relative des métaux lourds issus de la décharge d'Akouédo et risque de
354 contamination de la nappe du Continental Terminal (Abidjan-Côte d'Ivoire). Afrique
355 Sciences, 2006, 2(1): 39-56.
- 356 18. Ondo J. A. Vulnérabilité des sols maraîchers du Gabon (région de Libreville) :
357 acidification et mobilité des éléments métalliques. Thèse pour obtenir le grade de
358 Docteur de l'Université de Provence Spécialité Sciences de l'Environnement

- 359 Terrestre. Soutenue publiquement Le 29 Novembre 2011. École doctorale des
360 Sciences de l'Environnement. P 324
- 361 19. Toure Nantarie. Contamination des sols, eaux et produits maraîchers par des
362 Eléments Traces Métalliques dans la vallée du Nieké, dans le Sud Est de la Côte
363 d'Ivoire. Thèse de Doctorat Unique de l'Université Félix Houphouët-Boigny de
364 Cocody, 2012. Spécialité : *Pédologie*. 170 p.
- 365 20. Mench M, Baize D. Contamination des sols et de nos aliments d'origine végétale par
366 les éléments en traces, mesures pour réduire l'exposition, *Courrier de*
367 *l'environnement de l'INRA* n°52, septembre 2004, pp 31-56.
- 368 21. Bur T. Impact anthropique sur les éléments traces métalliques dans les sols
369 agricoles de midi-pyrenees. Implications en termes de limites et de charges
370 critiques, Thèse de Doctorat de l'Université de Toulouse, spécialité :
371 *Hydrogéochimie*, soutenue le 19 Décembre 2008, 399p.
- 372 22. Ahoussi K.E, Loko S, Koffi Y.B, Soro G, Oga Y.M. S, Soro N,. Evolution spatio-
373 Temporelle des teneurs en nitrates des eaux souterraines de la ville d'Abidjan
374 (CÔTE D'IVOIRE). *International Journal of Pure & Applied Bioscience*, 2013. 1 (3) :
375 45-60.
- 376 23. Gee G.W and Bauder J.W,. Particle-size Analysis. *In* : Klute A (eds) *Methods of Soil*
377 *Analysis. Part 1. Physical and Mineralogical Methods. Agronomy 9. Second Edition.*
378 ASA, SSSA, Madison, 1986. pp. 383-409.
- 379 24. Walkley A et Black I. A. An examination of Degtjareff method for determining soil
380 organic matter and a proposed modification of the chromic acid titration method. *I*
381 *Soil Science*, 1934, 37: 29-38
- 382 25. AFNOR. Norme française. AFNOR NF X31-415. Qualité des sols. Extraction des
383 éléments en traces solubles dans l'eau régale, AFNOR ,1995. 455-463.
- 384 26. Baize D, Sterckeman T, Piquet A, Ciesielski H, Béraud J, Bispo A. Dérogations
385 relatives à la réglementation sur l'épandage des boues de stations d'épuration.
386 Comment formuler une demande pour les sols à teneurs naturelles élevées en
387 éléments traces métalliques? 2005, 145 p.
- 388 27. Laurent C,. Les modèles d'adsorption des métaux en traces (Zn, Cd, Cu et Pb) dans
389 les sols: synthèse bibliographique et validation. Mémoire de stage de Diplôme
390 d'Etudes en Pollution Chimique et Environnement, Université Paris XI, Orsay 2003,
391 55 p.
- 392 28. Alloway B.J. *Heavy metals in soils*. Blackie Academic & Professional. Glasgow,
393 1990. 339p.
- 394 29. Ahoussi K.E., Koffi Y.B., Loko S., Kouassi A.M., Soro G., Biemi J. Caractérisation
395 des éléments traces métalliques (Mn, Ni, Zn, Cd, Cu, Pb, Cr, Co, Hg ,As) dans les
396 eaux superficielles de la commune de Marcory, Abidjan Côte d'Ivoire : cas du village
397 d'Abia Koumassi. *Geo-Eco-Trop* 2012, 36 : 159-174.

- 398 30. Colinet G., Lacroix D., Bock L..Importance of Pedogenic Factors in the Spatial
399 Distribution of some Soil Properties in the Belgian Loess Belt: II. Some Loess-
400 Substratum Contacts. *Geologica Belgica International Meeting*, 2002. Available at:
401 [http://orbi.ulg.ac.be/bitstream](http://orbi.ulg.ac.be/bitstream/2268/69487/1/ABS_SOL_COLINET%20et%20al_2002_09_Geologica%20Belgica_EN.pdf)
402 [/2268/69487/1/ABS_SOL_COLINET%20et%20al_](http://orbi.ulg.ac.be/bitstream/2268/69487/1/ABS_SOL_COLINET%20et%20al_2002_09_Geologica%20Belgica_EN.pdf)
2002_09_Geologica%20Belgica_EN.pdf
- 403 31. Coulibaly A. S., Mondé, Wognin V. A. & Aka K. State of anthropic pollution in the
404 estuary of Ebrié lagoon (Côte d'Ivoire) by analysis of the metal elements traces.
405 *European Journal of Scientific Research*, 2008. 19, 2: 372-390.
- 406 32. Soro G., Metongo S. B., Soro N., Ahoussi K. E., Kouame Koffi F. Zade S. G. P.
407 & Soro T.. Métaux lourds (Cu, Cr, Mn et Zn) dans les sédiments de surface d'une
408 lagune tropicale africaine : cas de la lagune Ebrié (Côte d'Ivoire). *International*
409 *Journal of Biological and Chemical Sciences*, 2009. 3, (6): 1408-1427.
- 410 33. Kone´ Y.J.M, Abril G, Delille B, Borges A.V. Seasonal variability of methane in the
411 rivers and lagoons of Ivory Coast (West Africa). *Biogeochemistry*, 2010. 100:21–37
- 412 34. Schwartz Ch, Zu Y, Li Y, Langlade L. Utilisation agricole des sols pollués par les
413 métaux lourds dans le Sud-ouest de la Chine. Colloque : Devenir et effets des
414 contaminants métalliques dans les agro écosystèmes et écosystèmes terrestres :
415 influence de l'usage des sols, 20 – 21 Mars 2003, Lile. Ministère de l'écologie et du
416 développement durable, Ingénieurs pour la terre, INRA, AFES2003. 103p.
- 417 35. Makela-Kurtto R. Effects of afforestation of agricultural land on heavy metal mobility
418 in soil (MEMO:FAIR3-CT96-1983): Individual final report of Partner 3 (MTT) for the
419 period 01-03-97 to 30-09-00. Agricultural Research Centre of Finland (MTT),
420 Resource Management Res, 2000. FI-31600 Jokioinen, Finland.
- 421 36. Foy, C. D., Chaney, R. L., and White, M. C. The physiology of metal toxicity in
422 plants, *Annu. Rev. Physiol.* 1978, 29 : 511- 566.
- 423 37. De Vries W, Bakker DJ, 1998. Manual for calculating critical loads of heavy metals
424 for terrestrial ecosystems. Guidelines for critical limits, calculation methods and input
425 data. Report 166, DLO Winand Staring Centre, Wageningen, the Netherlands, 144
426 pp.
- 427 38. Kabata-Pendia A. Trace elements in soils and plants. 4th edition, Taylor & Francis
428 and CRC Press, 2011. 467p.
- 429 39. Isermann, K. Method to reduce contamination and uptake of lead by plants from car
430 exhaust gases, *Environ. Pollut.* 1977, 12: 199-203.
- 431 40. Godin P. Les sources de pollution des sols : Essai de quantification des risques dus
432 aux éléments traces. AFES 2010: 73 – 87.
- 433 41. Jones, L. H. P., Jarvis, S. C., and Cowling, D. W. Lead uptake from soils by
434 perennial ryegrass and its relation to the supply of an essential element (sulphur),
435 *Plant Soil*, 1973. 38: 605 – 619.

- 436 42. Hassan N., Mahmood Q., Waseem A., Irshad M., Arshad P.F. Assessment of heavy
437 metals in wheat plants irrigated with contaminated wastewater. *Pol. J. Environ. Stud.*
438 2013. 22(1), 115 – 123.
439
- 440 43. Brookes PC, McGrath SP. Effect of metal toxicity on the size of microbial biomass.
441 *Journal of soil science*, 1984. 35:341-346
- 442 44. Adriano, D. C., Bollag, J. M., Frankenberger, Jr., W. T., and Sims, R. C.
443 *Bioremediation of Contaminated Soils*, Am. Soc. Agron.1999, Madison, WI, 800.
- 444 45. Barrios, Buresh R.J., Sprent, J.I. Organic matter in soil particle size and density
445 fractions from maize and legume cropping systems. *Soil Biology and Biochemistry*
446 1996. 28(2): 185 – 193.
- 447 46. Alexander P.D, Alloway B.J, Dourado A.M,. Genotypic variation in the accumulation
448 of Cd, Cu, Pb and Zn exhibited by six commonly grown vegetables. *Environ. Pollut.*
449 2006. 144:736–745.
- 450 47. Waseem A., Arshad J., Iqbal, F., Sajjad, A., Mehmood Z., Murtaza G. Pollution
451 Status of Pakistan: A Retrospective Review on Heavy Metal Contamination of
452 Water, Soil, and Vegetables. *BioMed Research International*, 2014., Hindawi
453 Publishing Corporation, 29p.
454