

Preliminary survey and ecological studies on ant fauna diversity in Asir region, Kingdom of Saudi Arabia

ABSTRACT

We report on twelve new records of ants species for the fauna in Asir region: *Camponotus aegyptiacus*, *Camponotus sericeus*, *Camponotus xerxes*, *Cataglyphis bicolor*, *Cataglyphis fortis*, *Cataglyphis niger*, *Lepisiota opaciventris*, *Paratrechina longicornis*, *Crematogaster aegyptiaca*, *Crematogaster senegalensis*, *Monomorium carbonarium*, and *Monomorium salomonis*. Genera like *Cataglyphis*, *Camponotus*, *Crematogaster* and *Messor* gradually dominate the ant fauna and are represented by a maximum number of endemic species with Myrmicinae accounting for the most numerous species followed by Formicinae. The maximum relative abundance and distribution of ant species in Asir province was observed in the Formicinae (*Cataglyphis bicolor*) and in the Myrmicinae (*Messor ebeninus*).
Keywords: *Ants*, *Formicidae*, *survey*, *diversity*, *Asir region*, *Kingdom of Saudi Arabia*.

1. INTRODUCTION

Ants belong to the family Formicidae of the order Hymenoptera with approximately 15,981 valid species and subspecies described worldwide (AntWeb, 2016), of an expected total of 22,000 species (Agosti and Johnson, 2003). They play an important role in maintaining the natural balance of ecosystem and can be used as bioindicators of environmental changes (Maryati, 1997; Peck et al., 1998; Toda and Kitching, 1999). Ants have been used extensively as bioindicators for used in land monitoring and assessment (Majer, 1983; Greenslade and Greenslade, 1984; Andersen, 1990, 1997) particularly to mine site restoration (Majer, 1984; Majer, 1985; Andersen, 1997), and also to disturbances such as fire, grazing and logging (Neumann, 1992; York, 1994; Vanderwoude et al., 1997a). Ants have been used as indicator taxa, most notably in Australia, where the diversity of ants is positively related to biodiversity in arid environments. They have also been used to monitor the effect of disturbances on biodiversity (Andersen et al., 1998; Andersen et al., 2007). Ants are considered to be an ideal candidate for use as an indicator group for a number of reasons: they are abundant and ubiquitous in areas of disturbance (Andersen, 1990), they exhibit strong interactions with all trophic levels (Briese, 1982). They are highly sensitive to environmental conditions and disturbance and they respond rapidly to environmental change (Vanderwoude et al., 1997a; Andersen 1990). Furthermore, sampling is relatively easy, without requiring enormous expertise and experience. Ants exhibit a greater resistance to pollutants in comparison to other invertebrates, especially to radioactivity (Torossian and Causse, 1968; Le Masne and Bonavita-Cougourdan, 1972) and to industrial pollutants (Petal et al., 1975). Ants are also excellent focal taxa for study and monitoring since they are found abundantly in most terrestrial ecosystems (Majer 1983; Hölldobler and Wilson 1990). The majority of ground dwelling ants live in perennial, sessile colonies with workers having limited foraging ranges (Hölldobler and Wilson 1990), allowing for reliable sampling and monitoring that is well-linked to characteristics of the local habitat (Alonso, 2010). Ants are very diverse and abundant, exhibit many types of relationships with other soil biota, and are found in most terrestrial environments of the planet. They have ability to maintain or restore soil quality and their value as bioindicators of soil quality (Lobry de Bruyn, 1999). The direct and indirect contribution of ants to soil health integrity, and the resilience of their response to disturbance. They also play important ecological roles, having direct interactions

Commentaire [1]: This part is new! But it is an almost textual copy of 4 lines from a new source (King et al. 1998, p. 1628, first paragraph of the introduction) that I suggested in my previous evaluations, and which is not cited.

Commentaire [2]: See lines 55-56 of the previous version. This part is still a textual copy from Glasier (2012), but the reference has been deleted.

Commentaire [3]: See lines 56-58 of the previous version. This part is also a textual copy from Glasier (2012), but the reference has been deleted.

Commentaire [4]: See lines 29-31 and 33-36 of the previous version. This part is still a textual copy from Orabi et al. (2011), but the reference has been deleted.

Supprimé: Coug ourdan

Commentaire [5]: This part is new! But it is a textual copy of 3 lines from a new source (Folgarait 1998, p. 1228, first paragraph on "Pollution") that I suggested in my previous evaluations, and which is not cited.

Commentaire [6]: See lines 50-55 of the previous version. This part is still a textual copy from Glasier (2012), but the reference has been deleted.

Commentaire [7]: This part is new! But it is also a textual copy of 3 lines from the previously mentioned Folgarait's article (1998, p. 1237, mid of the "Conclusion"), and the reference is still not cited.

Commentaire [8]: ???
None of these two sentences are complete.

with plants and animals at all trophic levels. Many of these roles are related to paedogenesis, nutrient cycling (Hutson, 1989), seed predation (Andersen, 1990; Majer, 1990) seed dispersal (Majer, 1990) and decomposition of organic matter. Ants may have positive or negative influences in cases where habitats are rehabilitated by seeding plants depending on their seed preferences or on the presence of elaiosomes (food ant rewards) in the seeds because the ants may act as granivores (negative effect) or dispersers (positive effect) (Majer, 1984). Ants are an excellent taxon for biodiversity studies because they are conspicuous and important faunistic elements in most ecosystems. For example, ants contribute as much to soil turnover as terrestrial annelids (Hölldobler and Wilson 1990). In areas with no termites, such as northern temperate regions, ants are also major contributors to the breakdown of wood (Hasen and Klotz, 2005).

2. MATERIAL AND METHODS

Study area (Asir region)

South western region (Fig.1) is located between latitude 17°30'-21°00' N and longitude 41°30'-44°30' E. It is entirely different from the rest of the kingdom. The mountain range of Al-Sarawat with its highest peak (Al-Sawdah mountain, 3000-3100 m), near Abha (2094 m) the capital of Asir province, divides the area into two distinct ecological regions. There are the high plateau with the slopes gently eastward from the escarpment to form the highlands of Asir, and to the west the highly jagged-edged mountains terrain to Tehamah Asir that merges westward into the narrow, sandy Red Sea coastal plain, making it the hottest part of the region. The Asir highland receives a variable seasonal rainfall which is higher than in the rest of the kingdom, i.e. 300-500 mm/year as compared with 50-100 mm/year elsewhere.

Field techniques

Two field techniques were used to collect and study ants mainly Pitfall trapping (Four traps for each location according to the four geographical directions and a distance between the pitfall traps of about 500 m) and direct sampling (Hand picking). Ants were collected in the field and then put into vials filled with 70% ethanol and tight-fitting caps to retard the evaporation of alcohol. For every sample, the vial was clearly labeled with a code number referring to the location, the date, the habitat....etc. These samples were collected every two weeks for one year (Nov.2013- Oct. 2014).

Sorting ant specimens

Two important divided into from the field, ants preparations was the sorting of the materials collected (mixed with soil and other organisms). After field work, ants specimens were important to remove from this other materials as soon as possible in order to prevent damage specimens (Lattke 2000). Salt-water extraction technique described by (Agosti et al., 2000) was used in order to separate ant specimens from the other materials because of its simplicity and low cost.

Identification of the ants to species

Examination and identification of ants to species rank was carried out by the use of a stereoscopic binocular microscope. Identification of ants to the species rank was obtained by the use of different taxonomic keys (Collingwood, 1985; Agosti, 1990; Bolton, 1995; Collingwood & Agosti, 1996; Cagniant, 2009).

Diversity data analysis

Data were analyzed by Incidence-based coverage estimator (ICE). Species richness and Alpha diversity were estimated by using Shannon-Wiener, and Simpson's D diversity indices. The program EstimateS (Colwell, 2009) was used to calculate these standard estimators.

Commentaire [9]: See lines 26-29 of the previous version. This part is still a textual copy from Orabi et al. (2011), but the reference has been deleted.

Commentaire [10]: This part is new! But it is also a textual copy of the 4 last lines of the paragraph on "Pollution" in the same Folgarait's article (1998, p. 1228), and the reference is still not cited.

Commentaire [11]: See lines 46-50 of the previous version. This part is still a textual copy from Glasier (2012), but the reference has been deleted.

Supprimé: was

Commentaire [12]: ??? What is the meaning?

Commentaire [13]: ??? What is the meaning?




Fig.1 Asir map showing the surveyed localities.

3. RESULTS AND DISCUSSION

Results of ant diversity studies

Table.1 Data of the surveyed sites in Asir province with number of specimens at each site.

	Area (Area more than 2000 m elevation)	Coordinates of latitude and longitude		Above sea level (altitude) asl	Total no. specimen collected	No. species
		N	E			
1	Abha (AB)**	18°14'460"	42°31'767"	2094 m	176	7
2	Al-Sawdah (AS)**	18°27'173"	42°38'400"	3000 - 3100 m	50	2
3	Sarat Abidah (SA)**	18°00'852"	43°09'00"	2380 m	56	2
4	Ahad Rafidah (AR)**	18°21'216"	42°84'378"	2090 m	28	2
5	Dhahran Al-Janoub (DJ)**	17°39'56.1 "	43°31'00.1 "	2100-2570 m	40	4
6	Ballahmar (BA)**	18°54'947"	42°40'899"	2250 m	150	9
7	Ballasmar (BS)**	18°79'296"	42°25'570"	2250 m	20	2
8	Tanomah (TA)**	18°89'215"	42°22'767"	2148 m	34	3
9	Al-Namas (AN)**	19°07'12"	42°07'48"	2600 m	73	4
(Area below 2000 m elevation)						
10	Muhayil (MU)*	18°34'760"	42°02'151"	456 m	115	5
11	Al-Majardah (AM)*	19°12'844"	41°92'460"	2000 m	33	4
12	Khamis Mushayt (KM)*	18°29'153"	42°73'874"	1850 - 2000 m	132	6
13	Rijal Almaa (RA)*	18°14'608"	42°16'500"	1000- 1500 m	113	5

Table.2 Mean Environmental parameters in both elevation gradients in 2013-2014.

Altitude	Less than 2000 m	More than 2000 m
Temperature (Mean)	25	15.5
Relative humidity	51	45
Wind speed	7	10
Precipitation	35	30
Specimen	393	627

Fig.2 Abundance effectiveness of both collection methods in both elevation




Table.3 Distribution of the ants in the Asir region, KSA

Region species		Al- Sawdah	Al-Namas	Dhahran Al Janoub	Sarat Abidah	Ballahmar	Ballasmar	Tanomah	Abha	Ahad Rafidah	Al- Majardah	Khamis Mushayt	Rijal Almaa	Muhayil
subfamily: Formicinae	<i>Camponotus aegyptiacus</i> Emery												++	
	<i>Camponotus xerxes</i> Forel								+					
	<i>Camponotus flavomarginatus</i> Mayr			+	++	+		+	++			+		
	<i>Camponotus sericeus</i> (Fabricius)									+	+	++	+	+
	<i>Cataglyphis albicans</i> (Roger)		+									+		
	<i>Cataglyphis nigra</i> (André)		+					++						
	<i>Cataglyphis bicolor</i> (Fabricius)			+	++				++		+	+		+
	<i>Cataglyphis fortis</i> (Forel)	+							+					
	<i>Lepisiota opaciventris</i> (Finzi)		+											
	<i>Paratrechina longicornis</i> (Latreille)												+	+
subfamily: Myrmicinae	<i>Crematogaster aegyptiaca</i> Mayr					+	+		+					
	<i>Crematogaster senegalensis</i> Roger										+		++	++
	<i>Monomorium carbonarium</i> (Smith)					+			+	+				
	<i>Monomorium sp.</i>					+					+	+		+
	<i>Monomorium salomonis</i> (Linnaeus)					+								
	<i>Messor ebeninus</i> Santschi	++		+		++	+	+	++			++		
	<i>Pheidole megacephala</i> (Fabricius)			+		+						+	++	
	<i>Tetramorium sericeiventris</i> Emery		+			+								
	<i>Tetramorium depressiceps</i> Menozzi					+								

(+) low density, (++) moderate density and (+++) high density

Table.4 Showing alpha diversity of ants in both elevational gradients

SAMPLE	ALPHA SD (ANALYTICAL)	SHANNON MEAN	SIMPSON MEAN
DIRECT SAMPLING	0.79	2.47	9.30
PITFALL TRAPPING	0.43	2.56	12.14

Diversity of Ants along elevational gradient

A total of 1,020 ants belonging to 19 species of 9 genera were collected. *Lepisiota*, *Paratrechina*, *Messor* and *Pheidole* genera were represented by a single species each, while *Camponotus*, *Cataglyphis*, *Monomorium*, *Crematogaster* and *Tetramorium* were represented by 4, 4, 3, 2 and 2 species respectively (Table.2 & Fig.2). Formicinae were represented by 4 genera and Myrmicinae by 5 genera. Relative abundance was higher for *Messor ebeninus*, *Cataglyphis bicolor* and *Crematogaster senegalensis* (17.65%; 13.93% and 11.67% respectively) while *Lepisiota opaciventris* had the lowest 0.49%. Hand picking yielded higher number of specimens (68.23%) than Pitfall Trapping (31.77%) (Fig.2).

At <2000 m the average temperature was 25°C, relative humidity 51%, wind speed was 7km/h and precipitation 35mm. The total catch in terms the number of specimens was 393. At the site of KM, the number of specimens collected was 132 (33.59%) with a maximum of 6 different species recorded. The genus *Crematogaster* was the most abundant with *Crematogaster senegalensis* 119 representing 30.28% of the total catch at this elevation and most the specimens were collected by hand picking method. Formicinae were represented mainly by the genus *Camponotus*, *Camponotus sericeus* has maximum catch of 78 specimens and forming the largest percentage 19.85%, again hand picking method was found to be the most effective followed by pitfall (Table.2 & Fig.2).

At >2000 m the average temperature was 15.5°C , relative humidity 45%, wind speed was 10 km/h and precipitation 30mm. The total catch in terms the number of specimens was 627. In terms of abundance, 176 (28.07%) specimens with 7 different species were recorded at the site of AB, while the higher diversity, with 9 different species, was observed at the site of BA. The genus *Messor* was the most abundant genus with *Messor ebeninus* representing 22.17% of the total catch and most of the specimens were collected by the hand picking method. Formicinae represented 51.35% of the specimens, with the genus *Cataglyphis* forming the bulk (32.69%), *Cataglyphis bicolor* accounting for the higher catch with 103 specimens. (Table.2 & Fig.2)

Discussion

Ants diversity along elevational gradient

Several studies have examined the effects of elevational gradients on ant diversity. In general these studies have revealed two patterns: one showing a steady decline in ant diversity from low to high elevations (Brühl *et al.*, 1998; Janzen *et al.*, 1976), and another a pattern of mid-elevation peak in species richness (Fisher & Robertson 2002; Sabu *et al.*, 2008; Samson *et al.*, 1997). One common element to these studies is that they have tended to quantify biodiversity using species diversity metrics that are based on morphospecies designations or morphologically determined taxonomic units (Gotelli & Colwell 2001).

In the present work, we also used morphology as a direct measure of the levels of ant functional diversity present at different elevational gradients. Therefore, we categorized our study area in two section one is <2000 m and the other >2000 m with 13 collecting sites. Also we tried to explain ant species to a set of guilds using both ecological information as well as a group of functionally significant morphological traits. In general, we found that functional diversity peaked at mid-elevation (although non-significantly). The results of these guild proportionality analysis show that at low and high elevations, ant communities contain uneven numbers of species from each guild and contain less of the overall variation in ant functional morphology suggesting that environmental factors like temperature; humidity etc.

Commentaire [14]: See comment 20 in the first revision.

Commentaire [15]: See comment 21 in the first revision.

may previously controlled these communities. As have mentioned we found a mid-elevational peak of species density that differs in both size and shape depending on the season and year. Abha (AB) and Ballahmar (BA) sites were dominant in terms of number of species (Table.1). These results are in accordance with studies that found mid-elevational peaks of ant species density (e.g. Fisher, 1999; Sanders, 2002; Bharti et al., 2013). This peak was that found to change in scale and position depending on the season (Table.2 & Fig.2). Larger community structure, abundance and variation were found in these sites, which may be because of suitable ecological factors like temperature, and humidity and these areas are known to have maximum rainfalls in kingdom (Table.2). Fewer species were observed in the dry season than in the wet season (Table.2). In our dataset, the seasonal change in the elevation–species density relationship is due to greater seasonal fluctuation at low elevations. These patterns of species density appear to be entirely explained by the effects of mean temperature and available area. In general, a weak linear decline in abundance across elevation was seen in the wet season (Table.2 & Fig.2). Species richness by different indices have been depicted in (Fig.2) and species abundance and effectiveness of sampling methods by Sobs (species observed) while Alpha diversity indices have been depicted in (Table.4). The data generated reflected that with decreasing in temperature and humidity, composition of species changes; as in case of Myrmicinae the generalist species are replaced by more high altitude specialists like *Messor ebeninus* Santschi, *Crematogaster senegalensis* Roger, and *Cataglyphis bicolor bicolor* (Formicinae). In the case of the Formicinae the interpretation resembles Myrmicinae as cold specialist *Camponotus sericeus* increases in abundance. But interestingly, the overall abundance increases from 1800 m to 2500 m with the number of species staying almost the same at both the elevations. For the time being, it is difficult to conclude that with more increase in altitude, the number of species and abundance would increase, but it is what we observed that with an increase in altitude in Asir.

Ants taxonomy with distribution

The results of the present study record the presence of nineteen species: *Camponotus aegyptiacus*, *Camponotus sericeus*, *Camponotus xerxes*, *Camponotus flavomarginatus*, *Cataglyphis albicans*, *Cataglyphis bicolor*, *Cataglyphis fortis*, *Cataglyphis nigra*, *Lepisiota opaciventris*, *Paratrechina longicornis*, *Crematogaster aegyptiaca*, *Crematogaster senegalensis*, *Messor ebeninus*, *Monomorium carbonarium*, *Monomorium* sp., *Monomorium salomonis*, *Pheidole megacephala*, *Tetramorium sericeiventris* and *Tetramorium depressiceps* from two subfamilies (Formicinae and Myrmecinae).

Twelve species were new record in Asir region: *Camponotus aegyptiacus*, *Camponotus sericeus*, *Camponotus xerxes*, *Cataglyphis bicolor*, *Cataglyphis fortis*, *Cataglyphis nigra*, *Lepisiota opaciventris*, *Paratrechina longicornis*, *Crematogaster aegyptiaca*, *Crematogaster senegalensis*, *Monomorium carbonarium* and *Monomorium salomonis*.

Five species were previously recorded and also found in new places: *Camponotus flavomarginatus*, *Messor ebeninus*, *Pheidole megacephala*, *Tetramorium sericeiventris* and *Tetramorium depressiceps* and two species were previously recorded in Asir region: *Cataglyphis albicans* and *Monomorium* sp.

4. CONCLUSION

Diversity of Asir region ant fauna in relation with the elevation gradient reveals that the functional diversity peaked at mid-elevation in Asir region. The data generated reflected that the composition of species changes in accordance with environmental factors like temperature, humidity etc. More ant individuals and species were found in the wetter parts of the elevation gradient rather than in hot and dry areas. These results suggest a strong environmental, control on ant diversity patterns in Asir region of KSA. At higher elevations, genera like *Cataglyphis*, *Camponotus*, *Crematogaster* and *Messor* gradually dominate the

Commentaire [16]: See comment 22 in the first revision.

Commentaire [17]: English ???

Commentaire [18]: See comment 18 in the second revision.

Commentaire [19]: See comment 24 in the first revision.

Commentaire [20]: See comment 26 in the first revision.

Commentaire [21]: See comment 27 in the first revision.

ant fauna and are represented by a higher number of endemic species. Maximum relative abundance and distribution of ant species in Asir province were observed for the Formicinae *Cataglyphis bicolor* and the Myrmicinae *Messor ebeninus*.

ACKNOWLEDGEMENTS

The authors are indebted to the King Abdulaziz City for Science and Technology (KACST) Saudi Arabia, for financial support of this work (KACST: AI-34-0225).

REFERENCES

- Agosti, D. (1990). Review and reclassification of *Cataglyphis* (Hymenoptera Formicidae). *Journal of Natural History*, 24: 1457-1505.
- Agosti, D. Majer, J.D. Alonso, L.E. and Schultz, T.R. (Eds.) (2000). *Ants: Standard Methods for Measuring and Monitoring Biodiversity*. Smithsonian Institution Press, Washington and London. 280 pp.
- Alonso, L.E. (2010). Ant conservation: current status and call for action. In L. Lach, C.L. Par & K.L. Abbott (Eds), *Ant Ecology*. Oxford University Press, pp. 59-74.
- Andersen, A.N. (1990). The use of ant communities to evaluate change in Australian terrestrial ecosystems: a review and a recipe. *Proceedings of the Ecological Society of Australia*, 16: 347-357.
- Andersen, A.N. (1997). Measuring invertebrate biodiversity surrogates of ant species richness in the Australian seasonal tropics. *Mem. Mus. Victt.*, 56:355-359.
- Andersen, A.N., Morrison, S. Belbin, L., Ashwath, N. and Brennan, K. (1998). The role of ants in minesite restoration in the Kakadu region of Australia's Northern Territory, with particular reference to their use as bioindicators. *Supervising Scientist Report 130*, Supervising Scientist, Canberra. 98 pp.
- Andersen, A.N., Parr, C.L., Lowe, L.M. and Müller, W.J. (2007). Contrasting fire-related resilience of ecologically dominant ants in tropical savannas of northern Australia. *Diversity and Distributions* 13: 438-446.
- Bharti, H., Sharma, Y.P. Bharti, M. and Pfeiffer, M. (2013). Ant species richness, endemism and functional groups, along an elevational gradient in the Himalayas. *Asian Myrmecology*. 5: 79-101.
- Bolton, B. (1995). *A New General Catalogue of the Ants of the World*. Harvard Univ. Press, Cambridge, Mass. 504 pp.
- Briese, D.T. (1982). The effect of ants on the soil of a semi-arid saltbush habitat. *Insectes Sociaux*, 29: 375-382.
- Brühl, C.A. Gunsalam, G. and Linsenmair, K.E. (1998). Stratification of ants (Hymenoptera, Formicidae) in a primary rain forest in Sabah, Borneo. *Journal of Tropical Ecology*, 14: 285-297.
- Cagniant, H. (2009). Le genre *Cataglyphis* Foerster, 1850 au Maroc (Hyménoptères Formicidae). *Orsis*, 24: 41-71.
- Collingwood, C.A. (1985). Hymenoptera: Fam. Formicidae of Saudi Arabia. *Fauna of Saudi Arabia*, 7: 230-302.
- Collingwood, C.A. and Agosti, D. (1996). Formicidae (Insecta: Hymenoptera) of Saudi Arabia (Part 2). *Fauna of Saudi Arabia*, 15: 300-385.
- Colwell, R.K. (2009). EstimateS: Statistical estimation of species richness and shared species from samples. Version 8.2. <http://viceroy.eeb.uconn.edu/EstimateS>.
- Fisher, B.L. (1999). Ant diversity patterns along an elevational gradient in the Réserve Naturelle Intégrale d'Andohahela, Madagascar. *Fieldiana Zoology* (n.s.) 94: 129-147.
- Fisher, B.L. and Robertson, H.G. (2002). Comparison and origin of forest and grassland ant assemblages in the High Plateau of Madagascar (Hymenoptera: Formicidae). *Biotropica* 34: 155-167.

- Gotelli, N.J. and Colwell, R.K. (2001). Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters* 4, 379-391.
- Greenslade, P.J.M. and Greenslade, P. (1984). Invertebrates and environmental assessment. *Environment and Planning*, 3: 13-15.
- Hansen, L.D. and Klotz, J.H. (2005). *The Carpenter Ants of the United States and Canada*. Comstock Publishing Associates, Ithaca, New York. 204 pp.
- Hölldobler, B. and Wilson, E.O. (1990). *The Ants*. The Belknap Press of Harvard University Press, Cambridge, Massachusetts, USA. 732 pp.
- Hutson, B. R. (1989). The role of fauna in nutrient turnover. In: J. D. Majer (ed.), *Animals in primary succession: The role of fauna in reclaimed lands*. Cambridge University Press, Cambridge, pp. 51-70.
- Janzen, D.H., Ataroff, M., Fariñas, M., Reyes S., Rincon, N., Soler A., Soriano, P. and Vera, M. (1976). Changes in arthropod community along an elevational transect in the Venezuelan Andes *Biotropica*, 8: 193-203..
- Latke, J.E. (2000). Specimen processing—building and curating an ant collection. In D. Agosti, J.D. Majer, L.E. Alonso & T.R. Schultz (eds.), *Ants: Standard Methods for Measuring and Monitoring Biodiversity*. Smithsonian Press, Washington. Pp.155–171.
- Le Masne, G. and Bonavita-Cougourdan, A. (1972). Premiers results tsd'une irradiation prolongee au cesium sur les populations de fourmis en Haute-Provence. *Ekologia Polska* 20, 129-144.
- Lobry de Bruyn, L. (1999). Ants as bioindicators of soil function in rural environments. *Agriculture, ecosystems & environment*, 74: 425-441.
- Majer, J.D. (1983). Ants: Bioindicators of minesite rehabilitation, land-use, and land conservation. *Environmental Management*, 7: 375-383.
- Majer, J.D. (1984). The influence of ants on seeding operations in Northern Australian mined areas. *Reclam. and Reveg. Resear*, 2: 299-313.
- Majer, J.D. (1985). Recolonisation by ants of rehabilitated mineral sand mines on North Stradbroke island, Queensland, with particular reference to seed removal. *Aust. J. Ecol.* 10: 31-48.
- Majer, J.D. (1990). Rehabilitation of disturbed land: long-term prospects for the recolonization of fauna. *Proceedings of the Ecological Society of Australia*, 16: 509-519.
- Maryati, M. (1997). Ants: an indicator for the tropical rain forest. *Manual for International Plot Course on Environmental Evaluation Using Insects as Indicator of Biodiversity: Ant Ecology, Taxonomy Collecting Methods and Identification* 17 March - 7 April 1997. Kota Kinabalu. Tropical Biology and Conservation Unit, Universiti Malaysia Sabah and International Institute of Entomology.
- Neumann, F.G. (1992). Responses of foraging ant populations to high-intensity wildfire, salvage logging and natural regeneration processes in *Eucalyptus regnans* regrowth forest of the Victorian central highlands. *Australian Forestry*, 55: 29-38.
- Peck, S.L. Mcquaid, B. and Campbell, C.L. (1998). Using ant species (Hymenoptera: Formicidae) as a biological indicator of agroecosystem condition. *Environmental Entomology*, 27: 1102-1110.
- Petal, J., Jakubczyk, H., Chmielewski, K. and Tatur, A. (1975). Response of ants to environmental pollution. In *Progress in Soil Zoology* (J. Vabek, ed.) pp. 363-373. Prague: Czechoslovakia Academy of Sciences.
- Sabu, T.K., Vineesh, P.J. and Vinod, K.V. (2008). Diversity of forest litter-inhabiting ants along elevations in the Wayanad region of the Western Ghats. *Journal of Insects Science*, 8: 69.
- Samson, D.A., Rickart, E.A. and Gonzales, P.C. (1997). Ant diversity and abundance along an elevational gradient in the Philippines. *Biotropica*, 29: 349-363.
- Sanders, N.J. (2002). Elevational gradients in ant species richness: area, geometry, and Rapoport's rule. *Ecography*, 25:25-32.

316 Toda, M. and Kitching, R.L. (1999). IBOY-DIWPA: Biodiversity Assessment Program in the
 317 Western Pacific and Asian region. International Net Work for Biodiversity Research
 318 Diversity (IBOY). Western Pacific and Asia.
 319 Torossian, C. and Causse, R. (1968). Effets des radiations gamma sur la fertlité et la
 320 longévité des colonies de *Dolichoderus quadripunctatus*. Compte rendu du Colloque
 321 ``Isotopes and radiation in entomology". International Atomic Energy Agency, Vienna.
 322 pp. 155-164.
 323 Vanderwoude, C., Andersen, A.N. and House, A.P.N. (1997). Ant communities as bio-
 324 indicators in relation to fire management of spotted gum (*Eucalyptus maculata* Hook.)
 325 forests in south-east Queensland. Memoirs of the Museum of Victoria, 56: 671-675.
 326 York, A. (1994). The long term effects of fire on forest ant communities: management
 327 implications for the conservation of biodiversity. Mem. Queensl. Mus. 36: 227-239.
 328