

# Assessment and Importance of Sesame Webworm, *Antigastra Catalaunalis* (Duponchel) in Western Tigray, North Ethiopia

## ABSTRACT

*Sesame (Sesamum indicum L.) is one of the important oil seed crops in Ethiopia. It is famous for international market and its production is challenged by inappropriate agronomic practices, insect pest infestation and weather uncertainties. Sesame webworm, Antigastra catalaunalis is a major pest in western Tigray. To assess the incidence and severity of A. catalaunalis, a survey was conducted in western zone of Tigray (Kafta Humera, Tsegede and Welkait) in 2015 cropping season starting from seedling up to capsule development stages. A total of 48 farm plots was assessed for prevalence, incidence and severity. All surveyed farm plots were infested at all stages of the crop. In the study area incidence of the pest was 66% and 15% capsule damage/severity. Capsule developmental stage of the crop was the most infested crop growth stage in the surveyed area and 100% incidence was also observed in late sown sesame (end of July). An increasing infestation trend of the pest was also observed as altitude is decreased. Therefore, A. catalaunalis is an economic pest in western zone of Tigray.*

**Key words:** *Antigastra catalaunalis, sesamum indicum, survey, incidence, severity*

## 1. INTRODUCTION

Sesame (*Sesamum indicum L.*) is an annual plant that belongs to the family *Pedaliaceae*. It is one of the world's oldest oil seed crop grown mainly for its high oil content of the seeds that contain 52 to 57% oil and 25% protein content [1, 2]. Sesame is mainly cultivated between the 25° N and 25° S latitudes. It is productive under high temperature condition and can grow on residual moisture at the end of a rainy season. Despite its ideal adaptation to dry sites, sesame can also be cultivated in humid, tropical and sub-tropical regions. Although, the order of leading sesame producer countries is changing from time to time but Ethiopia is the Sixth largest sesame producer in the world following Myanmar, India, China, Sudan and Tanzania, respectively and third in Africa preceding Uganda and Nigeria, respectively [3, 4].

In Ethiopia, sesame seed production is the leading oil crop and more than four million smallholder farmers are relying on the sesame production for their livelihood [5]. It is mainly being produced in Tigray (36%), Oromia (17%) Benishangul Gumuz (15%), Amara (31%) in the country [6].

Within Tigray the Western zone is the main production area with many large commercial and small scale farmers (about 90,000 producers) and it is a good source of income in these areas. Production of sesame in the country is very crucial in many aspects, but there are many hurdles for its production and productivity, such as pest infestation, seasonal delay, low yielding, post-harvest lost, poor storage facility, low fertilizer response, difference in capsule maturity, shattering etc. Of these, insect pest is one of the most important factor affecting production of sesame both in quality and quantity [1, 7].

Insect pests such as Sesame webworm (*Antigastra catalaunalis*), sesame seed bug (*Elasmolomus sordidus*), gall midge (*Asphondilia sesami*), and Indian meal moth (*Plodia interpunctella*) are the major insects of sesame during its different growing stages and post-harvest. Sesame webworm (*Antigastra catalaunalis*) is the most serious pest during various sesame growth stages starting 2-3 weeks after emergence up to harvesting[8-15].

- The objective of this study was to assess the incidence and severity of sesame webworm in western zone of Tigray.

## 2. MATERIALS AND METHODS

### 2.1 Description of the Study Areas

The Tigray regional state, where the study was conducted is located in the northern tip of Ethiopia and is bordered by Eritrea in the north, Sudan in the west, Afar region in the east, and Amhara region in the south. The survey districts include *Kafta Humera*, lowlands of *Tsegede* and *Wolkait* extends from 13°26' to 14°05'N latitude and 36°57' to 37°27' E longitude. Mixed farming system is the main occupation of the farm house holds. The districts are known for their high potential for sesame, sorghum and cotton production. Except for the very small areas under vegetables and fruits, crops are grown under rain fed condition. According to the Zonal Offices of Agriculture, western zone of Tigray has a total of 582,030.8 ha agricultural land, of which about 305,065 ha (99,704 ha of investors) is conducive for sesame production and has 89974 (577 investors) producers. *Kafta Humera* has an average monthly temperature and average annual rainfall of 28.2°C and 576.4mm, respectively. The lowlands of *Tsegede* and *Wolkait* have an average monthly temperature and annual rainfall of 27°C and 888.4 mm and 24.3°C and 750 mm, correspondingly. And the altitude of the surveying district ranges from 582 to 1063 m.




Figure 1 Map of the study areas

### 2.2 Surveying Method

Survey of *A. catalaunalis* was conducted in western Tigray, north Ethiopia (Figure 1) four times starting from seedling up to maturity (Figure 2). From the three woreda four *Tabias* each were selected purposely (sesame growing areas) and from each *Tabia* four farms/farm plots were selected randomly. Generally, a total of 48 sesame farm plots was assessed from both small scale farmers and investors and in each growth stage data was recorded from same

74 farm plots. The farm plots were sampled randomly at intervals of about 5-10 km [16] along  
 75 accessible roads until sesame field are found nearby, starting from *Maygebato Maydelie*. In  
 76 each farm plot, to assess the plot level of *A. catalaunalis* infestation at various crop growth  
 77 stages, sample were taken at 10-20-meter interval once at four points diagonally along the  
 78 field using 1 x 1 m quadrant randomly (Figure 1). The quadrant was thrown randomly and  
 79 plants inside the quadrant (both healthy and sesame-webworm infected) were counted to  
 80 determine the incidence of the pest. In addition to incidence, capsule damage was recorded  
 81 during capsule development and/or maturity stage through selection of five infected plants with  
 82 in the quadrant; then infected and non-infected capsules were counted for each of the five  
 83 plants in order to calculate capsule damage by dividing the infected capsules to the total  
 84 capsules in all the five plants. Infestation of *A. catalaunalis* was assessed also on early and  
 85 late grown cultivated and uncultivated sesame. During surveying elevation (altitude) was  
 86 measured using Garmin GPS for each location (farm plots).



87  
 88  
 89 • Figure 2 Sampling for *A. catalaunalis* infestation at different growth stages of sesame

### 90 2.3 Data Collection

- 91 • **Incidence:** mean percentage of infected plants showing typical symptoms (like  
 92 webbing of leaf and/or flower, leaf feeding, hole in capsule, the larva it self.) of  
 93 sesame webworm per total plant in the quadrant was recorded (Figure 3a) so as to  
 94 calculate incidence as follow:

$$\text{Incidence} = \frac{\text{infested plants}}{\text{total visited plants}} \times 100 \dots \dots \dots 1$$

- 95 • **Severity/Capsule Damage:** any capsule that showed sesame webworm symptom like  
 96 bored hole was considered as a damaged capsule (Figure 3b). Percentage of damaged

capsules by sesame webworm was calculated by counting the total damaged capsules of five plants in the quadrant using the following equation:

$$\text{Capsule Damage} = \frac{\text{no. of infested capsules}}{\text{total visited capsules}} \times 100 \dots \dots \dots 2$$

- **Agronomic data:** variety, planting date, growth stage of the crop and altitude of the fields were recorded.



Figure 3 Damaged sesame leaf, shoot and flower (A); damage sesame capsule and seed (B) by *Antigastra catalaunalis*

## 2.4 Data Analysis

Survey data was analyzed by Genstat-14 in split plot design arrangement. *Woredas* were taken as whole plots, while *Tabias*/localities were considered as sub plots, and means were compared through Fisher's LSD.

## 3. RESULTS AND DISCUSSIONS

### 3.1 Prevalence of *Antigastra catalaunalis* in Western Tigray

All the 48 sesame farm plots assessed from the three major sesame producing districts were found affected by *A. catalaunalis*, which clearly indicated that buildup of the pest across the districts. The survey result showed that in all the crop growth stages (at seedling, vegetative, flowering and fruiting stages) and irrespective of location all the sesame fields were found positive to *A. catalaunalis*.

### 3.2 Incidence of *A. catalaunalis* Affected by Sowing Time of Sesame (Cultivated and Uncultivated)

In addition to the cultivated one, sesame is also found as a wild grow/uncultivated in road sides, uncultivated lands and forest areas in western Tigray, north Ethiopia. And infestation of *A. catalaunalis* was observed in both cultivated and uncultivated sesame. The response of *A. catalaunalis* incidence to sowing/growing time of cultivated and uncultivated sesame is

demonstrated in (Figure 4). In the surveying areas 100% incidence was recorded from late sowing and late grows (end of July) in both farm and wild grown sesame, when a minimum incidence was recorded on the early sowing (early July) and a negligible incidence (less than 5%) was observed in the mid-June grown uncultivated sesame (Figure 4). There were no sesame fields sown in mid-June, because there was no ample rainfall during the specified month in the surveyed areas. The overall *A. catalaunalis* incidence was higher (68.12%) for cultivated sesame, when a minimal incidence (41.67%) was recorded for uncultivated sesame. The point behind the lower incidence on the uncultivated sesame might be that most uncultivated sesame grown early (June), because they do not demand too much rainfall/moisture like that of the cultivated sesame for emergence since they were buried shallowly in the soil and might emerge as immediate as the first shower since there were no soil disturbance also. Finally, incidence become lower because of the unfavorable weather conditions in the early grown/sown sesame. This study is in harmony with Abdel-rahman [17] who reported that delay in sesame sowing increased incidence of *A. catalaunalis*. Similarly investigations on the effect of sowing time of sesame on the occurrence of *A. catalaunalis* revealed that the incidence of the pest was highest (19.5%) in sesame sown late and lowest (8%) in early sown (June) [18].




Figure 4 Response of *A. catalaunalis* incidence to sowing time farm and wild grown sesame in western Tigray

### 3.3 Response of *A. catalaunalis* Incidence to Sesame Growth Stages

The means and significant variations ( $p < 0.01$ ) of *A. catalaunalis* incidence, among the growth stages (Figure 5), districts and specific locations (Table 1) surveyed are illustrated. The highest incidence (65.79%) of *A. catalaunalis* was observed during the capsule developmental stage, while the lowest (15.04%) was at seedling stage in the area of survey. The point behind the higher infestation at the later growth stage of the plant could be that the pest hides itself in bored pods and flowers, which the probability of the pest being in contact with the foliar sprayed insecticide become low. In a sense the pest is serious at pod developmental stage. This is in line with Muzaffar [19] who reported that during capsule formation phase of the crop infestation of the pest was higher than vegetative growth and flowering stages. Ahirwar, Gupta [8] also reported higher incidence of *A. catalaunalis* during

flowering and pod formation than the other crop stages. Similarly Egonyu, Kyamanywa [20] reported that the larvae spin silken webs around terminal leaves, eat the foliage and bore pods, causing substantial damage to grain inside. Lambda-cyhalothrin application significantly reduced the pest density in the early stage but not reduced at capsule damage[21].

With regard to *A. catalaunalis* infestation in the different locations *Kafta Humera* had the highest incidence (51.75%), while *Welkait* had the lowest (28.74%) during all growth stages of the plant (seedling, vegetative growth, flowering and capsule development stages) (Figure 5).




Figure 5. *A. catalaunalis* incidence at various growth stages and across the three districts in western Tigray.

### 3.4 *A. catalaunalis* Incidence and Damage in Sesame as Affected by Locations

The means and significant variations ( $p < 0.01$ ) of incidence and capsule damage caused by *A. catalaunalis* among districts and specific locations is illustrated in (Table 1). *A. catalaunalis* was observed in all surveyed fields of the three districts of sesame producing areas, however, higher incidence was recorded at *Kafta Humera* as compared to the other two districts. The average mean incidence over the districts varied from 54.14% at *Welkait* Woreda to 87.58% at *Kafta Humera*. A minimum of mean incidence 26.78% was recorded for the locality *Maidelie (tsegede)*, while the maximum mean incidence of 100% was recorded at *Rawian* and *Lugdi* localities of *kafta Humera*. The relatively higher incidence of *A. catalaunalis* in *Kafta Humera* could be due to the lower elevation (628m) and weather conditions of the area (higher temperature and low rainfall) as compared to the other two districts, which are located in high elevation (838.2 m) for *Welkait* and (764.3m) for *Tsegede* (Table 1). *Kafta Humera* has higher temperature and lower rainfall than *Welkait* and *Tsegede*. It is noticeable that the higher the altitude the cooler the weather condition and the lower the altitude the warmer the weather condition. Temperature decrease with altitude raise at a constant rate of  $-1^{\circ}\text{C}$  per 150 m up to tropopause [20] and the pest is in favor of warmer temperature. According to Ahirwar, Gupta [8], the level of incidence caused by the *A. catalaunalis* larvae is higher at higher temperature and dry weather conditions.

The overall pest incidence of 66% was recorded at fruiting stages of the crop in western zone

of Tigray. Therefore, the pest is economically very important in western Tigray. The economic threshold level for *A. catalaunalis* was reported as 10% of incidence [23] and in Ethiopia Mandefro, Girma [24] was stated that the economic injury level of the pest is 2-3% of infestation and SBN [25] has reported that control measure should start, when incidence of *A. catalaunalis* reached about 3 to 5 percent.

A minimum capsule damage of sesame caused by *A. catalaunalis* (7.11%) was noted in *Welkait Woreda*, while the highest (25.42%) was recorded in *Kafta Humera*. The districts of *Tsegede* and *Welkait* were similar in *A. catalaunalis* incidence (Table 1), but different in capsule damage, possibly because of the higher temperature for the low lands of *Tsegede*, which resulted in higher capsule damage next to *Kafta Humera*. Among the sesame fields surveyed (specific location) the minimum mean of severity/capsule damage (6.24%) was recorded in *Mogue* locality and the highest (28.87%) was recorded in *Rawian*. Sesame fields cultivated at an altitude below 700m, had the highest capsule damage as compared to other higher altitude areas (Table 1). Similar to the incidence, high severity of damage by *A. catalaunalis* in the lowlands was observed and may be because of higher temperature and lower rainfall. As altitude increases, temperature is getting decreased and this may have an impact on the duration and speed of *A. catalaunalis* reproduction, which clearly indicated that increment in altitude reduces reproduction rate of the pest.

Table 1 Means of incidence and capsule damage  $\pm$  (SD) of *A. catalaunalis* in western Tigray

| Districts           | Specific Locations | Altitude (m) | Incidence (%)                                  | Capsule Damage (%)                             |
|---------------------|--------------------|--------------|------------------------------------------------|------------------------------------------------|
| <b>Kafta Humera</b> | Rawian             | 592.8        | 100.00 <sup>c</sup> $\pm$ 0.0                  | 28.87 <sup>f</sup> $\pm$ 4.8                   |
|                     | Lugdi              | 621.2        | 100.00 <sup>c</sup> $\pm$ 0.0                  | 23.45 <sup>def</sup> $\pm$ 14.5                |
|                     | Mikadira           | 624.0        | 86.07 <sup>cde</sup> $\pm$ 17.3                | 28.77 <sup>ef</sup> $\pm$ 12.3                 |
|                     | Adebay             | 674.0        | 64.26 <sup>bc</sup> $\pm$ 11.9                 | 20.61 <sup>c-f</sup> $\pm$ 8.7                 |
|                     | <b>Mean</b>        | <b>628.0</b> | <b>87.58<sup>B</sup> <math>\pm</math> 17.8</b> | <b>25.42<sup>C</sup> <math>\pm</math> 10.4</b> |
| <b>Tsegede</b>      | Rubalemin          | 645.2        | 46.70 <sup>bc</sup> $\pm$ 17.2                 | 13.89 <sup>b-c</sup> $\pm$ 1.1                 |
|                     | Dansha             | 673.8        | 94.57 <sup>de</sup> $\pm$ 10.9                 | 10.17 <sup>abc</sup> $\pm$ 3.0                 |
|                     | Kebabo             | 729.0        | 54.53 <sup>b</sup> $\pm$ 18.6                  | 14.58 <sup>b-c</sup> $\pm$ 3.5                 |
|                     | Maydelie           | 1009.2       | 26.78 <sup>a</sup> $\pm$ 5.2                   | 10.99 <sup>a-d</sup> $\pm$ 4.5                 |
|                     | <b>Mean</b>        | <b>764.3</b> | <b>55.64<sup>A</sup> <math>\pm</math> 28.4</b> | <b>12.41<sup>B</sup> <math>\pm</math> 3.5</b>  |
| <b>Welkait</b>      | Maylemin           | 744.0        | 56.78 <sup>b</sup> $\pm$ 30.4                  | 9.15 <sup>ab</sup> $\pm$ 3.3                   |
|                     | Selam              | 816.0        | 33.12 <sup>a</sup> $\pm$ 20.9                  | 6.75 <sup>a</sup> $\pm$ 3.4                    |
|                     | Maygeba            | 859.8        | 59.51 <sup>bc</sup> $\pm$ 21.0                 | 6.31 <sup>a</sup> $\pm$ 1.9                    |
|                     | Mogue              | 933.0        | 67.14 <sup>bcd</sup> $\pm$ 8.7                 | 6.24 <sup>a</sup> $\pm$ 3.5                    |
|                     | <b>Mean</b>        | <b>838.2</b> | <b>54.14<sup>A</sup> <math>\pm</math> 23.4</b> | <b>7.11<sup>A</sup> <math>\pm</math> 3.0</b>   |
| <b>Grand Mean</b>   |                    | <b>743.5</b> | <b>65.79 <math>\pm</math> 27.8</b>             | <b>14.98 <math>\pm</math> 10.1</b>             |
| <b>CV(%)</b>        |                    |              | 13.8                                           | 17.3                                           |

Means followed by the same superscript are not statistically different from each other (at  $p < 0.01$ ), means followed by capital letters are means for *woreda*, means followed by small letters are for specific location (*tabia*)

### 3.5 Correlation of Incidence, Capsule Damage and Altitude

Capsule damage/severity was significantly related to incidence and altitude (Table 2).

Capsule damage was negatively and strongly correlated with altitude ( $r=-0.5719$ ) and positively with incidence ( $r=0.4704$ ). *A. catalaunalis* incidence correlated negatively and strongly with altitude ( $r=-0.5779$ ). Therefore, SWW incidence and altitude have the great contribution to the severity of sesame capsules. According to Reddy [26] incidence of *A. catalaunalis* correlated positively with temperature and sunshine hours and negative correlation with the rainfall.

Table 2 Correlation of incidence, capsule damage and altitude

|         | INC (%)   | CD (%)    | ALT(m) |
|---------|-----------|-----------|--------|
| INC (%) | -         |           |        |
| CD (%)  | 0.4704**  | -         |        |
| ALT(m)  | -0.5779** | -0.5719** | -      |

\*\* Indicates highly significant at 0.001% probability, INC=incidence, CD= capsule damage, ALT=altitude

#### 4. CONCLUSIONS

All surveyed fields of the three districts are found to be positive to *A. catalaunalis* in all growth stages of sesame. Late sown (end of July) sesame fields are totally infested by the pest. An increased trend of infestation is observed as altitude is decreasing. *A. catalaunalis* is economically very important pest in western Tigray. Therefore, planting of sesame should be on the onset of rainfall to minimize *A. catalaunalis* infestation and to boost sesame productivity. And further studies should be carried out on its biology, ecology, distribution across years and integrated managements.

#### REFERENCES

- [1] Khan, MHA, Sultana, NA, Islam, MN, & Hasan-uz-zaman, Mirza. (2009). Yield and yield contributing characters of sesame as affected by different management practices. *American Eurasian Journal of Scientific Research*, 4(3), 195-197.
- [2] Umar, HS, Okoye, CU, & Mamman, BD. (2010). Resource use efficiency in sesame (*Sesamum indicum* L.) production under organic and inorganic fertilizers applications in Keana Local Government Area, Nasarawa State, Nigeria. *Research Journal of Agriculture and Biological Sciences*, 6(4), 466-471.
- [3] FAOSTAT. (2012). Food and agriculture organization of the united nations statistical data. Retrieved 17, 2015, from <http://faostat.fao.org/faostat>
- [4] FAOSTAT. (2015). Food and agriculture organization of the united nations statistical data. Retrieved 22, 2016, from <http://faostat.fao.org/faostat>
- [5] Dennis, Weller. (2014). *Ethiopian Pulses, Oilseeds and Spices Processors-Exporters Association*. Paper presented at the Fourth International Conference on Pulses, Oilseeds, and Spices, Addi Ababa, Ethiopia.
- [6] Adefris, T., Tadele, A., & Tesfaye, B. . (2011). Sesame cultivation and use in Ethiopia.
- [7] Muez, Berhe, Berhanu, Abraha, Geremew, Terefe, & Walle, Melaku. (2008). Sesame harvest loss caused by sesame seed bug, *Elasmolomus sordidus* F. at Kafta-Humera

243 sesame fields.

244 *SINET: Ethiopian Journal of Science*, 31(2), 147-150.

245 [8] Ahirwar, M., Gupta, P., & Banerjee, S. (2010). Bio-ecology of leaf roller/capsule borer

246 *Antigastra catalaunalis* Duponchel. *Advances in Bio Research*, 1(2), 90-104.

247 [9] Selemun, Hagos. (2011). *Study on the potential of some botanical Powders and*

248 *Nimbecidine for the management of Sesame seed bug (Elasmolomus sordidus,*

249 *Fab.)(Hemiptera: Lygaeidae) in Humera, Northwest Ethiopia.* aau, Addis

250 Ababa, Ethiopia. 9

251 [10] Geremew, Terefe, Wakjira, Adugna, Muez, Berhe, & Hagos, Tadesse. (2012). Sesame

252 Production Manual. *EIAR/Embassy of the Kingdom of Netherlands, Ethiopia*, 49p.

253 [11] Egonyu, JP, Kyamanywa, S, Anyanga, W, & Ssekabembe, CK. (2005). *Review of pests*

254 *and diseases of sesame in Uganda.* Paper presented at the African Crop Science

255 Conference Proceedings.

256 [12] Suliman, E. H. , Bashir, N. H. H. , El tom, E., M.A. , , & Asad, Y. O. H. (2013).

257 Biology and Webbing behaviour of Sesame webworm, *Antigastra catalaunalis*

258 Duponchle (Lepidoptera: Pyraustidae). *Global Journal of Medicinal Plant Research*,

259 1(2), 210-213.

260 [13] Negash, Geleta Ayana. (2015). Status of Production and Marketing of Ethiopian Sesame

261 Seeds (*Sesamum indicum* L.): A Review. *Agricultural and Biological Sciences*

262 *Journal*,

263 1(5), 217-223.

264 [14] Karuppaiah, V. 2011. Evaluation of sesame genotypes for resistance to sesame leaf roller

265 and capsule borer, *Antigastra catalaunalis* (Duponchel) (Pyraustidae: Lepidoptera).

266 *Archives of Phytopathology and Plant Protection*, 44(9):882-887.

267 [15] Karuppaiah, V.& Nadarajan, L. 2013. Host plant resistance against sesame leaf webber

268 and capsule borer, *Antigastra catalaunalis* (Duponchel) (Pyraustidae: Lepidopter).

269 *African Journal of Agricultural Research*, 8(37): 4674-4680.

270 [16] Okonya, Joshua, Mwanga, Robert, Syndikus, Katja, & Kroschel, Jürgen. (2014). Insect

271 pests of sweetpotato in Uganda: farmers' perceptions of their importance and control

272 practices. *A SpringerOpen Journal*, 3(303), 1-10.

273 [17] Abdel-rahman, A. (2007). *Effect of sowing date on the performance of sesame*

274 *(Sesamum indicum L.) genotypes under irrigation conditions in northern Sudan.*

275 Paper presented at the African Crop Science Conference Proceedings.

276 [18] Abraham, E.V., Nataraja, K., & Ayaraj, S. (1990). *Investgaticns on the insecticidal*

277 *control of the phylldy disease of sesamun.* Paper presented at the Oil crops: Sesame

278 and sunflower subnetworks proceedings of the joint second workshop, Cairo, Egypt.

279 [19] Muzaffar, A., Rab-dino, K., Maqsood, A., & Imtiaz, A. (2002). insect pest associated

280 with sesame at Tanato Jam. *pakistan Journal of Applied Science* 2(7), 723-726.

281 [20] Egonyu, JP, Kyamanywa, S, & Ssekabembe, CK. (2009). Natural enemies of sesame

282 webworm and the effect of additive intercropping on its incidence in Uganda. *Journal*

283 *of Applied Biosciences*, 18, 1019-1025.

284 [21] Akinyemi, AO, Pitan, OOR, Osipitan, AA, & Adebisi, MA. (2015). Susceptibility of

285 sesame (*Sesamum indicum* L.) to major field insect pests as influenced by insecticide

286 application in a sub-humid environment. *African Entomology*, 23(1), 48-58. 10

287 [22] Cavcar, Mustafa. (2009). The International Standard Atmosphere. *Anadolu University,*

288 *Eskisehir, Turquia.*

289 [23] Kumar, Rakesh, Ali, Shamshad, & Dhoray, Umesh Chandra (2012). Incidence of

290 *Antigastra catalaunalis*, Dup. in Different Varieties of Sesame. *Molecular*

291 *Entomology*, 3.

- 292 [24] Mandefro, Nigussie, Girma, Anteneh, Anchala, Chimdo, & Kirub, Abebe. (2009).  
293 Improved technologies and resource management for Ethiopian Agriculture Training  
294 Manual.
- 295 [25] SBN. (2014). best agronomic practices for improvement of productivity and yield  
296 quality of sesame In SBN (Ed.), (pp. 1 -16). addis ababa, ethiopia SBN.
- 297 [26] Reddy, P.S., *Annual Report, 1995-96. Directorate of Oilseeds Research*, 1996:  
298 Hyderabad India.  
299

UNDER PEER REVIEW