

Original Research Article

Evaluation of Awir (*Ipomoea carnea*) for gully rehabilitation through different propagation techniques in north western zone of Tigray

Abstract

Since the past two decades, watershed management practices such as construction of stone bunds and establishment of exclosures have been widely implemented in the semi-arid low and highlands of northern Ethiopia to curb land degradation by soil erosion. The aim of this study was to identify scientific name and its impact of gully rehabilitation measure in the Serenta watershed of the Maitsebri agricultural research center experimental site, northern Ethiopia. Data were collected through semi-structured interviews, transect walks, field observation and field measurements to characterize the species. Scientific name of the species was identified and characterized as *Ipomoea carnea* Jac. by the national herbarium of Addis Ababa University. Based on the average result of the planted cuttings of stem/direct planting, its survival rate at side was greater than cuttings planted at top position of the gully. (100% at side and 95.84% at top). And for these cuttings planted by budding/layering, its survival rate at side and top positions are equal. (58.33% at side and 58.33% at top). This study demonstrated the potential of the species to bring remarkable restoration of eroded gully through reinforcement of gabion check dams that could serve as a basis for sustainable planning of future developments of areas experiencing severe land degradation due to water erosion and over grazing.

Keywords: *Ipomoea*, Gully, Propagation, Ethiopia

1. INTRODUCTION

Soil erosion caused by heavy rainfall and surface runoff is a serious problem in agricultural areas. Especially This is more serious on inclined slopes where the soil loss can reduce soil productivity and increase sediment and other pollution loads in receiving waters [1]. Soil erosion often occurs on steep slopes due to improper land use, monoculture and the use of tillage tools that leave the soil bare and pulverize it excessively [2]. After such treatments, the soil can be carried away by heavy rains. This problem affects crop productivity and the incomes of farmers. Soil erosion by runoff is often accepted as an unavoidable phenomenon associated with agriculture on sloping land. However, erosion removes the topsoil which is the part of the soil profile highest in

nutrients and organic matter [3]. The annual soil loss of Ethiopia is estimated to be 1.5 billion tons of soils ~~are eroded every per~~ year [4]. Due to the eroded top soil the productivity of the crop lands decreasing from time to time. One of the most wide spread threats to soil quality in this region is wind and water erosion that impacts the lives of the people in numerous ways. As a result of untreated catchment, poor vegetation covers and ~~sever~~ severe erosion have been forming gullies in the region. This formation of gullies leads to changing the productive land to marginal/unproductive/ land which in turns impacts the food security of the individual households since the size of land is small due to fragmentation. Using photogrammetric techniques, Shibru et al. (2003) cited in [5] to reverse the effect of gully erosion, the government of Tigray region has launched strategies which could improve the natural resources rehabilitation. Such as Among these strategies are catchment treatment, gully reclamation and other soil and water conservation structures. Recently introduced gully reshaping and revegetation efforts are found in various parts of the region with promising results. To be more effective, the physical soil and water structures should be integrated with biological measures such as, grasses, shrubs and trees to reinforce, strengthen and provides complete ground ~~cover~~ protection [6]. It is with this burning issue in mind that conserving soil and water through physical and biological measures are the current home work for the present generation to increase the productivity of the land for both short and long period of time.

However, very little or no research has been done so far as biological soil and water conservation tool. In addition, its name ~~is~~ was not identified and documented scientifically in Tigray in general and in the study area in particular. However, in Western Tigray, Tsegede woreda Zuria Dansha kebele, farmers used the species as small gully rehabilitation. The shrub is also useful for bee forage, fuel wood and planted as a fence for farmland.

Therefore, the objective of the study was to characterize and evaluate Awir as biological gully rehabilitation measure by determining the adaptability of environmental variables and morphological growth characteristics of different propagation techniques.

2. Materials and Methods

2.1. Study area description

The experiment was conducted at Medhani Alem kebele (the smallest political administration units in Ethiopia) Tselemti Woreda, North Western Tigray (Fig1). It is categorized as lowland and geographically located at 38°08' longitude and 13°05'

latitude with an altitude of 1350 m.a.s.l. Sorghum, maize, millet and *Tef* (*Eragrostis tef*) are the major crops. *Vegetations* such as onion, tomato, potato, pepper, carrot, garlic and groundnut grow in the study site. The area is also rich in *Boswellia papyrifera*, *Zizphus spina -Christ*, *Biospyros mespliformis*, *Cordia africana* and *Adansonia digitata* are the main vegetation types found in the area that serves as a source of income and utilizes for different purposes. Mai Tsebri is located 380 km from the regional city Mekelle. The altitudinal range of the worda falls between 800 and 2872 m.a.s.l. The land form is characterized as flat plain plateau, hills, mountains, gorge (valley) and some immediate break of slope.




Fig. 1: location map of the study area

The agro-climatic condition of the area is hot to dry semiarid lowland plains dictated by a very hot temperature. The maximum temperature ranges from 35.6°C in May to 36.4°C in April, while the minimum temperature is ranging from 15.7°C in December to 21.8°C in May (Tigray National Meteorological Agency (TNMA 2014)) (Fig. 2). The dry season occur between November to March whereas, the rainy season occurs between June to September, which follows a unimodal rainfall pattern.




Fig. 2. Six year rainfall and temperature of the study area (Source: TNMA, 2014)

Agriculture is the main livelihood of the community in the study area. It is known for its mixed farming system, where the rural people depend on crop and livestock production and in addition with agroforestry practices for their living.

2.2. Identification of Awir and Source of cuttings

It is a shrub species, which is found in western zone of Tigray. Traditionally local communities of the area have been planting Awir on their homestead lands to used to protect their lands from erosion by planting Awir on their homestead lands. Literatures were reviewed to identify and understand its agro-ecological distributions, management requirements and scientific nomenclature, but information was not found about the history of the species. In contrast, these areas have been covered with Awir resources that have not provide multi- service functions for to the local communities in particular and the region in general. So to utilize and manage the species properly as biological soil and water conservation and other socio-economic importance, its local and scientific name should be known and registered and Known. Hence, sample was taken to the national herbarium of Addis Ababa University for identification and characterization. The study was conducted through analysis of secondary data and supported using a participatory reconnaissance survey by employing different participatory tools, such as personal observation, transect walks, group discussions and focused group interviews. Representative peasant associations were selected for the actual study and then its main function, area converges and distribution of the species was recorded with the help of key informants and development agents.

Awir growing localities, namely Zuria Dansha and Baeker were selected for cutting collection based on agro-ecology similarity, representativeness, accessibility, distance and distance from the experimental site. This was done through consultation of local people and agricultural expertise by employing different tools of participatory reconnaissance

Formatted: Font: Italic

Comment [11]: What is the exact number you interviewed?

survey. And then branch cuttings were collected from altitudinal range of 1000 to 1350 m.a.s.l to minimize variations among sites and cuttings. Local communities were well recognized about the merits and demerits of the species. So that they recommend us to use as biological soil and water conservation techniques. Branch cuttings were taken from shrubs that look healthy at least phenotypically with a size range of 3 cm–5 cm in DBH and 2m–4 m in height. Then cuttings were collected, loaded and transported with a great care to avoid damages and scrapes to the planting site. As soon as arrived at the planting site, cuttings were trimmed to the required size using a sharp axe and planted in the prepared pits to protect cuttings against desiccation.

Comment [12]: This sentence is not clear

2.3. Experimental design and treatments

Site (gully) selection

The establishment of protective vegetation in a gully is usually much more difficult if runoff is not handled through the channel at the same time the plants are being planted. Therefore gullies that are treated with mechanical measures was selected for the satisfactory flood control and revegetation (Fig. 3). Our gully assessment finding revealed that the dimensions of the gullies were about two meters depth and 8m width leading to severe farm land deterioration [7].



Fig. 3. Active gully with free grazing problem of the experimental site.

2.4. Planting techniques and managements

Many different types of control measures for handling different kinds of erosion problems have been **done** over the last decades, even though effective conservation practices have not yet been developed for all numerous problems and combination of problems relating to the contrasting agricultural lands and practices. **The question is now, how and when Awir can plant to achieve the required objectives?** So the study is designed to answer these questions. Two propagation techniques (stem, layering) **and** planting position on the gully was used to investigate which type of planting technique and position is appropriate for successful revegetation.

Comment [13]: Do you mean.... can be planted to....

Cuttings of the species was collected from **Kafta Humera** where **Awir** found naturally exist. At each study experiment a total of 120 cuttings (stem **and** layering) was considered at side and top position of the gully. The cutting length of stem (1.5m) **and** layering (60cm) with a common branch girth was used in this experiment. There were 4 branch cuttings planted in double rows at spacing of 1m **between** plants and blocks when stem and layering propagation techniques used. The plot size was used 5m by 4m. Each branch cutting was planted in a pit that has 30-45cm depth and a cross-sectional area of 900cm² **(Personal communication to local farmers) in July, 2010**. It was placed within the fine soil of the pit and slanting lower cut will remain horizontal by keeping the cutting in an oblique position. Revegetation of gully is the plantation of the reshaped gully sides and bottom with biological species so that it reduces run off and control erosion and make the land productive [6]. Gully revegetation implies reaching a dense vegetation cover over the entire gully surface, i.e. both sides and bottom by planting Awir species. But for successful revegetation, three conditions and steps are important: Exclude the cattle throughout the year, use cut and carry and reshape the gully sides. Reshaping can be done either by cutting the edges or shaping the slope in to series of small steps or micro benches every 0.75-1m distance (Fig. 4).



Fig. 4: Gabion check dam construction and planted branch cuttings by direct and layering on reshaped gully

Source: (Photo taken by Girmay D. in 21/07/2010)

2.5. Data collection

To evaluate its growth rate 30% samples were taken at the month of 1, 2, 3, 6 & 12 from the total cuttings planted and then data on total shoot number, sample shoot length, total

leaf number and sample leaf length and survival rate of the cuttings were collected periodically.

2.6. Data analysis

Descriptive analysis such as mean of data was conducted on survival rates and planting methods on different growth response of morphological characteristics of planted branch cuttings and supported by photo evidences to show the effect of the species on sediment deposition and gully rehabilitation of the experiment.

3. Result and Discussion

3.1. Uses and characterization of Awir

It is an exotic shrub species, originally came from Sudan. ~~local~~Local communities of the area have been used to protect their lands from erosion and live fence by planting on their homestead lands. The socioeconomic analysis reveals that the species has two local names; *Awir* and *Serka-abeba* (meaning flowering all year round). Scientific name of the species was identified and characterized as *Ipomoea carnea* Jac. by the national herbarium of Addis Ababa University. Respondents under lined that, the main function of Awir is protecting land from erosion, live fence, increase soil fertility by reducing soil erosion, honey bee flora and can be used as shade for wild life if planted at protected areas (Fig. 5). Overall, we found a robust indication of positive effects of Awir on economic conditions of the households of the study area by protecting the small land from soil erosion. This is in line with the findings of Gebramlak et al [⁸] who found that lowland bamboo positively affected the income of rural households in Serako, Tigray, Ethiopia. In respondent's point of view, it is important to scale up this as alternative technology at areas that have gully and grazing problems. The species does not need much management but if we plant at farm lands, pruning will be important to decrease its holding area coverage. it has the ability to tolerate moisture stress and grow at poor soil fertility and water logging areas.

Comment [14]: This is not clear



Honey bee flora

Irrigation site is protected from flood damage



Gully in homesteads are protected by Awir

Fig. 5. The different use of Awir by the local communities.

3.2. Propagation techniques and its effect on gully reclamation

After cuttings were planted, it started to give new shoots and leaves within 5-7 days. Based on the collected data for stem/Budd planting at both side and top positions of the gully, the average result of planted cuttings characteristics was are provided-presented in Table 1.

Table 1: descriptive analysis of branch cuttings

S.N	Planting method	shoot number	sample shoot length	sample leaf length	sample crown diameter	Survival rate
1	Stem	18.17	207.78cm	22.45cm	2.92m	97.92%
2	Bud	6.83	160.11cm	22.00 cm	1.47m	58.33

209 Based on the average result of the planted cuttings of stem/direct planting, its survival
210 rate at side was greater than cuttings planted at top position of the gully; **100% at side and**
211 **95.84% at top position of the gully.** And for these cuttings planted by budding/layering,
212 its survival rate at side and top positions are equal. (58.33% at side and 58.33% at top).
213 For budding propagation method of Awir, the survival percent is decreasing due to huge
214 flooding (side) and water logging (top) effect. **This study is in line with the findings of**
215 **Reubens et al [⁶] who reported that Mortality rate for Dodonaea tree seedlings planted at**
216 **gully flour was especially high during the summer rains.** Hence, when we see almost all
217 the morphological growth characteristics of planted cuttings in stem/direct planting,
218 promising result has shown than cuttings planted by budding/layering propagation
219 method. Different and relevant stakeholders such as individual farmers, extension
220 workers, researchers, the media and other concerned individuals and professionals were
221 invited to visit the observation trial at different occasions. All the visitors and/or the
222 respondents **gave their views** on the newly vegetative propagation technology **and**
223 underlined to scaling up for other **similar** areas that have gully problems.

224 | In addition to the **regularly**-measured plant characteristics, it was also observed how **Awir**
225 cuttings planted were capable to re-sprout. In line to this finding many studies conclude
226 trees or shrubs that have good re-sprout behavior is good in rehabilitation of degraded
227 gullies [^{4, 6}].

Comment [15]: Some are in bold and italic, some simple word, and other only in italic. Please harmonize.



228

229 | [\(Source: Photo taken by Girmay D. in 21/01/2011\)](#)



Fig. 6. Showing by arrow large tree on the upper catchment to see the effect of the biological species on gully rehabilitation over time.

Source: (Photo taken by Girmay D. in 21/07/2011)

A study conducted by Gebreegziabher [⁹] in Ethiopia revealed that runoff volume is significantly ($P < 0.05$) reduced due to permanent bed planting system as compared to the *Terwah*¹ system and conventional ploughing.

¹ Terwah ploughing: The same tillage operations as described for the traditional system were applied but additionally contour furrows were made at 1.5–2m interval after the third superficial ploughing.

This demonstrates that not only biological soil and water conservation practices but also ploughing techniques can reduce runoff and soil erosion.

4. Conclusion and recommendation

The results suggest that *Ipomoea carnea* Jac., were the most suitable for biological soil and water conservation in active gully rehabilitation where free grazing is the main problem. The study concludes that farmers should select the species due to high performance to adapt to the harsh environmental conditions, easily established, locally available and a lower cost that bring better gully rehabilitation and environmental conservation. The role of the species in sediment trap performance and ecological effect should be studied.

Acknowledgment

The authors would like to acknowledge the generous financial support by Maitsebi agricultural research center to perform the study. Many thanks go to the three anonymous reviewers for their valuable comments on an earlier version of this paper.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

5. References

1. Gebremeskel, G., Gebremicael, T. G. & Girmay, A. Economic and environmental rehabilitation through soil and water conservation , the case of Tigray in northern Ethiopia. *J. Arid Environ.* 1–12 (2017). doi:10.1016/j.jaridenv.2017.12.002
2. Donjadee, S. & Tingsanchali, T. Soil and water conservation on steep slopes by mulching using rice straw and vetiver grass clippings. *Agric. Nat. Resour.* **50**, 75–79 (2016).
3. Fenta, A. A., Yasuda, H., Shimizu, K., Haregeweyn, N. & Negussie, A. Dynamics of Soil Erosion as Influenced by Watershed Management Practices : A Case Study of the Agula Watershed in the Semi-Arid Highlands of Highlands of Northern Ethiopia. *Environ. Manage.* **58**, 889–905 (2016).
4. Taha, N., Behailu, M. & Alemayhu, F. *Impact of Watershed Management in*

- 268 *controlling Desertification in Atsbi district of northern Ethiopia. (2006).*
- 269 5. Nyssen, J. *et al.* The effectiveness of loose rock check dams for gully control in
270 Tigray, northern Ethiopia. *Soil use Manag.* 55–64 (2004).
271 doi:10.1079/SUM2003223
- 272 6. Reubens, B. *et al.* Establishment and management of woody seedlings in gullies in
273 a semi-arid environment Tigray , Ethiopia. **324**, 131–156 (2009).
- 274 7. Bekele, Y., Tadesse, N. & Konka, B. Preliminary Study on the Impact of Water
275 Quality and Irrigation Practices on Soil Salinity and Crop Production, Gergera
276 Watershed, Atsbi-Wonberta, Tigray, Northern. *Mekelle Univ. Dep. Earth Sci. Coll.*
277 *Nat. Comput. Sci.* **4**, 29–46 (2012).
- 278 8. Gebramlak, G. D., Abadi, N. & Hizikias, E. B. Socioeconomic contribution of
279 oxytenanthera abyssinica (A.Rich) munro and determinants of growing in
280 homestead agroforestry system in Northern Ethiopia. *Ethnobot. Res. Appl.* **14**,
281 479–490 (2016).
- 282 9. Gebreegziabher, T. *et al.* Soil & Tillage Research Contour furrows for in situ soil
283 and water conservation , Tigray , Northern Ethiopia. **103**, 257–264 (2009).

284