

# Assessment of Richness of Diversity between Various Species of vegetation type and their Threat status in Mukundpur Forest Area, Satna District, Madhya Pradesh, India

P.K. Singh \*, Prachi Singh \*\* and Sherendra Sahu \*\*\*

\*Madhya Pradesh Forest Department, Satpura Bhavan, Bhopal (MP)-462004

\*\*Department of Botany, Govt. Girls PG College, Rewa, (MP)-486001

\*\*\*Department of Biotechnology, A.P.S. University, Rewa (MP)-486220

**ABSTRACT:** Mukundpur forest range is situated in Amarpatan Tahsil in Satna district of Madhya Pradesh India. The species of 58 trees, 28 shrubs, 08 lianas, 20 herbs and 19 grasses (total 133) had been found by using vegetation sampling. By evaluating IVI (Important Value Index) for the species of various vegetation types, the threat and conservation status was assessed by Normal Distribution Principle. The richness of species of study area was assessed by taking the two parameters i.e. number of species and their average IVI between various vegetation types and threat and conservation categories. The result of richness of diversity in numbers and their IVI for different vegetation types were expressed in terms of significant or non significant. The present study provided the current status of species diversity current diversity and it also designing procedure for optimal species diversity in study area by developing various alternative strategies to assess the number of species and their IVI between various vegetation types with optimum species diversity and minimizing the threat parameters simultaneously. Study area was the forest area of 111.55 km<sup>2</sup> of Mukundpur range of Satna Forest division, Madhya Pradesh, India. Field work was carried out during October 2015 to January 2016.

The study about the current status reveals that there was a nonsignificant richness of species diversity between various vegetation types and threat and conservation categories when number of the species was considered as a parameter. The species diversity was in the significant state of threat and conservation categories, but same was insignificant between various vegetation types when average IVI of the species was considered as a other parameter. The optimization technique was used to find out the number and average IVI of the species between various vegetation types (Trees, Shrubs, Lianas, Herbs and Grasses) by making the variance significant. The same technique is used to reduce the threat status between various threat and conservation categories by making the variance of number and average IVI of the species as non significant.

The optimal number species of 58 tree, 28 shrubs, 30 lianas, 20 herbs, 19 grasses (Total 155) and optimal average IVI between threat and conservation category 1 were assessed as 23.073, 19.284, 9.085, 25.321 and 10.067 within trees, shrubs, lianas, herbs and grasses respectively with total of 86.83 were assessed to make the significant diversity and at the same time maintaining the non significant status of threat and conservation status. The number of lianas species should be increased from 8 to 30 which lowered the average IVI of the species from 44.125 to the average value of 9.085.

**Keywords:** Analysis of Variance, Frequency Class, IVI, Level of Significance, Normal Distribution Principle, Optimization Technique, Significant and Non Significant, Threat and Conservation Categories.

## 1. INTRODUCTION

The study and analysis has been carried out with the purpose of conservation, propagation of diversity of vegetation with an innovative methodology employing modern statistical tool, for more meticulous and precise information on species diversity in terms of number and IVI with respect to different kinds of vegetation. The relevance of the study provides method of ecological modeling and procedure for designing species diversity in a forest area. In designing procedure, the various alternative strategies are developed to find the number of species between various vegetation types to make the species diversity optimum in the study area and at the same time minimizing the threat parameters. This will help in future research on the vegetation and allied conservation measures of various species with higher degree of precision

and accuracy. In the study area the present status of species diversity in terms of number and average IVI had been assessed in the study area. After evaluating the present status of species diversity, the optimal number of species and average IVI was evaluated to optimize species diversity and minimizing threat and conservation status in study area.

There are two popular diversity indices [1, 2]. These indices do not represent the species which are responsible threatening status in particular ecosystem. About 42% of world forests are dry forest [3] at global level. At national level, India accounts for 8% of the global biodiversity with only 2.4% of the total land area & the world [4, 5]. The tropical dry forests occupy 38% of the total forest area in India [6]. Tropical forests are often referred to as one of the most species diverse terrestrial ecosystems [7]. An Assessment of threatened plants of India has been made by [8]. Conservation and economic evaluation of biodiversity has been done by [9]. At regional level, threat assessment of Vindhyan region of Madhya Pradesh has been made by [10, 11, 12, 13, 14].

Mukundpur Range situated in Amarpatan Tahsil of Satna district in Madhya Pradesh, India. The first white Tiger safari is established at this village. The Mukundpur range is surrounded by mining areas of Bauxite and Limestone. The nearby located cement factories are always in search of new areas, besides exploiting existing known areas. Thus area is encountering impact of temperature rise, industrialization, desertification, shifting in the growing seasons of plants, loss of pollinators and seed dispersers, causing extinction of precious plants.


Looking towards the ecological stress in study area, the threat and conservation status between various vegetation types trees [15], shrubs [16], lianas [17], herbs [18] and grasses [19] had done in study area.

In the present work richness of plant diversity is assessed by the Analysis of Variance (ANOVA) between various species of vegetation types and their threat categories in terms of kinds of species and their IVI. This work also discusses the various alternative strategies to find the number of species between various vegetation types to make the species diversity optimum in the study area at the same time reducing the threat parameters which is providing a procedure for ecological modeling. This study will help in future research on the vegetation and allied conservation measures of various species with higher degree of precision and accuracy.

## **2. STUDY AREA:**

Mukundpur region mainly comprises the present area of Mukundpur range of Satna forest division and lies between north latitude of  $24^{\circ}11'35''$  N to  $24^{\circ}26'25''$  N and longitude of  $81^{\circ}6'35''$  E to  $81^{\circ}22'20''$  E The map of the study area is shown in **figure 1**. The range has geographical area of  $589.71 \text{ km}^2$  with forest area  $111.55 \text{ km}^2$  as discussed by Singh, 2018. Northern boundary lies with Beehar River demarcating Satna and Rewa district. The forest of Mand reserve is situated in this area where first white tiger safari is established. Eastern boundary lies mainly with the district boundaries bifurcating Rewa and Satna districts. The famous Charaki ghati forms one of its boundaries. Southern boundary lies mainly with submerged area of Son River and it extends to district boundaries of Shahadol and Satna districts. The average annual rainfall in study area was noticed from 354.1 mm to 1748.4 mm with mean annual rainfall of 1074.26 mm.

Figure 1- Map of the Study Area



The area receives nearly 51 rainy days in year. South western monsoon plays the active role of precipitation in study area starting from middle of June month. The average highest daily temperature ranges within 24.06<sup>0</sup> C to 41.73<sup>0</sup> C with mean temperature of 32.24<sup>0</sup> C. The highest daily temperature recorded was 47.7<sup>0</sup> C. Similarly the average lowest daily temperature was 8.85<sup>0</sup> C to 27.72<sup>0</sup> C with minimum daily temperature of 1.7<sup>0</sup> C.

### 3. MATERIAL AND METHOD:


For the assessment of biodiversity of Mukundpur region, the vegetation sampling was done for the trees, shrubs, herbs, lianas and grasses. Stratified systematic random sampling method was used for sampling for the vegetation Anon [26]. For determining minimum number of sample points, the formula used is  $n = z^2 \frac{pq}{E^2}$  where E= difference between population proportion mean and sample proportion average, p = population proportion, q= 1- p, z=1.96 for a level of significance of 95% [21].

Based on the secondary data from Mukundpur range and Satna forest division, the sample size for various tree parameters i.e. number of trees per hectare, volume of trees per hectare and established regeneration per hectare was calculated at 10% error (E) between population and sample proportion at 95% level of significance keeping in view time and other resources [22].

Minimum 95 numbers of sample points were calculated from the above formula to assess the vegetation. The forest maps of Mukundpur range on survey of India topo sheet is of the scale of 1:15000. The grids at 35''x 35'' and 30''x30'' intervals are drawn by trial and error, for systematic random sampling. The 111 and 151 random points were recorded on above grid. The 151 sample points at 30''x 30'' were selected on safer side, so that points may fall in river bed, submergence and encroachments. The longitudes and latitudes of 151 points were noted from topo sheets and list of 151 points are prepared.

Each sample points were located on ground with the help of GPS.


**Figure 2- sample plot with their Quadrat**



At each sample points, the layout of sample plot of 0.16 hectare with 9 quadrat of 2 ×2 m on ground as shown in **figure 2** was done with the help of prismatic compass Anon [20]. The combined map with sample plot in the study area is presented figure 3.



Figure 3- Combined Map of the Study Area with sample plot




At these points recording of data of the girth and species of the trees, along with species of shrubs and lianas (numbers) were taken on whole sample plot of 0.16 hectare and data for species of herbs, grasses and established regeneration was recorded at each 9 quadrat of  $2 \times 2$  m. The results were analyzed by developing a Microsoft access program to calculate the number, regeneration of trees per hectare and volume in  $m^3$  per hectare by using local volume table, prepared for Satna forest division. The calculations for the density, frequency, basal area and IVI of the all species of trees have been done with same program [23]. The name and number of shrubs, lianas herb plants and established regeneration of plant species was also evaluated. For grasses, only the names of the species and their presence were recorded, in each quadrat. All the IVI for all the vegetation type species have been summarized in decreasing order and analyzed further to assess the conservation and protection status of species by using the NORMAL DISTRIBUTION PRINCIPLE [24] as represented in **figure 4**, normal distribution curve of IVI values. Principle is as under:

$\mu$  = mean of IVI of all species,  $\sigma$  = standard deviation of IVI,

Then normal distribution principal states that there should be:-

- (a) 68% of total number of species whose IVI is between  $\mu + \sigma$  and  $\mu - \sigma$ .
- (b) 95% of the total number of species whose IVI is between  $\mu + 1.96\sigma$  and  $\mu - 1.96\sigma$ .
- (c) 99% of the total number of species whose IVI is between  $\mu + 2.58\sigma$  and  $\mu - 2.58\sigma$ .

Figure 4- Normal distribution curve of IVI values



For safer evaluation for IVI, for conditions (b) and (c)  $\mu - 2\sigma$  to  $\mu + 2\sigma$  and  $\mu - 3\sigma$  to  $\mu + 3\sigma$  have been calculated and used in further study. Now again here,  $\mu$  is the population mean and is equivalent to sample average and  $\sigma$  is population standard deviation and here for sample it is replaced by  $\sigma/\sqrt{n}$  i.e. standard error(SE).

Now, with the help of this principle [25], categorization is done as follows:

$IVI < \mu - 3\sigma$  (species having IVI less than 1%) - category 4.

$\mu - 3\sigma \leq IVI < \mu - 2\sigma$  (species having IVI between 1 to 5%) - category 3.

$\mu - 2\sigma \leq IVI < \mu - \sigma$  (species having IVI between 5 to 32%) - category 2.

$IVI \geq \mu - \sigma$  (species having IVI greater than 32%) - category 1.

The species in category 4 require highest degree of protection. The species in category 3 require lesser protection than category 4. The species in category 2 require lesser protection than category 3. The species in category 1 require least protection and are available in plenty and they are available for harvesting. The species wise results for trees [15], shrubs [16], lianas [17], herbs [18] and grasses [19] have been discussed for various threat and conservation categories for the study area.

After analyzing the above results have been tabulated between numbers of various species and their average IVI of different vegetation types and their number of species and average IVI for conservation and threat categories. After the tabulation the Null and alternate hypothesis have been formulated at 5% level of significance for the variance of number of different species and variance of average IVI between various vegetation types and their threat and conservation categories.

The two way Analysis of Variance (ANOVA) has been done by using 'F statistics' at 5% level of significance. At 5% level of significance, the testing of hypothesis has been done by following decision rules:

1. If  $F_{\text{calculated}} < F_{\text{tabulated}}$ , Null hypothesis is not rejected. It means there is no significant difference in variance of number of species and average IVI between various vegetation types and between various threat and conservation categories.
2. If  $F_{\text{calculated}} > F_{\text{tabulated}}$ , Null hypothesis rejected. It means there is a significant difference in variance of number of species and average IVI between various vegetation types and between various threat and conservation categories.

After testing the significance, our objective is to optimize the variance in number and their average IVI of the species between various vegetation types and tries to reduce the same between various threat and conservation categories. The optimization technique is used to find out the number an average IVI of the species between various vegetation types (Trees, Shrubs, Lianas,

Herbs and Grasses) by making the variance significant. The same technique is used to reduce the threat status between various threat and conservation categories by making the variance of number and average IVI of the species as non significant.

The optimization technique, [26] uses the iterative processes which consist of first designing a basic feasible solution and proceed towards OPTIMAL SOLUTION and testing the each feasible solution for optimality to know whether the solution on hand is optimal or not. If not an optimal, redesign the program and test for optimality until the test confirms the OPTIMALITY. The iterative steps are repeated until a finite optimal solution, if exists, is found.

#### **4. RESULT AND DISCUSSION:**

##### **4.1 The results for the 58 species of trees, for various threat and conservation categories are as follows:-**

**1. Category 4** No species exists in this category.

**2. Category 3** The species in this category are *Bridelia squamosa*, *Holoptelia integrifolia*, *Bombax ceiba*, *Bauhinia racemosa*, *Mitragyna parvifolia*, *Albizia procera*, *Sterculia urens*, *Carissa opaca*, *Ficus benghalensis*, *Solanum amricanum*, *Pterocarpus marsupium*, *Dalbergia sissoo*, *Boswellia serrata*, *Ziziphus mauritiana*, *Ficus religiosa*, *Schleichera oleosa*, *Dendrocalamus strictus*, *Ficus benjamina*, *Annona squamosa*, *Acacia leucophloea*, *Grewia tilaefolia*, *Gardenia latifolia*, *Woodfordia fruticosa*, *Careya arborea*, *Lannea coromandalica*, *Semecarpus anacardium*, *Gardenia resinifera*, *Terminalia arjuna*, *Ixora arborea*, *Vitex negundo*, *Cordia macleodii*, *Acacia ferruginea*, *Kydia calycina* (33 species with average IVI of 0.531). These species requires improvement in presence, density and basal area as a management strategy.

**3. Category 2** The species in this category are *Miliusa tomentosa*, *Shorea robusta*, *Phyllanthus emblica*, *Azadirachta indica*, *Holarrhena pubescens*, *Ziziphus xylopyrus*, *Albizia odoratissima*, *Cassia fistula*, *Cassine glauca*, *Terminalia alata*, *Aegle marmelos*, *Feronia elephantum*, *Garuga pinnata*, *Terminalia bellirica* and *Ougeinia oojeinensis* (14 species with average IVI of 2.039).

**4. Category 1** The species in this category are *Diospyros melanaxylon*, *Lagerstroemia parviflora*, *Tectona grandis*, *Butea monosperma*, *Anogeissus latifolia*, *Wrightia tinctoria*, *Ailanthus excelsa*, *Strychnos potatorum*, *Buchanania lanzan*, *Acacia catechu* and *Madhuca longifolia* (11 species with average IVI of 23.073).

##### **4.2 The results for 28 species of shrubs, for various threat and conservation categories are as follows:-**

**1. Category 4** No species exists in this category.

**2. Category 3** There are 11 species with average IVI of 0.557. These are *Grewia tilifolia*, *catunaregam spinosa*, *Nyctanthes arbor-tristis*, *Terminalia arjuna*, *Leucaena*



*leucocephala*, *Bridelia squamosa*, *Anogeissus pendula*, *Jatropha curcas*, *Abrus precatorius*, *Buchanania lanzan* and *Terminalia bellirica*.

3. **Category 2** The 8 species are observed with average IVI of 2.536. These species are *Aegle marmelos*, *Ziziphus xylopyrus*, *Ziziphus mauritiana*, *Cassia fistula*, *Artemisia vulgaris*, *Feronia elephantum*, *Miliusa tomentosa* and *Annona squamosa*.
4. **Category 1** The 9 species are observed with average IVI of 19.284. The species are *Lantana camara*, *Helicteres isora*, *Carissa opaca*, *Dendrocalamus strictus*, *Holarrhina pubescens*, *Woodfordia fruticosa*, *Chloroxylon swietenia*, *Alanium* spp, and *Solanum nigrum*.

Some of the species of trees are also appearing as a shrub in the results.

#### **4.3 The results for 08 species of lianas, for various threat and conservation categories are as follows:-**

1. **Category 4** There are 3 species with average IVI of 2.69 are under this category. They are *Bauhinia vahlii*, *Bauhinia purpurea* and *Butea superba*.
2. **Category 3** Only one species with average IVI of 15.5 is found in this category i.e. *Asparagus racemosus*.
3. **Category 2** No species are found in this category.
4. **Category 1** There are 4 species with average IVI of 44.125 in this category. These are *Hemidesmus indicus*, *Acacia donaldi*, *Clitoria ternatea* and *Cocculus hirsutus*.

#### **4.4 The results for 20 species of herbs, for various threat and conservation categories are as follows:-**

1. **Category 4** There are no species under this category.
2. **Category 3** There are 5 species with average IVI of 0.414 are classified under this category. These are *Mimosa pudica*, *Sida veronicaefolia*, *Sida cordifolia*, *Solanum virginianum* and *Rauvolfia serpentina*.
3. **Category 2** There are 8 species with average IVI of 2.583 are found in this category. These are *Cyperus rotundus*, *Vigna trilobata*, *Tribulus terrestris*, *Coleus barbatus*, *Andrographis paniculata*, *Enicostemma littorale*, *Coix lacryma jobi* and *Gymnema sylvestre*. These species require more protection than category 1 species, though they have little threat to extinct.
4. **Category 1** There are 7 species with average IVI of 25.321 in this category. These species are *Ocimum basilicum*, *Convolvulus microphyllus*, *Phyllanthus amarus*, *Aconitum chasmanthum*, *Eclipta alba*, *Borreria articularis* and *Bacopa monnieri*.

**4.5 The results for 19 species of grasses for various threat and conservation categories are as follows:-**

1. **Category 4** There are 6 species with average IVI of 0.12 under this category which require highest degree of protection. These species are *Dichanthium annulatum*, *Vigna trilobata*, *Paspalidium punctatum*, *Peucedanum dhana*, *Grewia hirsuta* and *Ziziphus oenoplia*.
2. **Category 3** There are 3 species with average IVI of 1.75 in this category. These are *Setaria intermedia*, *Ipomea reniformis* and *Saccharum spontaneum*.
3. **Category 2** There are only one species with average IVI of 3.42 are found in this category i.e. *Eragrostis tenella*.
4. **Category 1** There are 9 species with average IVI of 10.067 in this category. These are *Heteropogon contortus*, *Cenchrus ciliaris*, *Euphorbia thymifolia*, *Aristida funiculata*, *Cynodon dactylon*, *Oxalis stricta*, *Paspalum notatum*, *Dichanthium annulatum* and *Cassia tora*.

**4.6 Present status of species diversity in study area in terms of Number of kinds of species:-**

The results for number of species in various vegetation types ( $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$ ) and their threat and conservation categories ( $C_1$ ,  $C_2$ ,  $C_3$  and  $C_4$ ) From 4.1, 4.2, 4.3, 4.4 and 4.5 the number of various trees, shrubs, lianas, herbs and grasses with their threat and conservation status is presented in table 1.

**Table 1:- Number of species in various vegetation types and their threat and conservation categories**

Vegetation type	Threat and conservation category				Total
	category 1 ( $C_1$ )	category 2 ( $C_2$ )	category 3 ( $C_3$ )	category 4 ( $C_4$ )	
Trees ( $R_1$ )	11	14	33	0	58
Shrubs ( $R_2$ )	9	8	11	0	28
Lianas ( $R_3$ )	4	0	1	3	8
herbs ( $R_4$ )	7	8	5	0	20
Grasses ( $R_5$ )	9	1	3	6	19
Total	40	31	53	9	133 (Grand Total)

**For table 1 the following hypothesis at 5% level of significant is formulated:**

**A. Between conservation and threat categories:**

1. **Null hypothesis ( $H_0$ ):** There is no significant difference in the variance of number of species.
2. **Alternate hypothesis ( $H_i$ ):** There is significant difference in the variance of number of species.

## B. Between various vegetation types:

1. **Null hypothesis ( $H_0$ ):** There is no significant difference in variance of number of species.
2. **Alternate hypothesis ( $H_1$ ):** There is significant difference in variance of number of species.

The results of two ways analysis for table 1 is presented below in **table 2**:

**Table 2 :- ANOVA table for table 1**

Source of Variation	Sum of squares	Degree of freedom	Mean Sum of Square	F <sub>calculated</sub>	F <sub>tabulated at 5% level of significance</sub>
Between conservation category	205.75	4 - 1 = 3	68.583	1.601172	F <sub>(3,12)</sub> = 3.49
Between vegetation type	358.8	5 - 1 = 4	89.7	2.09418	F <sub>(4,12)</sub> = 3.26
Residuals	514	19 - 7 = 12	42.833		
Total	1078.55	20 - 1 = 19			

For variance of the number of species between conservation and threat categories  $F_{\text{calculated}} < F_{\text{tabulated}}$ , null hypothesis is accepted. It means there is no significant difference in variance of the number of species between threat and conservation category.

For variance of the number of species between various vegetation type  $F_{\text{calculated}} < F_{\text{tabulated}}$ , null hypothesis is accepted. It means there is no significant difference in variance of the number of species between various vegetation types.

At present in the study area above results indicate that variance of the number of species between various vegetation types (trees, shrubs, lianas, herbs and grasses) and between threat and conservation categories are not significant at 5% level of significance. Regarding comparing the results from other literatures suggest that the systems requires urgent development of programmes for biodiversity assessment and monitoring [27, 28], but standard methods are missing. The area is dominated by majority of tree species which are restricting the growth of shrubs, lianas, herbs, grasses and tubers. The species of *Lantana camara* and *Ocimum basilicum* are invading the study area. These species are appearing as weeds which are checking the growth of other species in the area. The various tree species are not converting into trees due to excessive invasion of *Lantana camara* as majority of species remaining in bushy form. The invasion of *Ocimum basilicum* as a weed, is limiting the growth of herbs and grass species.

#### 4.7 Present status of species diversity in study area in terms of Number of kinds of species

The results for average IVI of species in various vegetation types ( $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$ ) and their threat and conservation categories ( $C_1$ ,  $C_2$ ,  $C_3$  and  $C_4$ ) From 4.1, 4.2, 4.3, 4.4 and 4.5 the IVI of various trees, shrubs, lianas, herbs and grasses with their threat and conservation status is presented in table 3.

**Table 3: - Average IVI of species in various vegetation types and their threat and conservation categories**

Vegetation type	Threat and conservation category for Average IVI				
	category 1 ( $C_1$ )	category 2 ( $C_2$ )	category 3 ( $C_3$ )	category 4 ( $C_4$ )	Total
Trees ( $R_1$ )	23.073	2.039	0.531	0	25.643
Shrubs ( $R_2$ )	19.284	2.536	0.557	0	22.377
Lianas ( $R_3$ )	44.125	0	15.5	2.69	62.315
herbs ( $R_4$ )	25.321	2.583	0.414	0	28.318
Grasses ( $R_5$ )	10.067	3.42	1.75	0.12	15.357
Total	121.87	10.578	18.752	2.81	154.01

For table 2 the following hypothesis at 5% level of significant is formulated:

**A. Between conservation and threat categories:**

1. **Null hypothesis ( $H_0$ ):** There is no significant difference in variance of the average IVI of the species.
2. **Alternate hypothesis ( $H_i$ ):** There is significant difference in variance of the average IVI of the species.

**B. Between vegetation types:**

1. **Null hypothesis ( $H_0$ ):** There is no significant difference in variance of the average IVI of the species.
2. **Alternate hypothesis ( $H_i$ ):** There is significant difference in variance of the average IVI of the species.

The results of two ways analysis is presented below in table 4:

**Table 4 :- ANOVA for table 3**

Source of Variation	Sum of squares	Degree of freedom	Mean Sum of Square	F calculated	F tabulated at 5% level of significance
Between conservation category	1878.795	4 - 1 = 3	626.265	15.807	$F_{(3,12)} = 3.49$
Between vegetation type	333.85	5 - 1 = 4	83.463	2.107	$F_{(4,12)} = 3.26$
Residuals	475.438	19 - 7 = 12	39.620		
Total	2688.083	20 - 1 = 19			

For variance of the average IVI of species between conservation and threat categories  $F_{\text{calculated}} > F_{\text{tabulated}}$ , null hypothesis is rejected. It means there is a significant difference in variance of the average IVI of species between threat and conservation category.

For variance of the average IVI of species between various vegetation type  $F_{\text{calculated}} < F_{\text{tabulated}}$ , null hypothesis is accepted. It means there is no significant difference in variance of the average IVI of species between various vegetation types.

The present status of species diversity in study area, when the variance of average IVI between various vegetation types is taken, there is no richness of species diversity. But there is a significant difference in variance of average IVI of the species between the threat and conservation categories. Hence average IVI of the species is an important parameter as IVI of particular species includes the frequency, density and dominance in itself. Thus, the study area does not show the richness in species diversity but it shows the high stress in threat category. Regarding comparing these results from other literatures much information are not available but Chiarucci [29] states that no reliable method yet exists for estimating species richness in an area. Many authors have suggested diversity indices appropriate for their own studies, no one of which can be considered a priori correct for general application [30, 31, 32, 33, 34]. To improve the richness in species diversity the variance in the average IVI of threat category should be reduced and variance in the average IVI between various vegetation types should be increased. For that the different types of species and their population and frequency and basal area should be increase by management intervention in the study area.

#### 4.8 Evaluation of optimum species diversity in study area in terms of Number of kinds of species with minimizing the threat and conservation status

In order to optimize the species diversity in number of kinds of species and at the same time minimizing the threat and conservation status  $F_{\text{calculated}} > 3.26$ , say  $F_{\text{calculated}} = 3.3$ . For making the significant diversity in number of species ANOVA **table 2** is redesigned by backward calculation for mean sum of squares and total sum of squares between various vegetation types and it is expressed in bold and small brackets and modified ANOVA table is presented in **table 5**.

**Table 5:- Modified ANOVA table for significant diversity in number of species**

Source of Variation	Sum of squares	Degree of freedom	Mean Sum of Square	$F_{\text{calculated}}$	$F_{\text{tabulated}}$ at 5% level of significance
Between conservation category	205.75	4 - 1 = 3	68.583	1.601172 (2.68)	$F_{(3,12)} = 3.49$
Between vegetation type	358.8 (565.4)	5 - 1 = 4	89.7 (141.35)	2.09418 (3.3)	$F_{(4,12)} = 3.26$
Residuals	514 (307.4)	19 - 7 = 12	42.833 (25.62)		
Total	1078.55	20 - 1 = 19			



Thus modified mean sum of squares =  $42.833 * 3.3 = 141.35$  and modified sum of squares between various vegetation types =  $141.35 * 4 = 565.4$  by maintaining the total sum of square same value = 1078.55 and correcting the value of residuals ( $1078.55 - 565.4 = 513.15$ ). To check it for no significant variance in numbers between various threat and conservation categories, modified mean sum of squares between residuals =  $513.15/12 = 42.76$ . Then modified  $F_{\text{calculated}} = 2.68 < F_{\text{tabulated}} (3.49)$ . Hence both the objectives are satisfied by making the variance in number of species between threat and conservation categories non significant as well as making the variance of number of species between various vegetation types significant.

Thus to make the variance of number of between various vegetation types to be significant the necessary and sufficient condition is,

**Sum of squares between various vegetation type = 565.4**

$$R_1^2/4 + R_2^2/4 + R_3^2/4 + R_4^2/4 + R_5^2/4 - 884.45 = 565.4$$

$$R_1^2 + R_2^2 + R_3^2 + R_4^2 + R_5^2 = 4*(565.4 + 884.45)$$

$$R_1^2 + R_2^2 + R_3^2 + R_4^2 + R_5^2 = 4* 1449.85 = 5799.4$$

To find out the number of species in each vegetation type we have developed the criteria by assuming the existing value of four categories and calculating the other one thus five criteria have been developed which are as follows:

**Criteria 1:-**  $R_1$  is calculated assuming  $R_2 = 28, R_3 = 8, R_4 = 20, R_5 = 19$

$$R_1^2 + 28^2 + 8^2 + 20^2 + 19^2 = 5799.4 \text{ or } R_1 = 64$$

**Criteria 2:-**  $R_2$  is calculated assuming  $R_1 = 58, R_3 = 8, R_4 = 20, R_5 = 19$

$$58^2 + R_2^2 + 8^2 + 20^2 + 19^2 = 5799.4 \text{ or } R_2 = 40$$

**Criteria 3:-**  $R_3$  is calculated assuming  $R_1 = 58, R_2 = 28, R_4 = 20, R_5 = 19$

$$58^2 + 28^2 + R_3^2 + 20^2 + 19^2 = 5799.4 \text{ or } R_3 = 30$$

**Criteria 4:-**  $R_4$  is calculated assuming  $R_1 = 58, R_2 = 28, R_3 = 8, R_5 = 19$

$$58^2 + 28^2 + 8^2 + R_4^2 + 19^2 = 5799.4 \text{ or } R_4 = 35$$

**Criteria 5:-**  $R_5$  is calculated assuming  $R_1 = 58, R_2 = 28, R_3 = 8, R_4 = 20$

$$58^2 + 28^2 + 8^2 + 20^2 + R_5^2 = 5799.4 \text{ or } R_5 = 35$$

Thus there are five alternative criteria's, which are expressed in **table 6**, are available to explain the number of species to be maintained the variance in number of species between various vegetation types to be significant as well as maintaining non significant status of variance in number between various threat and conservation categories.

**Table 6:- Various alternative criteria's for different vegetation types**

Vegetation type	Number of species of vegetation types					Existing No. of species
	Criteria 1	Criteria 2	Criteria 3	Criteria 4	Criteria 5	
Trees	<b>64</b>	58	58	58	58	58
Shrubs	28	<b>40</b>	28	28	28	28
Lianas	8	8	<b>30</b>	8	8	8
herbs	20	20	20	<b>35</b>	20	20
Grasses	19	19	19	19	<b>35</b>	19
Total	139	145	<b>155</b>	148	149	133

**Criteria 3 are the feasible solution for optimality as this gives the highest total number of species of 155.** Thus 58 tree species, 28 shrubs, 30 lianas, 19 grass species (Total 155) are required to make the significant diversity in number. Thus the 22 number of more lianas species are required to enrich the study area for significant variance in number of species as well as non significant status between threat and conservation categories.

#### **4.9 Evaluation of optimum species diversity in study area in terms of average IVI of species with minimizing the threat and conservation status**

In order to make the significant species diversity and making the threat and conservation categories for average IVI of the species, non significant the **F** calculated < **3.49**, say **F** calculated = **3.49**. For making the non significant threat and conservation category in average IVI, ANOVA **table 4** is reformulated by backward calculation for mean sum of squares and total sum of squares between various vegetation types and it is expressed in bold and small brackets and modified ANOVA table is presented in **table 7**.

**Table 7:- Modified ANOVA table for non significant diversity for average IVI between threat and conservation categories**

Source of Variation	Sum of squares	Degree of freedom	Mean Sum of Square	F calculated	F tabulated at 5% level of significance
Between conservation category	1878.795 ( <b>414.81</b> )	4 - 1 = 3	626.265 ( <b>138.27</b> )	15.807 ( <b>3.49</b> )	F <sub>(3,12)</sub> = 3.49
Between vegetation type	333.85 ( <b>1797.835</b> )	5 - 1 = 4	83.463 ( <b>449.46</b> )	2.107 ( <b>11.344</b> )	F <sub>(4,12)</sub> = 3.26
Residuals	475.438	19 - 7 = 12	39.620		
Total	2688.083	20 - 1 = 19			

Between threat and conservation categories the mean sum of squares = 39.620 \* 3.49 = 138.27 and sum of squares = 138.27 \* 3 = 414.81, by maintaining the total sum of square same value = 2688.083 and correcting the value of Sum of squares between various vegetation types (2688.083 & 414.81 & 475.438 = 1797.835).

Thus modified Mean Sum of Square between vegetation types =  $1797.835/4 = 449.46$

To check the variance of average IVI between various vegetation types  $F_{\text{calculated}} = 449.46/39.62 = 11.344 > F_{\text{tabulated}} (3.26)$ . Hence maintaining the variance of average IVI of the species non significant between threat and conservation categories, the variance of average IVI between various vegetation types is significant as  $11.344 > 3.26$ . Thus both the objectives are full filled by making the variance of average IVI of the species between threat and conservation categories as **non significant** as well as the variance of average IVI between various vegetation types as **significant**.

Thus the necessary and sufficient condition for average IVI to be non significant between threat and conservation categories:

**Sum of squares between threat and conservation category = 414.81**

$$C_1^2/5 + C_2^2/5 + C_3^2/5 + C_4^2/5 \ \& \ 1185.95 = 414.81$$

$$C_1^2 + C_2^2 + C_3^2 + C_4^2 = 5*(1185.95 + 414.81)$$

$$C_1^2 + C_2^2 + C_3^2 + C_4^2 = 8003.6$$

To find out the average IVI between threat and conservation categories, four criteria's have been developed by assuming the three existing value of average IVI of threat and conservation categories to be constant and calculating the other one, these alternative criteria are:

**Criteria 1:-**  $C_1$  is calculated assuming  $C_2 = 10.578$ ,  $C_3 = 18.578$ ,  $C_4 = 2.81$ ,

$$C_1^2 + 10.578^2 + 18.578^2 + 2.81^2 = 8003.6 \text{ or } C_1 = 86.83$$

**Criteria 2:-**  $C_2$  is calculated assuming  $C_1 = 121.87$ ,  $C_3 = 18.578$ ,  $C_4 = 2.81$ ,

$$121.87^2 + C_2^2 + 18.578^2 + 2.81^2 = 8003.6 \text{ or } C_2 = \text{Imaginary}$$

**Criteria 3:-**  $C_3$  is calculated assuming  $C_1 = 121.87$ ,  $C_2 = 10.578$ ,  $C_4 = 2.81$ ,

$$121.87^2 + 10.578^2 + C_3^2 + 2.81^2 = 8003.6 \text{ or } C_3 = \text{Imaginary}$$

**Criteria 4:-**  $C_4$  is calculated assuming  $C_1 = 121.87$ ,  $C_2 = 10.578$ ,  $C_3 = 18.578$ ,

$$121.87^2 + 10.578^2 + 18.578^2 + C_4^2 = 8003.6 \text{ or } C_4 = \text{Imaginary}$$

Thus average IVI between threat and conservation categories in  $C_2$ ,  $C_3$  and  $C_4$  are not feasible and criteria 1 is important for non significant average IVI between threat and conservation categories i.e.  $C_1 = 86.83$ . Thus threat and conservation category 1 is very important category for average IVI of the species. The average IVI of the species in threat and conservation category 1 should be reduced by increasing the number of species between various vegetations types.

Thus to find out the average IVI within various vegetation types and between threat and conservation category 1 different alternatives are assessed by assuming the existing value of average IVI of within four vegetation types to be constant and calculating the other one, thus five

alternative criteria have been developed, which are as follows:

$$R_1 + R_2 + R_3 + R_4 + R_5 = 86.83$$

**Criteria 1:-**  $R_1$  is calculated assuming  $R_2 = 19.284$ ,  $R_3 = 44.125$ ,  $R_4 = 25.321$ ,  $R_5 = 10.067$

$$R_1 + 19.284 + 44.125 + 25.321 + 10.067 = 86.83 \text{ or } R_1 = \text{Negative}$$

**Criteria 2:-**  $R_2$  is calculated assuming  $R_1 = 23.073$ ,  $R_3 = 44.125$ ,  $R_4 = 25.321$ ,  $R_5 = 10.067$

$$23.073 + R_2 + 44.125 + 25.321 + 10.067 = 86.83 \text{ or } R_2 = \text{Negative}$$

**Criteria 3:-**  $R_3$  is calculated assuming  $R_1 = 23.073$ ,  $R_2 = 19.284$ ,  $R_4 = 25.321$ ,  $R_5 = 10.067$

$$23.073 + 19.284 + R_3 + 25.321 + 10.067 = 86.83 \text{ or } R_3 = 9.085$$

**Criteria 4:-**  $R_4$  is calculated assuming  $R_1 = 23.073$ ,  $R_2 = 19.284$ ,  $R_3 = 44.125$ ,  $R_5 = 10.067$

$$23.073 + 19.284 + 44.125 + R_4 + 10.067 = 86.83 \text{ or } R_4 = \text{Negative}$$

**Criteria 5:-**  $R_5$  is calculated assuming  $R_1 = 23.073$ ,  $R_2 = 19.284$ ,  $R_3 = 44.125$ ,  $R_4 = 25.321$

$$23.073 + 19.284 + 44.125 + 25.321 + R_5 = 86.83 \text{ or } R_5 = \text{Negative}$$

Thus there are five alternative criteria's, which are expressed in **table 8**, are available to explain the average IVI of species to be maintained the variance in average IVI of the species between various vegetation types to be significant as well as maintaining non significant status of variance in average IVI of the species between various threat and conservation categories.

**Table 8:- Various alternative criteria's, between different vegetation types for average IVI**

Vegetation type	Different criterias for average IVI					Existing average IVI in C <sub>1</sub>
	Criteria 1	Criteria 2	Criteria 3	Criteria 4	Criteria 5	
Trees	<b>-ve</b>	23.073	23.073	23.073	23.073	23.073
Shrubs	19.284	<b>-ve</b>	19.284	19.284	19.284	19.284
Lianas	44.125	44.125	<b>9.085</b>	44.125	44.125	44.125
herbs	25.321	25.321	25.321	<b>-ve</b>	25.321	25.321
Grasses	10.067	10.067	10.067	10.067	<b>-ve</b>	10.067
<b>Total</b>	<b>86.83</b>	<b>86.83</b>	<b>86.83</b>	<b>86.83</b>	<b>86.83</b>	121.87

The Criteria 3 is optimal and feasible solution for the average IVI in threat and conservation category 1. The other four criteria's (criteria 1, 2, 4 and 5) show the negative value of average IVI with in various vegetation types. The average IVI of the species between threat and conservation category 1 is the critical value of 86.83. This suggests that average IVI between

threat and conservation category 1 should be 23.073, 19.284, 9.085, 25.321 and 10.067 with in trees, shrubs, lianas, herbs and grasses respectively. The number of lianas species should be increased to lower the average IVI of the species to the value of 9.085 as compare to 44.125. The liana species is the critical one which should be enriched in study area so that average IVI between threat and conservation category 1 assumes the lower value 9.085.

## 5. CONCLUSION:

Presently in the study area there were species of 58 trees, 28 shrubs, 08 lianas, 20 herbs and 19 grasses (total 133). The species under threat and conservation category 4 were likely to be extinct in near future locally. The category 3 species could go to category 4 in near future. These species were mentioned in the study area which would be monitored local forest officer.

At present status of the richness in the study area indicated that variance of the number of species between various vegetation types (trees, shrubs, lianas, herbs and grasses) and between threat and conservation categories are not significant at 5% level of significance.

The present status of species diversity in study area, when the variance of average IVI between various vegetation types was taken, there was no richness of species diversity. But there was a significant difference in variance of average IVI of the species between the threat and conservation categories. Hence average IVI of the species is an important parameter as IVI of particular species includes the frequency, density and dominance in itself. Thus, the study area does not show the richness in species diversity but it shows the high stress in threat category when average IVI of species were taken as parameter.

In order to maintain the variance in number of species between various vegetation types to be significant as well as maintaining non significant status of variance in number between various threat and conservation categories, the optimal numbers of species were 155 in study area. The species of 58 tree, 28 shrubs, 30 lianas, 20 herbs, 19 grasses (Total 155) were required to make the significant diversity in number. Thus the 22 number of more lianas species were required to enrich the study area for significant variance in number of species as well as non significant status between threat and conservation categories.

In order to reduce the stress and improve the richness the variance of average IVI of the species between threat and conservation categories as **non significant** as well as the variance of average IVI between various vegetation types as **significant**. In order to fulfill the above the two objectives simultaneously, the average IVI in threat and conservation category 1 was should be 86.83 which was optimum value of IVI. The average IVI between threat and conservation category 1 should be 23.073, 19.284, 9.085, 25.321 and 10.067 within trees, shrubs, lianas, herbs and grasses respectively. The number of lianas species should be increased to lower the average IVI of the species to the value of 9.085 as compared to 44.125. The liana species was the limiting species which should be enriched in study area so that average IVI between threat and conservation category 1 assumes the lower value 9.085.



## ACKNOWLEDGEMENT

Authors are thankful to staff member of forest department of Satna, Rewa and Shivpuri for providing help during study. Authors are also grateful to Nepal Singh Govindgarh, Ravishankar Yadav, Rajesh Kumar Kol and L.S. Kushwaha for providing valuable knowledge for identifying the species.

## REFERENCES

1. Shannon C. L. A Mathematical theory of communication. The Bell system Technical Journal. 1948; 27:373-423 and 623-656.
2. Simpson E. H. Measurement of diversity Nature. 1949; 688.
3. Joshi H.G. Vegetation structure, floristic composition and soil nutrient status in three sites of Tropical dry deciduous forest of West Bengal, India, Indian Journal of Fundamental and Applied Life Science. 2012; 2(2):355-364.
4. Hajara P.K. & Mudgal V. An overview, BSI India.1997.
5. Reddy C. S. Life Sci J. 2008; 5(2): 84-89.
6. Dixit A.M. Ecological evaluation of dry tropical forest vegetation: An approach to environmental impact assessment, Tropical Ecology. 1997; 38:87-100.
7. Kumar A., Marcor B.G. & Saxena A. Tree species diversity and distribution pattern in tropical forest of Garohills, Curr. Sci. 2006; 91(10) 1370-1381.
8. Jain S.K. & Rao R.R. An assessment of threatened plants of India, Botanical survey of India, Culcutta. 1983.
9. Nayar M.P., Puspangadan P. Ravi, K & Santosh, V. Conservation of rare and endangered species of Indian flora, strategies for botanic gardens. In (eds) conservation and economic evaluation of biodiversity, Oxford & IBH publishing co. New Delhi. 1997; vol.1, pp.47-57.
10. Myres N. Threatened Biotas: 'Hotspots' in tropical forests, the Environment. 1988;8 (3):987-208.
11. Nayar M. P. & Sastry A. R. K.. Red data book of Indian plants, vol. 1, Botanical Survey of India, Culcutta, CAMP, CAMP Workshop, Shimla. 1987.
12. Nayar M. P. & Sastry A. R. K. Red data book of Indian plants, vol. 2, Botanical Survey of India, Culcutta, CAMP, CAMP Workshop, Shimla. 1988.
13. Nayar M. P. & Sastry A. R. K. Red data book of Indian plants, vol. 3, Botanical Survey of India, Culcutta, CAMP, CAMP Workshop, Shimla. 1990.
14. Nautiyal Sunil. Plant Biodiversity and its Conservation in Institute for Social and Economic change (ISEC) campus, Bangalore; A case study. J. Biodiversity. 2003;2(1)9-26.
15. Singh Prachi, Singh Neeta, Singh P. K., Sahu Sherendra & Singh Anoop. ECOLOGY, ENVIRONMENT AND CONSERVATION, accepted paper with Ref. No. EEC-3003. 2017; September Supplement Issue.
16. Singh Prachi, Sahu Sherendra K., Singh P. K. & Singh Neeta. International Journal of Fauna Biological Studies. 2017; 4(4):120-125.
17. Singh Prachi, Singh Neeta, Singh P. K. & Sahu Sherendra. VINDHYA BHARTI. 2017;No. 14 JulyVol. I.
18. Singh Prachi, Singh P. K., Sahu Sherendra, Singh Anoop, Gaurav N. & Singh Neeta. Journal of Pharmacognosy and Phytochemistry. 2017;6(4), 544-549.
19. Singh Prachi, Singh P. K., Sahu Sherendra, Singh Anoop, Gaurav N. & Singh Neeta. Journal of Medicinal Plants Studies. 2017; 5(4), 07-12.
20. Anonymous. Working plan code published by Chief Conservator Forests, working plan Satpuda Bhawan Bhopal, Madhya Pradesh Forest department. 1996.
21. Elhance D.L. Book Estimating sample size for population proportion, Sarojaini Naidu marg, Allahabad. 1994; Page No. 21.14 published by kitabmahal, 22-A.
22. Jain Atul Kumar IFS, Forest resource survey, Government of Madhya Pradesh Forest Department, Chapter of Forest resource Survey. 2008; Part-1, Page 47 to 53, working plan of Satna revised for 2008-09 to 2017-18.
23. Mishra R. Ecology work book, Page (36 to 48) Professor and head of Department of Banaras Hindu University Varansi India. 1968; published by Oxford & IBH Co. 17, Park street Calcutta-16.
24. Elhance D.L. Properties of normal distribution, book of Fundamentals of Statistics. 1994; page 18.32.

25. Singh P. Study of Diversity of Angio spermic flora of Mukundpur range (District Satna) with special reference to conservation of endangered plants. 2018; PhD thesis; 02: 22.
26. Murthy Rama P. Book of Operation Research, Published at NEW AGE INTERNATIONAL (P) LIMITED 4835/24, Ansari Road, Daryaganj, New Delhi-110002. 2007.
27. Pereira, H. & Cooper, H. Towards the global monitoring of biodiversity change. *Trends Ecol. Evol.* 21. 2006: 123-129.
28. Pereira, H. M., Paul W. L., Vania P., Rob A., Jorn P. W. S., Juan F. FM., Miguel B. A., Patricia B., Reinette B., William W. L. C., Louise C., Cooper H.D., Eric L.G., Sylvie G., George C. H., Henry P. H., Georgina M.M., Thierry O., Carmen R., Patricia R., Rpnert J. S., Ussif R.S. & Matt W. Scenarios for global biodiversity in the 21st century. *Science* 330. 2010: 1496-1501.
29. Chiarucci A. Estimating species richness: still a long way off ! *Journal of Vegetation Science* 23(6). 2012: 3-21.
30. Fager E. W. Diversity: a sampling study. *Am. Natur.* 106. 1972: 293-310.
31. Hill M. O. Diversity and evenness: a unifying notation and its consequences. *Ecology* 54. 1973: 427-432.
32. Hurlbert S.H. The nonconcept of species diversity: a critique and alternative parameters. *Ecology* 52. 1971: 577-586.
33. Liold M., Zar J. H. & Karr J. R. On the calculation of information theoretical measures of diversity. *Am. Midl. Natur.* 79. 1968: 257-272.
34. McIntosh R. P. An index of diversity and the relation of certain concepts to diversity. *Ecology* 48. 1967: 392-404.