

GROWTH AND YIELD ATTRIBUTES OF CAULIFLOWER AS INFLUENCED BY MICRONUTRIENTS AND PLANT SPACING

ABSTRACT

The experiment was conducted in the Horticulture Farm, Sher-e-Bangla Agricultural University, Dhaka from November 2015 to February 2016 with the aim of investigating the growth and yield attributes of cauliflower as influenced by different micronutrients and plant spacing. The experiment consisted of two factors, such as Factor A: Plant spacing (3 levels) as- S_1 : 50 cm \times 50 cm, S_2 : 50 cm \times 40 cm, S_3 : 50 cm \times 30 cm and Factor B: Levels of micronutrients (4 levels) as- T_0 : B_0Mo_0 (control), T_1 : $B_{1.0}Mo_{0.5}$ kg/ha, T_2 : $B_{2.0}Mo_{1.0}$ kg/ha, T_3 : $B_{3.0}Mo_{1.5}$ kg/ha. The experiment was laid out in Randomized Complete Block Design with three replications. All the studied parameters were significantly influenced by micronutrients and plant spacing. In case of plant spacing, the highest curd yield (39.89 t/ha) was found from S_2 and the lowest curd yield (35.00 t/ha) was found from S_1 . For micronutrients, T_2 treatment produced the highest curd yield (46.85 t/ha) and the lowest (24.41 t/ha) was from control. In case of combined effect, the highest curd yield (51.56 t/ha) was obtained from S_2T_2 and the lowest curd yield (20.33 t/ha) from S_1T_0 . Therefore, it can be suggested that the highest curd yield and good shape cauliflower curd can be obtained in plant spacing 50 cm \times 40 cm with the combined application of B 2.0 kg/ha and Mo 1.0 kg/ha.

Keywords: Growth, yield, micronutrients, spacing, cauliflower

1. INTRODUCTION

Cauliflower (*Brassica oleraceae* var. *botrytis* L.) is one of the most popular cruciferous vegetable crops cultivated for its white curds as edible part. It is being grown round the year for its white and tender curd vegetables and thrives best in a cool moist climate and it does not withstand very low temperature or too much heat [1]. Cauliflower is a very tasty and much popular vegetable in Bangladesh as well as all over the world. Due to increasing consumption of cauliflower products, the crop is becoming promising. Although Bangladesh is producing a good amount of cauliflower and it is using for the preparation of different delicious food but average yield of cauliflower is low in Bangladesh compared to other countries. Plant density as management practices and micronutrients is prerequisite for increasing the production of cauliflower in Bangladesh [2]. Plant spacing is an important aspect of crop production for maximizing the yield [3]. It helps to increase the number of leaves, branches and healthy foliage. Densely planted crop obstruct the proper growth and development. On the other hand, wider spacing ensures the basic requirements but decrease the total number of plants as well as total yield. Crop yield may be increased up to 25% by using optimum spacing [4]. Cauliflower responds well to macro nutrients–nitrogen, phosphorus and potassium. However, micro–nutrients are also essential for its proper growth and yield especially boron and molybdenum [3]. Boron application increased plant height, number of leaves per plant, length and width of the leaf, plant spread, main head weight and head yield both per plant and per hectare [5]. On the other hand, due to boron deficiency water soaked areas appear on the stem and head surface, gradually the stem becomes hollow and curd turns brown. Again the molybdenum deficiency appears on young plant with chlorosis of leaf margins and gradually the whole leaf turns white. They

also become cupped and wither, eventually. The leaf dies and the growing point also collapses [6]. It was known that there could be many genetic and environmental effects on the yield [23]. Considering the above all perspective, the present study was undertaken to investigate the effect of plant spacing and different levels of boron and molybdenum on cauliflower to find out the suitable combination of plant spacing and micronutrients which can ameliorate the growth and yield attributes of cauliflower.

2. MATERIAL AND METHODS

2.1 Experimental Site

The experiment was conducted at the Horticultural farm of Sher-e-Bangla Agricultural University, Dhaka, Bangladesh from November 2015 to February 2016. The location of the experimental site was 23°74'N latitude and 90°35'E longitude and at an elevation of 8.2 m from sea level. The climate of experimental site was under the subtropical climate, characterized by three distinct seasons, the winter season from November to February and the pre-monsoon or hot season from March to April and the monsoon period from May to October. The soil of the experimental area belongs to the Modhupur Tract (AEZ No 28). It had shallow red brown terrace soil. The selected plot was medium high land and the soil series was Tejgaon.

2.2 Planting Material

The seeds of cauliflower cv. Snowball were collected from the Seed Wing, BADC (Bangladesh Agriculture Development Corporation), Dhaka.

2.3 Experimental Design and Treatments

The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications. The experiment consisted of two factors, such as Factor A: Plant spacing (3 levels) as- S₁: 50 cm × 50 cm, S₂: 50 cm × 40 cm, S₃: 50 cm × 30 cm and Factor B: Levels of micronutrients (4 levels) as- T₀: B₀Mo₀ (control), T₁: B_{1.0}Mo_{0.5} kg/ha, T₂: B_{2.0}Mo_{1.0} kg/ha, T₃: B_{3.0}Mo_{1.5} kg/ha. There were 36 unit plots and the size of each plot was 2.0 m × 1.8 m. The distance between two blocks and two plots were 1.0 m and 0.5 m, respectively. The soil was having a texture of sandy loam with pH and organic matter 6.1 and 1.18%, respectively. Total N (%), available P (µg/g), exchangeable K (meq/100g) were 0.091, 16.0 and 0.32 respectively. The micronutrients i.e. boron and molybdenum were applied as three consecutive foliar spray at 30, 40 and 50 days after transplanting (DAT) as per treatments. The sources of micronutrients boron and molybdenum were obtained from Borax (contain 11% boron) and Ammonium molybdate (contain 98.99% tracer element).

2.4 Growth condition of Cauliflower & Measurements of Parameters

Seedlings were grown following proper methods and all of the cultural practices were done properly. Seeds (5g) were sown in the well prepared seedbed (3×1) m² on November 01, 2015. Application of manure and fertilizers were applied as per treatment. The crop was grown with the recommended dose of N: P: K (138:96:150 kg/ha) and FYM@ 20 ton/ha. Full dose of phosphorus, potash and half dose of nitrogen were applied as soil application before transplanting. The remaining half dose of nitrogen was applied at 30 days after transplanting (DAT). Healthy and uniform seedlings of 20 days old seedlings were transplanting in the experimental plots on 20 November, 2015. Intercultural practices viz. gap filling, weeding, earthing up, irrigation, pest and disease control etc. were done as per requirements. The hand weeding was done 15, 30 and 45 days after transplanting to keep the plots free from weeds. For controlling leaf caterpillars Nogos @ 1 ml/L water were applied two times at an interval of 10 days starting soon after the appearance of infestation. All cauliflower curd was not matured at a same time, harvesting was done at 15 February to 02 March. Different yield contributing data have been recorded from the mean of five harvested plants which was selected at random of each unit plot of every harvesting stage.

2.5 Data Collection and Analysis

Five plants were randomly selected from each unit plot for the collection of data. The plants in the outer rows and the extreme end of the middle rows were excluded from the random selection to avoid the border effect. The height of the plants was measured from the ground level to the tip of the highest leaves using a meter scale. Dry matter content of leaves and curd was measured with the formula -Dry matter content of (leaves/curd) = $\frac{\text{Dry weight}}{\text{Fresh weight}} \times 100$. At first leaves and

curd were cut into pieces and was dried under sunshine for 3 days and then dried in an oven at 70°C for 72 hours. The sample was then transferred into desiccators and allowed to cool down at room temperature. The final weight of the sample was taken and then measured with the above formula. The data obtained for different parameters were statistically analyzed to find out the significance difference of variety and different fertilizer application on yield and yield contributing characters of cabbage. The mean values of all the characters were calculated and analysis of variance was performing by the 'F' (variance ratio) test. The significance of the difference among the treatment combinations means was estimated by the Duncan's Multiple Range Test (DMRT) at 5% level of probability [7].

3. RESULTS AND DISCUSSION

3.1 Plant height

Plant height of cauliflower showed significant influence due to different plant spacing at 30, 40, 50, 60 DAT and at harvest (Fig.1). The significantly superior plant height (59.84 cm) was observed from S₃ (50 cm × 30 cm) treatment at harvest which was statistically similar (55.87 cm) to S₂ (50 cm × 40 cm) treatment, while the shortest plant (46.56 cm) was found from S₁ (50 cm × 50 cm) treatment at the same growth stage (Fig.1). The variation in plant height as influenced by spacing was perhaps due to proper utilization of nutrient, moisture and light. [3] reported the maximum plant height (49.33 cm) where the plants were spaced 45×50 cm apart.

During growing period plant height gradually increased with time and reached to the maximum at harvest. Plant height was significantly influenced by micronutrients at 30, 40, 50, 60 DAT and at harvest (Fig. 2). At the time of harvest, the tallest plant (61.89 cm) was found from T₂ (B_{2.0}Mo_{1.0} kg/ha) treatment which was followed (55.89 cm and 53.57 cm) by T₃ (B_{3.0}Mo_{1.5} kg/ha) and T₁ (B_{1.0}Mo_{0.5} kg/ha) treatments and they were statistically similar, whereas the shortest plant (45.00 cm) was recorded from T₀ (B₀Mo₀ i.e. control) treatment (Fig. 2). The results indicate that the increasing rate of micronutrients significantly increase the plant height. [8] reported that application of boron increased the plant height of cauliflower and our finding is in agreement with their findings.




Figure 1: Effect of different plant spacing on plant height of cauliflower at different DAT




Figure 2: Effect of micronutrients on plant height of cauliflower at different DAT

Combined effect of different plant spacing and micronutrients showed statistically significant variation on plant height of cauliflower at 30, 40, 50, 60 DAT and at harvest (Table 1). The maximum plant height (69.23 cm) was recorded from S_1T_2 (50 cm \times 50 cm plant spacing with $B_{2.0}Mo_{1.0}$ kg/ha) treatment at harvest which was statistically similar (66.73 cm) to S_2T_2 (50 cm \times 40 cm plant spacing with $B_{2.0}Mo_{1.0}$ kg/ha), while the shortest plant (36.43 cm) was observed from S_1T_0 (50 cm \times 50 cm plant spacing with B_0Mo_0 kg/ha) treatment (Table 1).

3.2 Number of leaves per plant

The number of leaves per plant of cauliflower was significantly influenced by the different plant spacing at 30, 40, 50, 60 DAT and at harvest (Fig.3). An increasing trend in number of leaves per plant was found up to harvest for all the treatments. The highest number of leaves per plant (18.00) was recorded from S_2 treatment at harvest time which was statistically similar (17.67) to S_1 treatment, whereas the lowest number of leaves per plant (16.53) was found from S_3 (Fig.3) treatment at the same growth stage of plant. It was observed that the number of leaves was higher in plants with wider spacing and lower in closely plants. It is probably, due to reduce inter plant competition for access to nutrients, moisture and other resources. Similar trend was reported by [21].

Table 1: Combined effect of different plant spacing and micronutrients on plant height at different days after transplanting (DAT) and harvest of cauliflower

Treatments	Plant Height at				
	30 DAT	40 DAT	50 DAT	60 DAT	Harvest
S_1T_0	15.24 g	19.25 h	23.46 i	31.57 g	36.43 g
S_1T_1	18.19 ef	22.80 fg	27.59 h	35.95 fg	39.02 fg
S_1T_2	27.42 a	34.81 a	46.26 a	57.80 a	69.23 a
S_1T_3	21.01 de	25.30 ef	32.93 fg	44.18 de	41.56 f
S_2T_0	16.57 fg	21.37 gh	30.69 gh	39.12 ef	42.70 f
S_2T_1	20.80 de	29.65 b-d	40.22 cd	49.81 c	58.28 c-e

S ₂ T ₂	26.02 ab	34.45 ab	45.25 ab	56.14 ab	66.73 ab
S ₂ T ₃	23.17 bcd	30.94 bc	42.51 bc	53.91 abc	59.38 cd
S ₃ T ₀	20.62 de	26.64 de	35.43 ef	50.38 bc	55.88 de
S ₃ T ₁	24.08 bc	32.68 ab	45.06ab	51.85 bc	63.42 bc
S ₃ T ₂	21.34 cd	28.82 cd	38.22de	49.45 cd	53.32 e
S ₃ T ₃	25.88 ab	32.91 ab	44.95 ab	54.56 abc	63.12 bc
LSD(0.05)	2.735	3.193	3.286	5.279	4.825
CV (%)	8.64	7.17	5.15	6.51	5.27

In a column, means followed by same letter (s) do not differ significantly at 5% level of probability, Factor A: Plant spacing (3 levels) as- S₁: 50 cm × 50 cm, S₂: 50 cm × 40 cm, S₃: 50 cm × 30 cm and Factor B: Levels of micronutrients (4 levels) as- T₀: B₀Mo₀ (control), T₁: B_{1.0}Mo_{0.5} kg/ha, T₂: B_{2.0}Mo_{1.0} kg/ha, T₃: B_{3.0}Mo_{1.5} kg/ha.




Figure 3; Effect of plant spacing on number of leaves per plant at different DAT and harvest of cauliflower

The number of leaves per plant was found to be significantly influenced by the application of micronutrients at 30, 40, 50, 60 DAT and at harvest (Fig 4). At harvest, the highest number of leaves per plant (19.27) was recorded from T₂ treatment which was followed (18.62 and 17.71) by T₃ and T₁ treatment and they were statistically similar, while the lowest number of leaves per plant (14.00) was found from T₀ (Fig 4). Thakur *et al.*[8] reported that application of boron increased the number of leaves per plant of cauliflower. Sharma [9] in cauliflower, who stated that the probable reasons for enhanced plant height and the number of leaves, may be due to promoting effects of molybdenum on vegetative growth which ultimately lead to more photosynthetic activities. Similar findings were also reported by [10, 11, 12].




Figure 4: Effect of micronutrients on number of leaves per plant at different DAT and harvest of cauliflower

The combined effect of different plant spacing and micronutrients showed statistically significant variation on number of leaves per plant of cauliflower at 30, 40, 50, 60 DAT and at harvest (Table 2). At harvest, the highest number of leaves per plant (20.27) was recorded from S_2T_2 whereas the lowest number of leaves per plant (13.53) was found from S_1T_0 treatment combination (Table 2). [6] reported that the number of leaves/plant increased significantly with the different treatments of boron and molybdenum at every stage of observations and our findings is in conformity with their findings.

3.3 Days to curd initiation

Days to curd initiation of cauliflower showed significant differences due to different plant spacing (Table 3). The maximum days to curd initiation (57.50) was observed from S_1 treatment which was closely followed (53.75) by S_2 treatment, while the minimum days to curd initiation (49.42) was found from S_3 treatment (Table 3). The maximum days required for curd initiation in wider spacing might be attributed due to the less interplant competition, which resulted in better vegetative growth of plants. Similar result was observed by [2].

Table 2. Combined effect of different plant spacing and micronutrients on number of leaves per plant at different DAT and harvest of cauliflower

Treatments	Number of Leaves per Plant at				
	30 DAT	40 DAT	50 DAT	60 DAT	Harvest
S_1T_0	3.67 d	6.13 d	10.53 g	10.93 g	13.53 e
S_1T_1	6.00 ab	9.30 ab	14.73 bc	18.13 bc	19.20 ab
S_1T_2	6.60 ab	9.53 a	15.27 ab	19.93 ab	19.93 a
S_1T_3	5.87 b	8.00 c	13.67 cd	16.60 c-e	18.00 c
S_2T_0	4.80 c	6.93 d	11.53 fg	12.33 fg	14.20 e
S_2T_1	6.20 ab	9.30 b	15.13 b	19.80 b	19.27 b
S_2T_2	6.87 a	9.80 a	16.13 a	21.40 a	20.27 a
S_2T_3	6.13 ab	8.33 bc	14.47 bc	17.87 bc	18.27 bc
S_3T_0	4.60 c	6.87 d	11.20 fg	11.60 g	14.27 e
S_3T_1	4.53 cd	6.57 d	13.00 de	17.47 bcd	17.40 cd
S_3T_2	4.40 cd	6.27 d	12.00 ef	14.47 ef	17.60 cd

S ₃ T ₃	4.33 cd	6.60 d	12.20 ef	15.20 de	16.87 d
LSD(0.05)	0.799	1.015	1.069	2.453	0.994
CV (%)	9.57	8.09	5.13	9.46	6.58

In a column, means followed by same letter (s) do not differ significantly at 5% level of probability, Factor A: Plant spacing (3 levels) as- S₁: 50 cm × 50 cm, S₂: 50 cm × 40 cm, S₃: 50 cm × 30 cm and Factor B: Levels of micronutrients (4 levels) as- T₀: B₀Mo₀ (control), T₁: B_{1.0}Mo_{0.5} kg/ha, T₂: B_{2.0}Mo_{1.0} kg/ha, T₃: B_{3.0}Mo_{1.5} kg/ha.

Statistically significant variation was recorded for micronutrients in terms of days to curd initiation of cauliflower (Table 3). The maximum days to curd initiation (59.22) was found from T₀ treatment whereas the minimum days to curd initiation (48.00) was recorded from T₂ treatment (Table 3). The combined effect of different plant spacing and micronutrients combination showed significant variation on days to curd initiation of cauliflower (Table 4). The maximum days to curd initiation (65.33) was recorded from S₁T₀ which was statistically similar (62.00 and 61.67) to S₁T₃ and S₁T₁, while the minimum days to curd initiation (41.00) was found from S₁T₂ treatment combination (Table 4). It was observed that the curd initiation period required in plants decreased with the increasing levels of micronutrients application.; This might be due to the positive role played by the regulating micronutrients in balanced absorption of nutrients might improve physiological activities, which resulted the endogenous growth hormone synthesis responsible for early curd formation in plants. The present result is in agreement with the findings of [18].

3.4 Dry matter content of leaves

Dry matter content of leaves of cauliflower showed significant differences due to different plant spacing (Table 3). The highest dry matter content of leaves (12.57 %) was observed from S₂ treatment which was statistically similar (12.24%) to S₁ treatment, while the lowest dry matter content of leaves (11.90 %) was found from S₃ treatment (Table 3). Statistically significant variation was recorded for micronutrients in terms of dry matter content of leaves of cauliflower (Table 3). The highest dry matter content of leaves (12.94%) was found from T₂ treatment whereas the lowest dry matter content of leaves (10.84 g) was recorded from T₀ treatment (Table 3). [13] reported similar kind or of results. Combined effect of different plant spacing and micronutrients showed significant variation on dry matter content of leaves of cauliflower (Table 4). The highest dry matter content of leaves (13.62 %) was recorded from S₂T₂ while the lowest dry matter content of leaves (10.49 %) was observed from S₁T₀ treatment combination (Table 4).

Table 3. Effect of different plant spacing on yield attributes and yields of cauliflower

Treatments	Days to curd initiation	Dry matter content of leaves (%)	Dry matter content of curd (%)	Diameter of curd (cm)	Curd yield/plant (kg)	Curd yield hectare (ton)
S ₁	57.50 a	12.24 ab	13.29 ab	9.07 a	1.05 a	35.00 b
S ₂	53.75 b	12.57 a	13.93 a	8.21 b	0.90 b	39.89 a
S ₃	49.42 c	11.90 b	12.69 b	8.10 b	0.55 c	36.39 b
LSD(0.05)	3.812	0.379	0.703	0.390	0.038	1.840
CV (%)	8.16	4.39	6.75	4.87	5.31	5.86
Treatments	Days to curd initiation	Dry matter content of leaves (%)	Dry matter content of curd (%)	Diameter of curd (cm)	Curd yield/plant (kg)	Curd yield hectare (ton)
T ₀	59.22 a	10.84 c	12.56 c	7.56 c	0.53 d	24.41 d
T ₁	53.78 b	12.37 b	13.24 b	8.43 b	0.80 c	35.67 c
T ₂	48.00 c	12.94 a	14.10 a	9.00 a	1.05 a	46.85 a
T ₃	53.89 b	12.79 ab	13.31 ab	8.85 ab	0.94 b	41.44 b
LSD(0.05)	4.402	0.439	0.811	0.449	0.044	2.125
CV (%)	8.16	4.39	6.75	4.87	5.31	5.86

In a column, means followed by same letter (s) do not differ significantly at 5% level of probability, Factor A: Plant spacing (3 levels) as- S_1 : 50 cm \times 50 cm, S_2 : 50 cm \times 40 cm, S_3 : 50 cm \times 30 cm and Factor B: Levels of micronutrients (4 levels) as- T_0 : B_0Mo_0 (control), T_1 : $B_{1.0}Mo_{0.5}$ kg/ha, T_2 : $B_{2.0}Mo_{1.0}$ kg/ha, T_3 : $B_{3.0}Mo_{1.5}$ kg/ha.

3.5 Dry matter content of curd

It is obvious from the Table 3 that the dry matter content of curd significantly influenced by the different plant spacing. The maximum dry matter content of curd (13.93%) was observed from S_2 treatment which was statistically similar (13.29 %) to S_1 treatment, while the lowest dry matter (12.69%) from S_3 treatment (Table 3). Statistically significant variation was recorded for micronutrients in terms of dry matter content of leaves of cauliflower (Table 3). The highest dry matter content of curd (14.10 %) was found from T_2 treatment whereas the lowest dry matter content of curd (12.56%) from T_0 treatment (Table 3). [8] reported that application of boron increased the dry matter content of cauliflower.

Combined effect of different plant spacing and micronutrients showed statistically significant variation on dry matter content of curd of cauliflower (Table 4). The highest dry matter content of curd (15.63%) was recorded from S_2T_2 which was statistically similar (15.00%, 14.97% and 14.41%) to S_2T_3 , S_1T_2 and S_2T_1 , while the lowest dry matter content of curd (11.00%) was observed from S_1T_0 treatment combination (Table 4). This increase might be due to the collective effect of boron and molybdenum. Boron plays role in enhancing the translocation of carbohydrates from the site of synthesis to reproductive tissues in the curd, whereas, molybdenum stimulates the photosynthesis and enhance the metabolic process. Such significant response of micronutrients has also been reported by [14, 15, 16, 17]

3.6 Diameter of curd

Diameter of curd of cauliflower showed significant differences due to different plant spacing (Table 3). The highest diameter of curd (9.07 cm) was observed from S_1 treatment, while the lowest diameter of curd (8.10 cm) was found from S_3 treatment. Formation of bigger curd at the widest spacing was probably due to the availability of more nutrients, light, moisture to the plants. On the other hand, in closer spacing plants inter plants competition resulted in formation of small curd. [2, 3] reported similar kind or result and our findings is in corroboration with their findings. Statistically significant variation was recorded for micronutrients in terms of diameter of curd of cauliflower (Table 3). The highest diameter of curd (9.00 cm) was found from T_2 treatment whereas the lowest diameter of curd (7.56 cm) from T_0 treatment (Table 3). [18] reported that B and Mo application significantly increased curd diameter of cauliflower. Combined effect of different plant spacing and micronutrients showed statistically significant variation on diameter of curd of cauliflower (Table 4). The highest diameter of curd (9.90 cm) was recorded from S_1T_2 treatment combination while the lowest diameter of curd (7.07 cm) was observed from S_3T_0 treatment combination (Table 4). The formation of bigger curd with the application of higher levels of micronutrients might be done to higher synthesis of carbohydrate and their translocation to the curd, which subsequently helped in the formation of higher curd of cauliflower. Similar results have been reported by [14, 16, 18, 19].

3.7 Curd yield per plant

Weight of curd per plant of cauliflower showed significant differences due to different plant spacing (Table 3). The highest weight of curd per plant (1.05 kg) was observed from S_1 treatment which was closely followed (0.90 kg) to S_2 treatment, while the lowest weight of curd per plant (0.55 kg) was found from S_3 treatment (Table 3). Similar kind or result was reported by [2, 3]. Statistically significant variation was recorded for micronutrients in terms of weight of curd per plant of cauliflower (Table 3). The highest weight of curd per plant (1.05 kg) was found from T_2 treatment whereas the lowest weight (0.53 kg) from T_0 treatment which was followed (0.80 kg) by T_1 treatment (Table 3). [8] reported that the application of boron increased the curd yield of cauliflower. In case of combined effect of different plant spacing and micronutrients, statistically significant variation on curd yield per plant of cauliflower was found (Table 4). The highest weight of curd per plant (1.33 kg) was recorded from S_1T_2 which was statistically similar (1.27 kg) to S_1T_3 , while the lowest weight of curd per plant (0.44 kg) was observed from S_3T_0 treatment combination (Table 4). Increase in yield might be due to the combined application of boron and molybdenum at optimum levels under deficient condition increased uptake of major nutrients which resulted in sturdy plant growth and increased yield and quality. Present result confirms findings of [16] who found maximum yield with the combined foliar application of boron and molybdenum in cauliflower.

278
279
280

Table 4. Combined effect of different plant spacing and micronutrients on yield attributes and yield of cauliflower

Treatments	Days to curd initiation	Dry matter content of leaves (%)	Dry matter content of curd (%)	Diameter of curd (cm)	Curd yield per plant (kg)	Curd yield per hectare (ton)
S ₁ T ₀	65.33 a	10.49 f	11.00 d	8.10 de	0.61 de	20.33 e
S ₁ T ₁	61.67 ab	12.36 cd	13.69 bc	9.06 abc	0.99 c	33.00 d
S ₁ T ₂	41.00 f	13.46 ab	14.97 ab	9.90 a	1.33 a	44.33 b
S ₁ T ₃	62.00 ab	12.64 bc	13.48 bc	9.22 a-c	1.27 a	42.33 b
S ₂ T ₀	53.67 b-e	11.12 ef	11.67 d	7.50 ef	0.53 fg	23.56 e
S ₂ T ₁	55.33 b-d	12.57 c	14.41 ab	8.44 cd	0.94 c	41.78 bc
S ₂ T ₂	47.67 def	13.62 a	15.63 a	9.41 ab	1.16 b	51.56 a
S ₂ T ₃	54.33 b-e	12.97 a-c	15.00 ab	7.91 def	0.96 c	42.67 bc
S ₃ T ₀	46.00 ef	10.91 f	14.01 b	7.07 f	0.44 h	29.33 d
S ₃ T ₁	50.33 c-e	12.18 cd	11.62 d	7.81 def	0.48 gh	32.22 d
S ₃ T ₂	55.33 b-d	11.73 de	11.70 d	8.11 de	0.67 d	44.67 b
S ₃ T ₃	58.00 a-c	12.77 bc	12.45 cd	8.98 bc	0.59 ef	39.33 c
LSD(0.05)	7.625	0.758	1.407	0.779	0.076	3.681
CV (%)	8.16	4.39	6.75	4.87	5.31	5.86

281 In a column, means followed by same letter (s) do not differ significantly at 5% level of probability,
282 Factor A: Plant spacing (3 levels) as- S₁: 50 cm × 50 cm, S₂: 50 cm × 40 cm, S₃: 50 cm × 30 cm and
283 Factor B: Levels of micronutrients (4 levels) as- T₀: B₀Mo₀ (control), T₁: B_{1.0}Mo_{0.5} kg/ha, T₂: B_{2.0}Mo_{1.0}
284 kg/ha, T₃: B_{3.0}Mo_{1.5} kg/ha.

285

286 3.8 Curd yield per hectare

287 Curd yield per hectare of cauliflower showed significantly significant differences due to different plant
288 spacing (Table 3). The highest curd yield per hectare (39.89 t/ha) was observed from S₂ treatment,
289 while the lowest curd yield per hectare (35.00 t/ha) was found from S₁ treatment (Table 3). The crops
290 grow in such close spacing yield more though main heads are smaller and these mature slightly later
291 that case optimum spacing is followed. [3, 20] reported the maximum yield of cauliflower where the
292 plants were spaced 45x50 cm apart.

293 Statistically significant variation was recorded for micronutrients in terms of curd yield per hectare of
294 cauliflower (Table 3). The highest curd yield (46.85 t/ha) was found from T₂ treatment which was
295 closely followed by (41.44 t/ha) T₃ treatment, whereas the lowest curd yield (24.41 t/ha) was recorded
296 from T₀ treatment (Table 3). [18] reported that B and Mo application significantly increased curd
297 diameter, weight and yield of cauliflower. [21] reported the better cauliflower curd from the application
298 of boron.

299 Combined effect of different plant spacing and micronutrients showed statistically significant variation
300 on curd yield of cauliflower (Table 4). The highest yield (51.56 t/ha) was recorded from S₂T₂, while the
301 lowest curd yield (20.33 t/ha) was observed from S₁T₀ treatment combination (Table 4). These
302 findings are in confirmation with the findings of [14, 16, 18, 19, 22].

303

304 4. CONCLUSION

305

306 In the experiment plant spacing S₁ (50 cm × 50 cm) treatment gave higher curd yield per plant but
307 plant spacing S₂ (50 cm × 40 cm) treatment gave maximum curd yield per hectare.

Micronutrients combination T₂ (2.0 kg B/ha and 1.0 kg Mo/ha) was more effective than control T₀. Therefore, the results of the investigation suggests that the highest curd yield and good shape cauliflower curd can be obtained, in plant spacing 50 cm x 40 cm with the combined application of B 2.0 kg /ha and Mo 1.0 kg/ha.

REFERENCES

1. Din, F., Qasim M, Elahi N and Faridullah . Responce of different sowing dates on the growth and yield of cauliflower. Sarhad J. Agric.2007; 23(2):289-291.
2. Kannan, D., Kumar, S.D. and Kumar, J.S. Effect of Spacing, Boron and Their Combinations on Yield and Yield Attributing Characters of Cauliflower (*Brassica oleraceae* Var. Botrytis L.). J. Life Sci.2016; 13(3): 524-526.
3. Rahman, M., Iqbal, M., Jilani, M.S. and Waseem, K. Effect of different plant spacing on the production of cauliflower (*Brassica oleraceae* var. Botrytis) under the agro-climatic conditions of D.I. Khan. Pakistan J. Biol. Sci. 2007;10 (24): 4531-4534.
4. Hossain, M.F., Ara, N., Uddin, M.R., Islam, M.R. and Azam, M.G. Effect of sowing date and plant spacing on seed production of cauliflower. Bangladesh J. Agril. Res.,2015; 40(3): 491-500.
5. Moniruzzaman, M., Rahman, S.M.L., Kibria, M.G., Rahman, M.A. and Hossain, M.M.. Effect of boron and nitrogen on yield and hollowstem of broccoli. J. Soil. Nature.2007; 1(3): 24-29.
6. Ningawale, D.K; Singh, R; Bose,U.S;Gurjar,P.S; Anchal Sharma; Gautam, U.S. Effect of boron and molybdenum on growth, yield and quality of cauliflower (*Brassica oleracea* var botrytis) cv. Snowball 16.Indian J. of Agric. Sci.,2016; 86(6):825-829.
7. Gomez, K.A. and Gomez, A.A. Statistical Procedure for Agricultural Research (2nd edn.). Intl. Rice Res. Inst., A Willey Int. Sci.,1984; pp. 28-192.
8. Thakur, O.P., Sharma, P.P. and Sing, K.K. Effect of nitrogen and phosphorus with and without boron on curd yield and stalk rot incidence in cauliflower. Veg. Sci.,1991; 18(2): 115-121.
9. Sharma S K.. Effect of boron and molybdenum on seed production of cauliflower. Indian Journal of Horticulture; 2002; 59(2): 117–80.
10. Singh K P and Rajput C B S. Effect of molybdenum on cauliflower (*Brassica oleracea* var. *botrytis* L.) in sand nutrient culture. Experimental Agriculture;1976; 12(2): 195–199.
11. Muthoo A K Kumar S and Moury A N. Studies on the effect of foliar application of GA₃, NAA and Molybdenum on growth and yield of cauliflower (*Bassica oleracea* var. botrytis L.) cv. Snowball-16. Haryana Journal of Horticulture Science;1987; 16(1-2): 115–20.
12. Rahman A Matiar K M Hossain Monowar S M. and Monjur Hamid M . Effect of sulphur, boron and molybdenum on the growth, curd weight and seed yield of cauliflower. Punjab Vegetable Grower,1992; 27: 11–14.
13. Mengel, K., and Kirkby, E.A. Principles of Plant Nutrition (5th ed.) Dordrecht: Kluwer Academic Publishers.2001; p. 480.
14. Kotur, S.C. Synergistic effect of lime, boron and molybdenum on curd rot and curd yield of cauliflower (*Brassica oleracea* L. var. *botrytis*) on Alfisol. Indian J. Agri. Sci.1998; 68 (5): 268-70.
15. Farag, I.A., Faryhali MA, Refai EF and Shalaby GT. Response of two cultivars of cauliflower to nitrogen rates, boron and molybdenum. Assiut. J. Agri. Sci.1954; 25 (5): 221-33.

- 356 16. Singh, D.N. Effect of boron on the growth and yield of cauliflower in lateritic soil of Western Orissa.
357 Indian J. Hort ; 2003; 60 (3): 283-286.
- 358 17. Chattopadhyay, S.B., and Mukhopadhyay T.P. Effect of foliar application of boron and
359 molybdenum on growth and yield of cauliflower in tarai zone *West Bengal*. Environment and
360 Ecology.2003; 21(4): 955-59.
- 361 18. Kumar S and Choudhary D R. 2002. Effects of FYM, molybdenum and boron application on yield
362 attributes and yield of cauliflower. Crop Research, 2002; 24 (3): 494–6.
- 363 19. Prasad V M and Yadav D. Effect of foliar application of boron and molybdenum on the growth and
364 yield of cauliflower cv. Snowball-16. New Agriculture, 2003; 14(1-2): 121–128.
365
- 366 20. Farzana, L., Solaiman, A.H.M. and Amin, M.R. Potentiality of producing summer cauliflower as
367 influenced by organic manures and spacing. Asian J. Med. Biol. Res., 2016; 2(2): 304-317.
- 368 21. Khadka, Y.G., Rai, S.K. and Raut, S. Effect of Boron on Cauliflower Production. Nepal J. Sci. &
369 Tech., 2005; 6: 103-108.
- 370 22. Mahmud Z U Rahman M M Salam M A Saha S R and Alam M S. Effect of sulphur, boron and
371 molybdenum on curd yield of cauliflower. *Journal of Sub Tropical Agriculture Research and*
372 *Development*, 2005; 3(1): 82–86.
373
- 374 23. Yazici, N., Bilir, N. Aspectual Fertility Variation and Its Effect on Gene Diversity of Seeds in
375 Natural Stands of Taurus Cedar (*Cedrus libani* A. Rich.). *I.J.Genomics*, 2017; 2s960624:1-5.