

An Overview of Development of Quantitative Neurotoxicity Testing *In Vitro*

ABSTRACT

In vitro neurotoxicity testing has been hampered by the fact that the brain architecture is complex. However, a series of innovation of neuroculturing broke through the barrier. The establishment of culturing for the primary neuron and the immobilized neuroblastoma cells enables neurotoxicity testing in vitro. Following to necrotic cell death, extensive morphological changes as seen during neuronal differentiation was used for the endpoints of neurotoxicity. Two-dimensional imaging techniques facilitated quantitative analyses of toxicity of many neurotoxicants. Three-dimensional culturing of neurospheres *in vitro* has been expected to investigate the neurodevelopmental toxicity. The neurosphere assay in vitro also improved the sensitivity to estimate the neurotoxicity. In this paper, I will retrospectively take an overview about the in vitro neurotoxicity testing for rotenone, a dopaminergic pesticide as an environmental toxicant.

Keywords: PC12 cells; NB-1 cells; neural stem cells; neurospheres; rotenone; quantitative neurotoxicity

1. INTRODUCTION

In light of the large number of chemicals, there is a demand to develop rapid screening techniques. Historically, PC12 cells have been used for testing neurotoxicity. The PC 12 cell line was derived from rat pheochromocytoma, a tumor arising from chromaffin cells of the adrenal medulla and developed to study cell differentiation [1]. Neuronal differentiation is a complex process that induces both morphological and biochemical changes. The most obvious things are a decrease in cell proliferation, and the emergence of extending processes. During neuronal differentiation, cells also acquire excitability and start to express some chemical coding genes that provide their functional identity. Upon exposure to neuronal stimulation, PC12 cells gradually exit the mitotic cycle and begin to differentiate, developing axonal projections, electrical excitability, and the characteristics of cholinergic and catecholaminergic neurons. Therefore, the PC12 model enables the detection of environmental toxicants.


Following to rat PC12 model cells, human cell lines such as NB-1 and SH-SY5Y were used to investigate potential species-specific differences, rather than non-human cell origins. It was expected to extrapolate human toxicity.

Recent evidence points to important contributions of exposure to environmental neurotoxicants in the marked increase in neurodevelopmental disorders [2-4]. In response to the need for more efficient methods to identify potential developmental neurotoxicants, neurosphere assay has recently been established. Neural stem cells play an essential role in the development of central nervous system, having self-renewal potency and being multi-

potential. In 1992, it was demonstrated that cells from central nervous system of adult and embryonic mice can be isolated and propagated in culture [5]. In the presence of epidermal growth factor, cell agglomerations, termed neurospheres were formed. They proliferate in culture and have the ability to migrate and differentiate into neurons, astrocytes, and oligodendrocytes. Neurosphere culturing is three-dimensional cell systems and there a valuable in vitro model that mimics basic processes of brain development. Therefore, neurospheres are a useful tool for testing chemicals for their abilities to interfere with these processed: proliferation, migration, differentiation, and apoptosis.

In this paper, we compare a variety of cell-based neurotoxicity testing with several endpoints using rotenone, a dopaminergic pesticide (Fig.1, ref 6-8).

Figure 1. The structure of rotenone and its derived plant



2. MATERIAL AND METHODS

2.1 Culture of PC12 cells

PC12 cells (RCB 0009; RIKEN, Tsukuba, Japan) were grown in Dulbecco's modified Eagle's Medium (Sigma-Aldrich) supplemented with 10% fetal bovine serum (FBS; Life Technologies, Inc., Rockville, MD), 4.5 mg/ml glucose, penicillin (100 U/ml), and streptomycin (100 µg/ml) in a humidified atmosphere of 95% air, 5% CO₂ at 37 °C. The cells were subcultured (1:3) 2 to 3 times per week. Cell viability was determined by trypan blue exclusion method.

2.2 Culture of NB-1 cells

Human neuroblastoma NB-1 cells were cultured in 45% RPMI 1640 and 45% Eagle's minimum essential medium containing 10% FBS (Life Technologies, Inc.), sodium pyruvate, penicillin (100 U/ml), and streptomycin (100 µg/ml) in a humidified atmosphere of 95% air, 5% CO₂ at 37 °C. The cells were subcultured (1:6) once a week. The viable cell number of the NB-1 cells was estimated by crystal violet staining. Fixed and dried cells in a plate were rehydrated with distilled water and photographed under a phase-contrast microscope (DMIRB, Leica Microsystems, Tokyo, Japan) equipped with digital camera. The digital images obtained were then analyzed using image analysis software by counting the cell number and total neurite length in the image field. The degree of neurite extension is represented as the total length of neuritis in micrometer per cell in randomly chosen phase-contrast microscope fields.

2.3 Culture of neurosphere

Pregnant Wistar rats at embryonic day 14 (E14) were obtained from Clea (Tokyo, Japan). The animals were maintained in home cages at 22°C with a 12-h light-dark cycle. They received the MF diet (Oriental Yeast Corp., Tokyo, Japan) and distilled water ad libitum. All animal care procedures were in accordance with National Institute for Environmental Studies guidelines. The rats were sacrificed by diethyl ether overdose on E16. The embryos were removed and transferred to minimal essential medium (MEM; Sigma-Aldrich). Subsequently, the mesencephalons were dissected from the embryos, and were enzymatically digested with 50 U deoxyribonuclease I (Takara Corp., Kyoto, Japan) and 0.8 U papain (Sigma-Aldrich) at 32°C for 12 min. After stirring, the digestion mixture was passed through a 70-µm cell strainer (BD Biosciences). The run-through containing the neural stem cells was centrifuged at 800 x g for 10 min. It was then resuspended in Dulbecco's Modified Eagle's Medium (DMEM) and F12 medium (1:1; Invitrogen, Tokyo, Japan) supplemented with B27 (Invitrogen), 20 ng/ml basic fibroblast growth factor (bFGF; R&D Systems, Inc., MN) and 10 ng/ml epidermal growth factor (EGF; Roche Applied Science, Tokyo, Japan), and cultured in uncoated dishes without serum. Fresh culture medium containing EGF and bFGF was added after 3-4 days.

The neurospheres were seeded in an uncoated glass-bottomed dish (D110300; MATSUNAMI, Tokyo, Japan) in the presence of bFGF and EGF for 3 h, allow cells to adhere. The migrating distance of the cells was statistically measured from the edge of the sphere, using National Institute of Health ImageJ 1.38x software (public domain software).


TUNEL staining was carried out, as described previously [26, 27]. The cells were fixed in 4% paraformaldehyde, washed twice with PBS, and permeabilized in 0.5% Triton X-100 for 5 min on ice. TUNEL labelling was done with fluorescein dUTP (Roche Applied Science, Mannheim, Germany) in the presence of terminal deoxynucleotidyl transferase for 1 h at 37°C. Following labelling, the cells were washed with PBS twice and then directly surveyed under a fluorescence microscope. Images were captured using Viewfinder Lite ver.1.0 camera software through DP-50 digital camera (Olympus, Tokyo, Japan). For quantification of TUNEL-labeled cells, every field containing TUNEL-positive signals was photographed at 100x optical magnification. Then, TUNEL-positive cells were counted.

3. RESULTS

3.1. Cell viability of PC12 cells exposed to rotenone

Rotenone neurotoxicity in PC12 cells was examined by trypan blue exclusion method. PC12 cells were exposed to variety of concentration of rotenone for 3 days. Following fixing the treated cells, a number of cells were counted. Cell viability was decreased in a semilogarithmic-linear, dose-dependent manner. IC₅₀ was about 1 µM (Fig.2).

Figure 2. Rotenone neurotoxicity in rat pheochromocytoma PC12 cells.




119

120 3.1. 1. Inhibition of neurite outgrowth of NB-1 cells exposed to rotenone

121 Rotenone neurotoxicity in NB-1 cells was examined as an endpoint of neurite
 122 outgrowth. NB-1 cells were seeded on a culture plate and rotenone (0~50 μM) was
 123 added for 24h. The treated cells were fixed and the length of neurite outgrowth was
 124 measured by imaging analyses. The length was decreased in a semilogarithmic-
 125 linear, dose-dependent manner. IC₅₀ was about 1 μM (Fig.3).

126 **Figure 3. Neurite outgrowth in rotenone-exposed human neuroblastoma**
 127 **NB1 cells.**



128

129


130 3.1.2. Neurosphere assay for neurodevelopmental toxicity of rotenone

131 We isolated neural stem cells from E15 rat embryos (Fig.4a), using pooled mesencephalons
132 from 12 fetuses. After 2~3 weeks in culture, neurospheres appeared (Fig. 4c), suggesting
133 self-renewal occurred. Neurospheres of about 200 μm in diameter consisted of about 10^3
134 cells.

135

136 **Figure 4.** (a) An E15 rat fetus showing the mesencephalon. (b) Primary neurospheres after
137 7 days *in vitro* (c) Primary neurospheres after 2~3 days *in vitro*. Adapted from ref. [9].


138



139

140 **Figure 5. Identification of cultured mesencephalic neurosphere.** Neurospheres were
141 immunostained with (a) anti-nestin antibody, (b) anti-MAPs antibody, or (c) anti-GFAP
142 antibody. Adapted from ref. [9].

143



144

145

146

147

148

149

150

151 To identify neural stem cells, we stained the neurospheres with an anti-nestin antibody, as
152 shown in Figure 5a. The nestin-positive cells were localized both at the edge and within the
153 spheres. Since neural stem cells are multipotent for neural differentiation, we also
154 immunostained the neurospheres for MAPs, which were located in cells at the edge of the
155 spheres (Fig.5b). Since on E15, when we isolated the neuronal stem cells, rat embryos are
156 undergoing gliogenesis, we stained the neurospheres with anti-GFAP antibody, which mainly
157 stained cells at the sides of the spheres (Fig.5c). Our results suggested that heterogeneous
158 cell populations were present in neurospheres, at late embryonic stages.

During the culture, cells emerged from the plated neurospheres and migrated along the radial axis (Fig. 6 a and b). After 3 h, the plated cells were treated with various concentrations of rotenone (0-10 μM) for 24 h (Figs.7 b-f). The migration distance of the cells was measured from the edge of the neurospheres using NIH ImageJ 1.38x public domain software. Rotenone prevented the cells from migrating from the neurospheres in a linear, dose-dependent manner (Fig. 7g). The half-maximal inhibitory concentration (IC_{50}) was 0.32 μM .

Figure 6. Neural stem cells radically migrated from neurosphere. (a) The early stage, or (b) the later stage of the culturing





Figure 7. Rotenone inhibition of cell migration from neurospheres *in vitro* at various concentrations. (a) No migrating cells during the initial 3h, (b) 0 μM , (c) 0.1 μM , (d) 1 μM , (e) 5 μM , (f) 10 μM . The migration distance was quantitatively measured with NIH ImageJ 1.38x software (g). Adapted from ref. [10].



To compare the toxicity of 2 chemicals, neurosphere assay for bisphenol A, an endocrine disruptor, was carried out under the condition where rotenone was did (Fig.8). The percent inhibition of migration by bisphenol A and rotenone at 1 μM was 35% and 70%, respectively.


211 Thus, the rank order of potency of chemicals was: bisphenol A< rotenone. The value of
212 other endpoints was summarized in Table 1.

213

214 **Figure 8. Comparison of neurotoxicity of 2 chemicals by neurosphere assay *in vitro*.**

215

Modified from ref. 11.



216

217

218

219 **Table 1. Multiple endpoints used in chemical neurosphere testing *In Vitro***

220

221

Modified from refs. 10 and 11.

| Endpoint | Chemical | |
|--------------------------------------|----------|-------------|
| | Rotenone | Bisphenol A |
| 1. Migration (IC ₅₀) | 0.32 μM | ↓ |
| 2. Proliferation (IC ₅₀) | 1.9 μM | 5.0 μM |
| 3. Apoptosis | ↑ | ND |
| 4. Log K _{ow} | 4.10 | 3.32 |

222 ↑:induction; ↓:inhibition; ND: not detected; values of Log K_{ow} were reported in International
223 Chemical Safety Cards (ICSC, Japanese Ver.).

224

225 **Table 2. Sensitivity to rotenone neurotoxicity in various *In Vitro* testing**

| Source | Half maximal effects | Period s | Endpoint | Reference |
|------------------|----------------------|----------|-------------------|------------------|
| Rat PC12 cells | 1 μM < | 24h | Cell viability | 24 |
| | 1 μM | 72h | Cell viability | Unpublished data |
| Human NB-1 cells | 1 μM | 24h | Neurite outgrowth | Unpublished data |

| | | | | |
|---------------------|--------------|-----|----------------|----|
| Human neurosphere | 4 μ M | 24h | Cell viability | 25 |
| Rat E16 neurosphere | 0.32 μ M | 24h | Migration | 10 |
| | 1.9 μ M | 24h | Proliferation | 10 |
| | 1.4 μ M | 24h | Apoptosis | 10 |

4. DISCUSSION

Rotenone is a botanical pesticide [6-8]. Neurotoxic nature of rotenone has been used to produce adult Parkinson model animals due to nigrostriatal dopaminergic lesions [12-14]. Upon the working hypothesis of the developmental origins of health and disease (DOHaD; ref.15), it is important to examine the neurodevelopmental toxicity of rotenone in neurosphere assays, comparing other cell lines. The hypothesis suggests that the environmental origins of human sporadic Parkinson disease occur early in life. One possible explanation for this phenomenon is that early exposure to neurotoxic chemicals reduces the number of dopaminergic neurons in the substantia nigra to levels below those needed to sustain normal function during the course of the neuronal attrition associated with aging.

There are many endpoints to evaluate neurotoxicity [Table 2]. Ultimate endpoint for neurotoxicity is cell death. Therefore, neuronal cell viability was used for evaluating its toxicity. Catastrophic cell death by neurotoxins is observed as perturbation of energy producing systems, cellular membrane defects, and increased influx of calcium ions. This is concomitant with necrotic action of the toxin. For more sensitive detection of the toxicity, a new biochemical marker is needed.

About 20 years ago, apoptotic nature of environmental toxins were discovered in several types of cells, including renal and neuronal cell [16,17]. Apoptosis is a programmed form of cell death mediating precisely controlled deletions of 'unwanted' cells. This phenomenon is initiated not only by physiological stimuli but also by an extensive array of nonphysiological agents. The characteristics of apoptosis are DNA fragmentation and chromatin condensation, in which are endpoints of toxicants as earlier phase than necrotic phase of toxins.

Development of techniques for Image acquisition enabled to open a new way to evaluate neurotoxicity: neurite outgrowth was also used for endpoint of neurotoxicity [18,19]. The growth of axonal and dendritic processes during brain development is a critical determinant of neural connectivity, and disruption of this process could lead to neuronal dysfunction. Neurite outgrowth can be recapitulated in vitro using a variety of cell models. These models have been become valuable tools for investigating the mechanism for known developmental neurotoxicants.

The developing human brain can be more susceptible to injury caused by toxic agents than the brain of an adult. Probably all potential neurotoxic compounds would also cause damage to the developing brain and at much lower doses. Indeed, neurodevelopmental disorders in children such as attention deficit disorder or autism have been associated with the exposure to chemicals in the environment during early fetal development [20].

Particularly, it has been suggested that one of possibility of autism-spectrum disorders are initiated in the embryonic neural stem cells [21]. Neural stem cells play an essential role in the development of central nervous system, having self-renewal potency and being

266 multipotential: they are able to differentiate to neurons, astrocytes and oligodendrocytes to
267 form neuronal architecture. Indeed, in the culture of neural stem cells, they form free-
268 floating three-dimensional structures. Therefore, application of neurosphere for
269 neurodevelopmental toxicity testing is reasonable.

270 Quantitative analyses in this study revealed the linearity in function as migration inhibition
271 versus chemical concentration. Owing to the linearity of the functional relationship between
272 the migration inhibition and the concentration of test chemicals, this approach could be
273 employed as a reliable quantitative assay system, excluding the issue of nonlinearity in low
274 dose of an endocrine disruptor such as bisphenol A [22,23].
275

276 5. CONCLUSION

277
278 In this paper, I take an overview about neurotoxicity of rotenone, investigating with rat PC12
279 cell, human NB-1 cells and rat embryonic neural stem cells. Furthermore, quantitative
280 analysis revealed a linear function between the cellular endpoints and the rotenone
281 concentration. This could be employed as a simple and rapid screening for neurotoxicity of
282 environmental chemicals. Particularly, neurosphere assay would be hoped to develop the
283 risk assessment methods for chemicals based on infant physiology.
284
285

286 REFERENCES

- 287
288
289 1. Greene, L.A. Tischler, A.S. Establishment of a noradrenergic clonal line of rat
290 adrenal pheochromocytoma cells which respond to nerve growth factor. Proc Natl Acad Sci
291 U.S.A.1976, 73, 2424-2428.
- 292 2. Jacobson, J.L. Jacobson, S.W. Intellectual impairment in children exposed to
293 polychlorinated bisphenyls in utero. N. Engl. J. Med. 1996, 335, 783-789.
- 294 3. Patandin, S., Lanting, C.I., Mulder, P.G., Boersma, E.R., Sauer, P.J., Weisglas-
295 Kuperus, N. J. Pediatr. 1999, 134, 33-41.
- 296 4. Rice, D.C. Parallels between attention deficit hyperactivity disorder and behavioral
297 deficits produced by neurotoxic exposure in monkeys. Environ. Health Perspect. 2000,
298 108(Suppl.3), 511-533.
- 299 5. Reynolds, B.A., Tetzlaff, W., Weiss, S. A multipotent EGF-responsive striatal
300 embryonic progenitor cell produces neurons and astrocytes. J. Neurosci. 1992, 12, 4565-
301 4574.
- 302 6. Ray, D.E. Pesticides derived from plants and living organisms. In Handbook of
303 Pesticide Toxicology, Hayes, W.J. and Laws E.R. Eds.; Academic Press, San Diego, US,
304 1991; pp585-636.
- 305 7. Ujvary, I. Pest control agents from natural products. In Handbook of Pesticide
306 Toxicology, 2nd ed.; Krieger, R. Ed.; Academic Press, California, US, 2001; pp109-180.
- 307 8. Ott, K.C. Rotenone. A brief review of its chemistry, environmental fate, and the
308 toxicity of rotenone formulations. New Mexico Council of Trout Unlimited. 2008; Available at:
309 <http://www.newmexicotu.org/Rotenone%20summary.pdf>.

- 310 9. Ishido M. Effects of *p*-nitrotoluene on cultured mesencephalic neural stem cells. J.
311 Health Sci (Tokyo) 2009; 55, 114-118.
- 312 10. Ishido, M., Suzuki J. Inhibition by rotenone of mesencephalic neural stem-cell
313 migration in a neurosphere assay in vitro. Toxicol in Vitro 2010; 24, 552-557.
- 314 11. Ishido, M., Suzuki, J. Quantitative analyses of inhibitory effects of bisphenol A on
315 neural stem-cell migration using a neurosphere assay in vitro. J. Health Sci.(Tokyo) 2010;
316 56, 175-181.
- 317 12. Betarbet, R., Sherer, T.B., MacKenzie, G., Garcia-Osuna, M., Panov, A.V.,
318 Greenamyre, J.T. Chronic systemic pesticide exposure reproduces features of Parkinson's
319 disease. Nat. Neurosci. 2000, 3, 1301-1306.
- 320 13. Dauer, W., Przedborski, S. Parkinson's disease: mechanisms and models. Neuron
321 2003, 39; 889-909.
- 322 14. Blesa, J., Phani, S., Jackson-Lewis, V., Przedborski, S. Classic and new animal
323 models of Parkinson's disease. J. Biomed. Biotech. 2012, doi:10.1155/2012/845618
- 324 15. Gilman, M.W., Barker, D., Bier, D., Cagampang, F., Challs, J., Fall, C., Godfrey, K.,
325 Gluckman, P., Hanson, M., Kuh, D., Nathanielsz, P., Nestel, P., Thornburg, K.L. Meeting
326 report on the 3rd international congress on developmental origins of health and disease
327 (DOHaD). Pediatr. Res. 2007; 61, 625-629.
- 328 16. Ishido, M. et al. Cadmium-induced DNA fragmentation is inhibitable by zinc in
329 porcine kidney LLC-PK₁ cells. Life Sci. 1995, 17, 351-356.
- 330 17. Kunimoto, M. Methylmercury induces apoptosis of rat cerebellar neurons in primary
331 culture. Biochem. Biophys. Res. Commun. 1994, 204, 310-317.
- 332 18. Pramanik, R., Ishido, M., and Kunimoto, M. Methylmercury-mediated down-
333 regulation of mtHSP70 and phospholipase A2 mRNA expression in human neuroblastoma
334 NB-1 cells identified by cDNA macroarray analysis. J. Health Sci.(Tokyo) 2002, 48, 381-
335 384.
- 336 19. Pramanik, R., Ishido, M., and Kunimoto, M. Effects of cadmium chloride on neurite
337 outgrowth and gene expression in human neuroblastoma NB-1 cells. J. Health Sci.(Tokyo)
338 2001, 47; 478-482.
- 339 20. Sai, Y., Wu, Q., Le, W., Ye, F., Dong, Z. Rotenone-induced PC12 cell toxicity is
340 caused by oxidative stress resulting from altered dopamine metabolism. Toxicology in Vitro
341 2008, 22; 1461-1468.
- 342 21. Li, J., Spletter, M.L., Johnson, D.A., Wright, L.S., Svendsen, C.N., Johnson, J.A.
343 Rotenone-induced caspase 9/3-independent and -dependent cell death in undifferentiated
344 and differentiated human neural stem cells. J. Neurochem. 2005; 92, 462-476.
- 345 22. Colborn, T. Neurodevelopment and endocrine disruption. Environ Health Perspect.
346 2004, 115, 924-931.
- 347 23. Li, H., Radford, J.C., Ragusa, M.J., Shea, K.L., McKercher, S.R., Zaremba, J.D.,
348 Soussou, W., Nie, Z., Kang, Y. J., Nakanishi, N., Okamoto, S., Roberts, A. J., Schwarz, J. J.,

349 Lipton, S. A. Transcription factor MEF2C influences neural stem/progenitor cell
350 differentiation and maturation in vivo. *Proc. Natl. Acad. Sci. U.S.A* 2008, 105, 9397-9402.

351 24. Welshons, W.V., Thayer, K.A., Judt, B.M., Taylor, J.A., Curran, E.M., vom Saal, F.S.
352 Large effects from small exposure. *Environ. Health Perspect.* 2003, 111; 994-1006.

353 25. vom Saal, F.S., Hughes, C. An extensive new literature concerning low-dose effects
354 of bisphenol A shows the needs for a new risk assessment. *Environ. Health Perspect.* 2005,
355 113; 926-933.

356 26. Ishido, M. Melatonin inhibits maneb-induced aggregation of α -synuclein in rat
357 pheochromocytoma cells. *J. Pineal Res.* 2007, 42:125-130.

358 27. Ishido, M. Activation of STAT3 by pituitary adenylate cyclase-Activating polypeptide
359 (PACAP) during PACAP-promoted neurite outgrowth of PC12 cells. *J. Mol. Neurosci.* 2010,
360 42; 349-358.

361

362

363 **DEFINITIONS, ACRONYMS, ABBREVIATIONS**

364 Here is the Definitions section. This is an optional section.

365 **Term:** Definition for the term

366

367 **APPENDIX**

368