

Original Research Article

The potential ameliorative effects of *Annona muricata* (Linn) on Sodium Fluoride-Induced Toxicity on Haematological indices and Fecundity of Adult Male Wistar Rats.

Abstract

Aim: Ameliorative potentials of *Annona muricata* (Linn) on Sodium fluoride-induced toxicity on haematology indices and fecundity of adult male Wistar rats. **Methods:** Eighty-five (85) adult male Wistar rats were divided into 17 groups of 5 rats each. NaF (10 mg/kg) + fruit juice, ethanol stem bark, and leaf extracts of *A. muricata* at five different doses of 500, 1000, 1500, 2000 and 2500 mg/kg body weight were administered to the rats for 6 weeks. Blood samples were taken after 6 weeks through the ocular puncture and the sera were used for testosterone, follicle stimulating hormone (FSH), luteinizing hormone (LH) Triiodothyronine (T3), Thyroxine (T4), and Thyroid Stimulating Hormone (TSH) tests, while whole blood was used for haematological parameters such as haemoglobin (Hb), packed cell volume (PCV), total white blood cell, platelets count, lymphocytes (%) and neutrophils (%). The testes and epididymis of the rats were harvested for histological studies and sperm analysis such as sperm motility, viability, count and sperm head abnormality. **Results:** Administration of NaF + fruit juice, NaF + Stem bark and NaF + leaf extracts caused an increase ($p < 0.05$) in epididymal sperm count, sperm motility, and live spermatozoa along with a simultaneous decrease in dead spermatozoa as compared to the rats of the group treated with NaF alone. Result also showed that treatment with doses above 500 mg/kg body weight of NaF + fruit juice, NaF + stem bark and leaf extracts produced significant increases ($p < 0.05$) in Hb concentration and PCV when compared with group 2 rats. Similarly, groups treated with NaF + 2000 mg/kg of fruit juice and NaF + 1000 mg/kg of stem bark, and leaf extracts, showed a significant increase ($p < 0.05$) in platelet count when compared with groups 1 and 2 rats. Histological examination showed that NaF treatment brought about severe testicular damage while treatment with the extracts ameliorated this effect. **Conclusion:** *A. muricata* fruit juice and extracts were found to increase testosterone concentration, thus validating its ameliorative potential in NaF-induced toxicity.

Keywords: Sodium fluoride, *Annona muricata* (Linn), fecundity, haematology, hormones, histology.

Introduction

Several clinical investigations and animal experiments suggest that fluoride has adverse impacts on male reproductive function producing structural and functional defects in spermatozoa, a decrease in sperm count, disturbances in the levels of reproductive hormones and reduced fertility [1, 2]. Spermatozoa undergo various processes to ultimately fertilize an oocyte, including spermatogenesis, capacitation, and the acrosome reaction. Fluoride has been shown to impair all three of these processes [3]. *In vitro* fluoride exposure at high concentrations affected certain signal pathways, such as inhibition of the cell cycle, apoptosis and proliferation [4]. Thyroid hormone disruption caused by fluoride results in abnormal function and development of testes, lowering libido, reducing sex hormones, interferes directly and indirectly with spermatogenesis, influencing steroid hormone receptors, inducing oxidative stress in testes. However, the most important mechanism by which fluoride reduces the level of testosterone is interference with steroidogenesis in the

45 Leydig cells. This interference has been demonstrated in several studies in which activity
46 levels of testicular steroidogenic marker enzymes 3β -hydroxysteroid dehydrogenase (3β
47 HSD) and 17β -hydroxysteroid dehydrogenase (17-HSD) decreased significantly in NaF-
48 treated rats [5].

49 *Annona muricata* fruit juice has been shown to possess antibacterial, antifungal,
50 anticancerous, antimalarial, antidiabetic, hepatoprotective, anti-inflammatory, hypotensive
51 and immune enhancing effect [6]. Phytochemical screening of *A. muricata* leaf ethanolic
52 extract shows the presence of saponins, triterpenoids, flavonoids, tannins, alkaloids, and
53 cardiac glycosides [7]. *A. muricata* leaf extract is believed to stabilize blood sugar level in a
54 normal range that is very useful for diabetic management [8]. Several types of research have
55 shown that *A. muricata* leaf has hypoglycemic activity and revealed regeneration of
56 pancreatic islet [9, 10, 11]. The ethanol leaf extract of *A. muricata* also is known to reduce
57 serum uric acid level [12], contain essential oils with parasiticidal, antibacterial, antidiarrheal,
58 rheumatological and antineuralgic properties [13, 14, 15]. The extract from *A.*
59 *muricata* induced necrosis of pancreatic cancer cells by inhibiting cellular metabolism [16]. *A.*
60 *muricata* leaf extract may possess anticancer properties by enhancing caspase-3 activity
61 which is a pro-apoptosis marker [17]. The use of different parts of *A. muricata* for the
62 treatment of these pathological disorders suggests it may possess anti-toxic properties and
63 stimulated our interest to study its ameliorative effect on NaF- induced toxicity on
64 haematology indices and fecundity of adult male Wistar rats.

65 **Materials and Methods**

66 **2.1 Animals**

67 Male adult albino rats (150-250 g) were obtained from the animal house of College of
68 Medicine, University of Nigeria, Enugu Campus. The animals were housed in steel cages
69 within the Laboratory Animals Facility of Brain-Phosphorylation Scientific Solution
70 Services, No9, Ogui Road Enugu, Enugu State, 5th Floor, Right Wing, maintained and given
71 standard feed and clean drinking water *ad libitum*. They were allowed to acclimatize for a
72 period of four weeks before use. All animal experiments were in compliance with the
73 National Institute of Health Guide for care and use of laboratory animal.

74 **Collection and Extraction of Plant Materials and Fruit Juice.**

75 Fresh stem bark, leaf, and fruits of *Annona muricata* were collected from Abua, Rivers State,
76 in March 2017. The stem bark and the leaf were cut to pieces, dried under room
77 temperature, ground and pulverized to a coarse powder using a Hammer mill (Gallenkamp,
78 U.S.A.). The plant materials were identified and authenticated by Mr. Alfred Ozioko of
79 International Centre for Ethnomedicines and Drug Development Nsukka, Nigeria and
80 deposited in herbarium with Voucher Number: Intercedd/16091. Known quantities (1.851kg)
81 of the dried stem bark powder and 1.016 kg of the dried leaf powder were extracted with
82 analytical grade ethanol using maceration method for 48 hours. The mixture was vacuum-
83 filtered through Whatman No 1 filter paper and concentrated using a vacuum rotary
84 evaporator (Eyla N-1000, Japan) to afford 97.352 g (5.257 % w/w) for stem bark extract and
85 126.312 g (12.432 % w/w) for leaf extract. The extractive yield was calculated using the
86 relation: Yield (%) = [Weight of extract (g)/Weight of plant material (g)]*100. The fruit juices
87 were used raw without concentrating it. The epicarps and the seeds of the ripe fruits were
88 removed with hand and the mesocarps were sliced with knife into small sizes and ground
89 with and an electric grinder into paste form. This was further sieved with a muslin cloth to

90 remove the fibres. The filtrate was transferred into clean glass container, sealed and
91 preserved in refrigerator at -10°C until use.

92 **Experimental Design**

93 Eighty five sexually matured male adult albino rats (150-250 g) were divided into 17 groups
94 of 5 rats each, according to their average weight, and received daily oral dose of the
95 treatment as follows:

- 96 Group 1: Normal feed and water (positive control)
- 97 Group 2: NaF (10mg/kg) (negative control)
- 98 Group 3: NaF (10mg/kg) + Fruit Juice Extract (500mg/kg)
- 99 Group 4: NaF (10mg/kg) + Fruit Juice Extract (1000mg/kg)
- 100 Group 5: NaF (10mg/kg) + Fruit Juice Extract (1500mg/kg)
- 101 Group 6: NaF (10mg/kg) + Fruit Juice Extract (2000mg/kg)
- 102 Group 7: NaF (10mg/kg) + Fruit Juice Extract (2500mg/kg)
- 103 Group 8: NaF (10mg/kg) + Leaf Extract (500mg/kg)
- 104 Group 9: NaF (10mg/kg) + Leaf Extract (1000mg/kg)
- 105 Group 10: NaF (10mg/kg) + Leaf Extract (1500mg/kg)
- 106 Group 11: NaF (10mg/kg) + Leaf Extract (2000mg/kg)
- 107 Group 12: NaF (10mg/kg) + Leaf Extract (2500mg/kg)
- 108 Group 13: NaF (10mg/kg) + Stem Bark Extract (500mg/kg)
- 109 Group 14: NaF (10mg/kg) + Stem Bark Extract (1000mg/kg)
- 110 Group 15: NaF (10mg/kg) + Stem Bark Extract (1500mg/kg)
- 111 Group 16: NaF (10mg/kg) + Stem Bark Extract (2000mg/kg)
- 112 Group 17: NaF (10mg/kg) + Stem Bark Extract (2500mg/kg)

113 Blood was taken after the 6th week of administration through **the** ocular puncture. Two ml of
114 the blood samples from each group (n=4) were collected in test tubes and put into centrifuge
115 tubes, spun at 3000 rpm for 10 min and the serum collected for hormonal assays which
116 include: testosterone, follicle stimulating hormone (FSH), luteinizing hormone (LH),
117 Triiodothyronine (T3), Thyroxine (T4) hormone and Thyroid stimulating hormone. Whole
118 **blood** (2 ml) for haematological studies were placed in EDTA tubes and assayed for full
119 blood count. The rats were sacrificed under chloroform anaesthesia after collection of blood
120 samples. **The testes and epididymis** were dissected out and rapidly fixed in buffered neutral
121 formalin (10%) for histological studies. The epididymis was processed for epididymal sperm
122 motility, viability, count and sperm head abnormality.

123 **Histopathological examination**

124 The tissues were subjected to standard routine histological procedures [18]. The slides were
125 viewed using the light microscope and histopathological changes were observed and
126 recorded at x400 magnification identifying both the normal and atrophied seminiferous
127 tubules and spermatocytes.

128 **2.3.2 Haematological studies**

129 **Determination of haematological parameters**

130 Determination of haematological parameters such as haemoglobin concentration (Hb),
131 packed cell volume (PCV), total white blood cell count (TWBC), platelet count, neutrophils
132 and lymphocytes) were done using standard operative procedures[19].

133 **Hormonal Assay**

134 Plasma Testosterone, Follicle-stimulating and Luteinizing hormones were determined by
135 fluorescence immunoassay (FIA) methods with commercial kits (Boditech Med Incorporated,
136 Republic of Korea), using the ichroma machine (Boditech: BOD13303, Korea).

137 **Sperm Analyses**

138 **Semen pH and sperm motility**

139 Immediately after dissection, a puncture was made in the epididymis with a sterile pin. The
140 semen smeared on the pin was rubbed on a pH paper of range 1.0-10.0. The colour change
141 corresponds to the pH and was read from the paper. The dissected epididymis was
142 measured and sliced into small pieces with a sterilized surgical blade and finally introduced
143 into a beaker. The epididymal sperm samples were obtained by macerating known weight
144 (100 mg) of cauda epididymis in physiological saline in the ratio of 1:10 weight by volume.
145 After vigorous shaking, two drops of sperm suspension was put on a microscope slide and a
146 coverslip was placed. The numbers of progressively motile sperm cells were counted under
147 $\times 40$ lenses.

$$148 \quad \% \text{ Motility} = \frac{\text{No of motile spermatozoa}}{\text{Total no of spermatozoa counted}} \times 100$$

150 **Percentage dead sperm cells**

151 The percentage of dead sperm cells was determined using "Eosin-Nigrosin one-step staining
152 technique" [20]. A portion of the sperm suspension was mixed with equal volume of Eosin-
153 Nigrosin stain and two (2) air-dried smears were prepared on glass slides for each sample.
154 Dead sperm cells took up the stain and appeared pinkish. Percentage of dead sperm cells
155 were calculated based on the number of dead sperm cells out of the total number of sperm
156 cells observed.

157 **Sperm viability**

158 The sperm viability test was determined using "Eosin-Nigrosin one-step staining technique"
159 [20]. A portion of the sperm suspension was mixed with equal volume of Eosin-Nigrosin
160 stain and two (2) air-dried smears were prepared on glass slides for each sample. The slides
161 were examined for percentage viability. Normal live sperm cells excluded the stain and
162 appeared whitish, whereas dead sperm cells took up the stain and appeared pinkish.
163 Percentage viability was calculated based on the number of live sperm cells out of the total
164 number of sperm cells counted.

$$165 \quad \text{Sperm viability count} = \frac{\text{Life cell (viable cells)}}{\text{Total cells (both dead \& alive)}} \times \frac{100}{1}$$

167 **Sperm count**

168 The dissected epididymis was measured and sliced into small pieces with a sterilized
 169 surgical blade and finally introduced into a beaker. The epididymal sperm samples were
 170 obtained by macerating this known weight of cauda epididymis in physiological saline in the
 171 ratio of 1:10 weight by volume. After vigorous shaking, two drops of sperm suspension was
 172 put on a microscope slide and a cover slip was placed. Epididymal sperm count was
 173 obtained by cytometry using the improved Neubauer cytometer and was expressed as
 174 million/ml of suspension [21].

175 **Sperm head abnormality test**

176 A known volume of the sperm suspension was mixed with 1% eosin solution (10:1) for 30
 177 min and air-dried smears were prepared on glass slides for each sample. The percentage of
 178 sperm head abnormality was calculated accordingly [21].

179 **Statistical analysis**

180 The data were analysed by (SPSS version 17.5, SPSS Inc.). Significant differences between
 181 means were determined by One-way ANOVA and regarded significant at $p < 0.05$. Results
 182 were presented as Mean \pm Standard Deviation

183 **Results**

184 **Haemoglobin Concentration, Packed Cell Volume, and Platelets Count**

185 Effects of fruit juice, ethanol extracts of stem bark and leaf of *A. muricata* on haemoglobin
 186 concentration of NaF-induced toxicity and fecundity profile of adult male Wistar rats is shown
 187 in Table 3.1. It was observed that NaF at a dose of 10 mg/kg caused a significant decrease
 188 ($p < 0.05$) in Hb concentration, percentage PCV and platelets count in the adult male rats
 189 when compared with the control. However, treatment with doses above 500 mg/kg body
 190 weight of NaF + fruit juice, NaF + stem bark and leaf extracts produced significant increases
 191 in Hb concentration and percentage PCV values when compared with group 2 rats. Similarly,
 192 groups treated with NaF + 2000 mg/kg of fruit juice and NaF + 1000 mg/kg of stem bark, and
 193 leaf extracts, showed a significant increase ($p < 0.05$) in platelet count in comparison with
 194 both the group treated with NaF alone and control group.

195 **Table 3.1: Haemoglobin concentration, percentage packed cell volume and platelet**
 196 **count of control and test groups of NaF-induced toxicity and fecundity profile of adult**
 197 **male Wistar rats after 6 weeks of treatment.**

Group	Hb Concentration (g/dl)	PCV (%)	Platelet count ($\times 10^3 \text{mm}^3$)
Control	9.13 \pm 0.42	25.67 \pm 3.79	118.00 \pm 7.00
NaF	5.87 \pm 0.29 ^a	18.33 \pm 0.58 ^a	112.00 \pm 2.65
Stem bark extract			
10mg/kg NaF + 500	6.87 \pm 0.23	17.47 \pm 13.05 ^a	96.00 \pm 2.59 ^a
10mg/kg NaF + 1000	9.00 \pm 0.00 ^b	24.00 \pm 3.00 ^b	145.50 \pm 5.50 ^{ab}
10mg/kg NaF + 1500	8.27 \pm 1.33 ^b	25.67 \pm 2.08 ^b	128.33 \pm 2.88
10mg/kg NaF + 2000	8.20 \pm 3.20 ^b	24.00 \pm 1.00 ^b	131.33 \pm 3.21 ^{ab}
10mg/kg NaF + 2500	8.87 \pm 1.50 ^b	24.33 \pm 4.93 ^b	140.33 \pm 4.16 ^{ab}

Leaf extract

10mg/kg NaF + 500	7.87±1.16	21.00±1.00	143.00±6.08 ^{αβ}
10mg/kg NaF + 1000	9.00±0.00 ^β	26.33±0.58 ^β	138.33±1.04 ^β
10mg/kg NaF +1500	7.93±0.91 ^β	23.33±3.51 ^β	126.00±1.05 ^β
10mg/kg NaF + 2000	7.80±1.14	23.67±3.06 ^β	128.33±1.89 ^β
10mg/kg NaF + 2500	7.65±0.35	23.00±1.00	132.50±1.75 ^{αβ}

Fruit Juice

10mg/kg NaF + 500	9.25±0.75 ^β	28.50±1.50 ^β	129.50±5.00
10mg/kg NaF + 1000	8.23±1.63 ^β	24.33±4.51 ^β	137.00±1.92 ^β
10mg/kg NaF +1500	8.73±0.55 ^β	26.67±1.53 ^β	128.33±1.04 ^β
10mg/kg NaF + 2000	8.70±0.36 ^β	25.00±1.00 ^β	147.33±6.43 ^{αβ}
10mg/kg NaF + 2500	9.10±0.10 ^β	27.00±0.00 ^β	140.00±1.00 ^{αβ}

198 Results are expressed as Mean±SD; n=4

199 The mean values with ^β as superscripts across the column compared with group treated
 200 with NaF alone are considered significant (p<0.05). The mean values with ^α as superscripts
 201 across the column compared with control group fed with water and feed only are considered
 202 significant (p<0.05).

203 Total White Blood Cell, Neutrophil and Lymphocyte Count

204 Table 3.2 shows the results of the effect of fruit juice, ethanol extracts of stem bark and leaf
 205 of *A. muricata* on total white blood cell of NaF-induced toxicity on fertility profile of adult male
 206 rats. There was no significant difference in total white blood cell, Neutrophil and lymphocyte
 207 count in the group treated with 10 mg/kg of NaF when compared with the control group fed
 208 with water and rat feed only. Similar, there was no significance difference in the total white
 209 blood cell, Neutrophil and lymphocyte count in the groups treated with different
 210 concentrations of the extracts when compared with the controls.

211 **Table 3.2: Total White Blood Cell, Neutrophil and Lymphocyte Count of control and**
 212 **test groups of NaF- induced toxicity and fecundity profile of adult male Wistar after 6**
 213 **weeks of treatment**

Group	WBC (x10 ³ mm ³)	Neutrophil (%)	Lymphocyte (%)
Control	9.33±1.15	44.67±4.51	55.33±4.51
NaF	10.20±7.21	51.43±7.51	49.57±7.51
Stem bark extract			
10mg/kg NaF + 500	09.47±8.39	44.00±5.20	55.67±4.93
10mg/kg NaF + 1000	07.00±5.00	41.00±1.00	58.50±1.50
10mg/kg NaF +1500	08.67±1.76	37.67±2.08	61.00±1.00
10mg/kg NaF + 2000	08.80±1.21	43.67±5.51	56.33±5.51

10mg/kg NaF + 2500	08.50±1.32	36.00±3.31	64.00±4.00
Leaf extract			
10mg/kg NaF + 500	10.10±1.00	39.67±17.62	58.00±14.00
10mg/kg NaF + 1000	11.33±5.77	39.00±1.00	61.00±1.00
10mg/kg NaF + 1500	11.000±1.00	37.67±2.52	62.00±2.65
10mg/kg NaF + 2000	10.00±1.06	53.33±14.15	46.00±13.86
10mg/kg NaF + 2500	53.50±4.25 ^{αβ}	39.00±0.00	60.50±0.50
Fruit Juice			
10mg/kg NaF + 500	9.20±2.20	47.50±2.50	54.50±4.50
10mg/kg NaF + 1000	10.93±1.05	47.33±15.31	49.33±13.58
10mg/kg NaF + 1500	6.57±2.03	46.00±5.29	54.00±5.29
10mg/kg NaF + 2000	9.73±2.31	37.00±2.65	62.67±2.52
10mg/kg NaF + 2500	05.90±3.50	42.50±1.50	67.00±1.00

214 Results are expressed as Mean±SD; n=4

215 The mean values with β as superscripts across the column compared with group treated
 216 with NaF alone are considered significant ($p<0.05$). The mean values with α as superscripts
 217 across the column compared with

218 Testosterone, FSH and LH Concentration

219 Effects of fruit juice, ethanol extracts of stem bark and leaf of *A. muricata* on Testosterone
 220 FSH and LH Concentration, of NaF-induced toxicity on fecundity profile of adult male Wistar
 221 rats shows that at a dose of 10 mg/kg, NaF caused a non-significant decrease ($p>0.05$) in
 222 the serum testosterone and LH concentration, and a significant decrease ($p<0.05$) in the
 223 FSH when compared with normal control group fed with water and feed only (Table 3.3).
 224 Groups treated with NaF + fruit juice, NaF + 2000 and 2500 mg/kg stem bark and Leaf
 225 extracts showed significant increase ($p<0.05$) in serum testosterone concentrations when
 226 compared with both group treated with NaF alone and control group. Groups treated with
 227 NaF + fruit juice, NaF + Leaf extracts showed significant decrease in serum FSH
 228 concentration, when compared with group tested with NaF alone. However, only the group
 229 treated with NaF + 500 mg/kg stem bark extract showed a significant increase ($p<0.05$) in
 230 serum FSH concentration, when compared with group treated with NaF alone. On the other
 231 hand, groups treated with NaF + 1000 and 2500 mg/kg fruit juice and NaF + 500 and 1000
 232 mg/kg leaf extracts exhibited significant decrease ($p<0.05$) in serum LH concentration when
 233 compared with the control group.

234

235

236 **Table 3.3: Testosterone, Follicle stimulating hormone (FSH) and Luteinizing hormone**
 237 **concentration of control and test groups of NaF-induced toxicity and fecundity profile**
 238 **of adult male Wistar after 6 weeks of treatment**

Group	Testosterone (ng/ml)	FSH (MI μ /l)	LH (MI μ /l)
Control	0.827 \pm 0.06	3.54 \pm 2.62	1.93 \pm 0.05
NaF	0.740 \pm 0.06	2.83 \pm 0.79 ^{α}	1.63 \pm 0.62
Stem bark extract			
10mg/kg NaF + 500	0.760 \pm 0.01	5.00 \pm 1.00 ^{β}	1.63 \pm 0.08
10mg/kg NaF + 1000	0.920 \pm 0.11 ^{β}	2.50 \pm 2.33	1.84 \pm 0.50
10mg/kg NaF + 1500	0.833 \pm 0.22	1.87 \pm 0.25	1.57 \pm 0.33
10mg/kg NaF + 2000	2.093 \pm 0.37 ^{$\alpha\beta$}	1.61 \pm 1.12 ^{α}	1.42 \pm 0.55
10mg/kg NaF + 2500	1.637 \pm 0.65 ^{$\alpha\beta$}	1.52 \pm 0.50 ^{α}	1.56 \pm 0.65
Leaf extract			
10mg/kg NaF + 500	0.633 \pm 0.13	1.05 \pm 0.03 ^{α}	0.80 \pm 0.03 ^{$\alpha\beta$}
10mg/kg NaF + 1000	0.817 \pm 0.03	1.05 \pm 0.13 ^{α}	1.04 \pm 0.22 ^{α}
10mg/kg NaF + 1500	0.863 \pm 0.22	1.20 \pm 0.21 ^{α}	1.13 \pm 0.06
10mg/kg NaF + 2000	0.953 \pm 0.13 ^{$\alpha\beta$}	1.93 \pm 0.29	1.63 \pm 0.46
10mg/kg NaF + 2500	1.850 \pm 0.84 ^{$\alpha\beta$}	1.33 \pm 0.50 ^{α}	1.85 \pm 0.11
Fruit Juice			
10mg/kg NaF + 500	1.653 \pm 0.70 ^{$\alpha\beta$}	1.41 \pm 0.42 ^{α}	1.45 \pm 0.84
10mg/kg NaF + 1000	1.900 \pm 0.10 ^{$\alpha\beta$}	1.38 \pm 0.50 ^{α}	0.83 \pm 0.24 ^{α}
10mg/kg NaF + 1500	1.213 \pm 0.27 ^{$\alpha\beta$}	1.83 \pm 0.16	1.96 \pm 0.01 ^{β}
10mg/kg NaF + 2000	2.403 \pm 0.57 ^{$\alpha\beta$}	0.97 \pm 0.03 ^{α}	1.34 \pm 0.58
10mg/kg NaF + 2500	1.653 \pm 0.22 ^{$\alpha\beta$}	1.66 \pm 0.13 ^{α}	0.84 \pm 0.05 ^{α}

239 Results are expressed as Mean \pm SD; n=4

240 The mean values with β as superscripts across the column compared with group treated
 241 with NaF alone are considered significant ($p < 0.05$). The mean values with α as superscripts
 242 across the column compared with control group fed with water and feed only are considered
 243 significant ($p < 0.05$).

244 **Thyroid Hormones**

245 Results in table 3.4 showed the ameliorative potential of fruit juice, ethanol extracts of stem
 246 bark and leaf of *A. muricata* on thyroid hormone concentration of NaF-induced toxicity on
 247 fertility profile of adult male Wistar rats. At a dose of 10 mg/kg, NaF, caused a non-significant
 248 ($p > 0.05$) increase in T_3 , non-significantly reduction ($p > 0.05$) in T_4 and a non-significant
 249 increase ($p > 0.05$) in TSH concentration respectively, when compared with normal control
 250 group. However, concomitant administration of NaF+2000 mg/kg of fruit juice, NaF+2000
 251 mg/kg of stem bark and NaF+2500 mg/kg of leaf extracts exhibited significant increase
 252 ($p < 0.05$) in serum thyroxine concentration when compared with group treated with NaF

253 alone and the control group. Similarly, groups treated with NaF + 2500 mg/kg fruit juice, NaF
 254 + 2500g/kg stem bark and NaF +500 - 2500 mg/kg leaf extracts exhibited significant
 255 increases ($p < 0.05$) in TSH concentration when compared with both the group treated with
 256 NaF alone and the normal control group fed with water and feed only.

257 **Table 3.4: Triiodothyronine (T₃), Thyroxine (T₄), and Thyroid Stimulating Hormone**
 258 **(TSH) concentration of control and test groups of NaF-induced toxicity and fecundity**
 259 **profile of adult male Wistar after 6 weeks of treatment**

Group	T ₃ (ng/ml)	T ₄ (Mlμ/l)	TSH (Mlμ/l)
Control	0.83±0.06	4.23±0.35	0.89±0.70
NaF	0.74±0.06	4.13±0.48	0.93±0.10
Stem bark extract			
10mg/kg NaF + 500	0.43±0.03 ^{αβ}	3.60±0.10	1.16±0.15
10mg/kg NaF + 1000	0.33±0.02 ^{αβ}	5.63±1.13	1.23±0.03
10mg/kg NaF +1500	0.36±0.05 ^{αβ}	4.53±1.70	1.03±0.16
10mg/kg NaF + 2000	0.30±0.08 ^{αβ}	6.17±1.04 ^{αβ}	1.67±0.29 ^β
10mg/kg NaF + 2500	0.37±0.15 ^{αβ}	4.70±0.26	2.33±0.49 ^{αβ}
Leaf extract			
10mg/kg NaF + 500	0.26±0.06 ^{αβ}	2.60±0.10	2.15±0.15 ^{αβ}
10mg/kg NaF + 1000	0.40±0.05 ^{αβ}	3.02±0.03	1.55±0.35 ^β
10mg/kg NaF +1500	1.21±0.27 ^{αβ}	3.15±0.05	1.67±0.02 ^β
10mg/kg NaF + 2000	2.40±0.57 ^{αβ}	3.45±0.05	2.05±0.05 ^{αβ}
10mg/kg NaF + 2500	1.65±0.22 ^{αβ}	6.16±5.67 ^{αβ}	1.86±0.06 ^{αβ}
Fruit Juice			
10mg/kg NaF + 500	0.44±0.06 ^{αβ}	5.50±0.61	0.87±0.08
10mg/kg NaF + 1000	0.25±0.05 ^{αβ}	5.05±0.95	0.84±0.01
10mg/kg NaF +1500	0.42±0.08 ^{αβ}	4.30±0.30	0.85±0.01
10mg/kg NaF + 2000	0.46±0.31 ^{αβ}	8.87±3.63 ^{αβ}	0.97±0.11
10mg/kg NaF + 2500	0.45±0.31 ^{αβ}	4.10±0.10	1.96±0.06 ^{αβ}

260 Results are expressed as Mean±SD; n=4

261 The mean values with β as superscripts across the column compared with group treated
 262 with NaF alone are considered significant ($p < 0.05$). The mean values with α as superscripts
 263 across the column compared with control group fed with water and feed only are considered
 264 significant ($p < 0.05$)

265 Sperm Count and Sperm Motility

266 Results obtained on the fecundity profile of adult male Wistar rats of NaF-induced toxicity
 267 (Table 3.5), showed that administration of 10 mg/kg NaF caused a concentration dependent
 268 and statistically significant ($p < 0.05$) reduction in sperm count in rats in comparison with

269 control group. However, all the groups treated with NaF + fruit juice, NaF + stem bark and
 270 NaF + leaf extracts showed significant increase ($p < 0.05$) in sperm count except the group
 271 treated with NaF + 500 mg/kg of leaf extract. Similarly, result for sperm motility showed that
 272 NaF at a dose of 10mg/kg caused significant decrease ($p < 0.05$) in sperm motility in
 273 comparison with the control groups. Fruit juice at higher doses of 1500 - 2500 mg/kg, NaF +
 274 2500 mg/kg of stem bark and NaF + 500 - 2500 mg/kg of leaf extracts caused a significant
 275 increase ($p < 0.05$) in sperm motility when compared with the NaF treated group.

276 **Table 3.5: Sperm count and Sperm motility of control and test groups of NaF-induced**
 277 **toxicity and fecundity profile of adult male Wistar after 6 weeks of treatment.**

Group	Sperm count (x 10 ⁶ /ml)	Sperm motility (%)
Control	960.03±5.00 ^β	94.00±4.00 ^β
NaF	207.04±2.00 ^α	30.00±1.00 ^α
Stem bark extract		
10mg/kg NaF + 500	240.00±2.00 ^{αβ}	31.00±1.00 ^α
10mg/kg NaF + 1000	464.00±2.65 ^{αβ}	32.00±3.00 ^α
10mg/kg NaF +1500	417.00±7.00 ^{αβ}	34.00±4.00 ^α
10mg/kg NaF + 2000	592.00±2.00 ^{αβ}	35.33±3.79 ^α
10mg/kg NaF + 2500	570.00±8.00 ^{αβ}	38.00±4.90 ^{αβ}
Leaf extract		
10mg/kg NaF + 500	126.00±2.00 ^{αβ}	20.00±1.00 ^{αβ}
10mg/kg NaF + 1000	241.00±1.00 ^{αβ}	26.00±5.00 ^α
10mg/kg NaF +1500	211.34±1.34 ^{αβ}	25.00±2.00 ^α
10mg/kg NaF + 2000	502.00±2.00 ^{αβ}	30.00±2.00 ^α
10mg/kg NaF + 2500	569.00±4.00 ^{αβ}	31.00±4.00 ^α
Fruit Juice		
10mg/kg NaF + 500	341.03±1.04 ^{αβ}	31.00±2.00 ^α
10mg/kg NaF + 1000	634.21±4.21 ^{αβ}	37.33±6.51 ^α
10mg/kg NaF +1500	450.00±5.00 ^{αβ}	40.00±4.00 ^{αβ}
10mg/kg NaF + 2000	694.51±4.51 ^{αβ}	57.00±3.00 ^{αβ}
10mg/kg NaF + 2500	651.52±1.51 ^{αβ}	41.00±2.00 ^{αβ}

278 Results are expressed as Mean±SD; n=4

279 The mean values with β as superscripts across the column compared with group treated
 280 with NaF alone are considered significant ($p < 0.05$). The mean values with α as superscripts
 281 across the column compared with control group fed with water and feed only are considered
 282 significant ($p < 0.05$)

283 Live/viable Sperm Cells and Percentage Dead Sperm Cells

284 Results of live/viable sperm cells of NaF-induced toxicity on fertility profile of adult male
 285 Wistar rats (Table 3.6) showed that administration of 10mg/kg of NaF caused a significant

286 decrease ($p < 0.05$) in the percentage of live and a significant increase ($p < 0.05$) in percentage
 287 of dead sperm cells in comparison with the control group. However, treatment with NaF +
 288 1500 - 2500mg/kg of fruit juice and NaF + 2000 - 2500 mg/kg stem bark extract caused a
 289 significant increase ($p < 0.05$) in the percentage of live sperm cells and a significant decrease
 290 ($p < 0.05$) in percentage of dead sperm cells when compared with the group treated with NaF
 291 alone. The leaf extract did not produce any visible ameliorative effect in the NaF-induced
 292 toxicity sperm cells.

293 **Table 3.6: Percentage Live/Viable sperm cells and Sperm dead cells of control and**
 294 **test groups of NaF-induced toxicity and fecundity profile of adult male Wistar** after 6
 295 **weeks of treatment**

Group	Live/Viable sperm cells (%)	Sperm dead cells (%)
Control	97.08±2.08 ^β	3.01±0.01 ^β
NaF	32.08±2.02 ^α	67.62±7.62 ^α
Stem bark extract		
10mg/kg NaF + 500	32.65±2.33 ^α	67.25±2.25 ^α
10mg/kg NaF + 1000	34.73±4.00 ^α	65.17±5.17 ^α
10mg/kg NaF + 1500	38.90±1.90 ^α	61.09±1.09 ^α
10mg/kg NaF + 2000	41.66±1.33 ^{αβ}	58.34±2.34 ^α
10mg/kg NaF + 2500	40.39±2.30 ^{αβ}	59.51±5.50 ^α
Leaf extract		
10mg/kg NaF + 500	28.68±3.30 ^α	71.32±1.32 ^α
10mg/kg NaF + 1000	32.52±2.02 ^α	67.28±3.21 ^α
10mg/kg NaF + 1500	32.04±3.04 ^α	67.86±2.11 ^α
10mg/kg NaF + 2000	33.15±3.10 ^α	66.85±6.85 ^α
10mg/kg NaF + 2500	36.07±4.07 ^α	64.93±4.00 ^α
Fruit Juice		
10mg/kg NaF + 500	34.78±4.00 ^α	65.12±3.12 ^α
10mg/kg NaF + 1000	40.43±1.23 ^{αβ}	59.24±5.24 ^α
10mg/kg NaF + 1500	43.47±3.40 ^{αβ}	57.54±2.32 ^{αβ}
10mg/kg NaF + 2000	60.92±5.02 ^{αβ}	49.07±7.07 ^{αβ}
10mg/kg NaF + 2500	47.81±7.01 ^{αβ}	53.18±3.18 ^{αβ}

296 Results are expressed as Mean±SD; n=4

297 The mean values with **β** as superscripts across the column compared with group treated
 298 with NaF alone are considered significant ($p < 0.05$). The mean values with **α** as superscripts
 299 across the column compared with control group fed with water and feed only are considered
 300 significant ($p < 0.05$)


301

302 Epididymal Sperm pH

303 The results obtained showed that the administration of 10mg/kg of NaF alone and
304 concomitant administration of NaF + fruit juice, NaF + stem bark and NaF + leaf extracts had
305 no significant effect on epididymal sperm pH. Epididymal sperm pH result after 6 weeks of
306 treatment was 6 for all the tested groups.

307 Histology Results


308 Photomicrographs of thin sections (5 μ m) of the testes of experimental rats harvested at the
309 end of 6 weeks of treatment with fruit juice of *A. muricata* (Plate 1) and stained with H&E
310 (400x).



311 Plate 1 shows photomicrographs of a section of the testes in the control groups and rats
312 treated with NaF and different concentrations of fruit juice of *A. muricata*.
313 A – Group 1 rats, that received feed and water only, showed normal testicular micro
314 architecture. There was normal spermatogenesis with different stages of differentiation and
315 maturation. Seminiferous tubules were lined with interstitial cells of the Leydig and well
316 enhanced spermatogenesis. B – Group 2 rats treated with 10 mg/kg NaF showed severe
317 testicular damage with severe spermatogenic arrest and severe apoptosis of the interstitial
318 cell of Leydig. The overall features are ghost like. There was lack of differentiation and
319 maturation of spermatogenesis and there was marked infiltration in the interstitial area of
320 seminiferous tubules. Severe spermatogenic arrest and severe apoptosis of the interstitial
321 cell of Leydig. C – Group 3 rats treated with 10 mg/kg NaF and 500 mg/kg of Fruit juice
322 showed mild restoration with mild enhanced spermatogenesis. However there are moderate
323 cellular apoptosis in some areas. D – Group 4 rats treated concomitantly with 10 mg/kg NaF
324 and treated with 1000 mg/kg of Fruit juice showed moderate restoration with moderate

325 enhanced spermatogenesis and moderate restoration of the interstitial cells of the Leydig. E
 326 – Group 5 rats treated concomitantly with 10 mg/kg NaF and 1500 mg/kg of Fruit juice
 327 showed moderate restoration with well enhanced spermatogenesis and interstitial cell of
 328 the Leydig appears normal. F – Group 6 rats treated concomitantly with 10 mg/kg NaF and
 329 2000 mg/kg of Fruit juice showed moderate restoration with moderate enhanced
 330 spermatogenesis and interstitial cells of the Leydig that appears normal. G - Group 7 rats
 331 treated concomitantly with 10 mg/kg NaF and 2500 mg/kg of Fruit juice showed mild cellular
 332 apoptosis otherwise normal with well enhanced spermatogenesis.

333 Photomicrographs of thin sections (5 µm) of the Testes of experimental rats harvested at the
 334 end of 6 Weeks of treatment with leaf extract of *A. muricata* (Plate 2) and stained with H&E
 335 (400x).




336 Plate 2. Photomicrographs of a section of the testes of rats treated with leaf extract of *A.*
 337 *muricata*

338 H Group 8 rats treated concomitantly with 10 mg/kg NaF and 500 mg/kg of leaf extract
 339 showed moderate regeneration with moderate enhanced spermatogenesis. However there
 340 are moderate apoptosis of the interstitial cells of the Leydig. I – Group 9 rats treated
 341 concomitantly with 10 mg/kg NaF and 1000 mg/kg of leaf extract showed moderate
 342 regeneration with moderate enhanced spermatogenesis. However there are moderate
 343 spermatogenic arrest. J – Group 10 rats treated concomitantly with 10 mg/kg NaF and 1500
 344 mg/kg of leaf extract showed mild regeneration with moderate arrest of spermatogenesis. K
 345 – Group 11 rats treated concomitantly with 10 mg/kg NaF and 2000 mg/kg of leaf extract
 346 showed mild regeneration with moderate arrest of spermatogenesis and severe apoptosis
 347 of the interstitial cell Leydig. L – Group 12 rats treated concomitantly with 10 mg/kg NaF and
 348 2500 mg/kg of Leave extract showed mild regeneration with severe apoptosis of the
 349 interstitial cell ledig.

350 Photomicrographs of thin sections (5 µm) of the Testes of experimental rats harvested at the
 351 end of 6 Weeks of treatment with Stem bark extract of *A. muricata* (Plate 3) and stained with
 352 H&E (400x).

353
 354



356

357 Plate 3. Photomicrographs of a section of the testes of rats treated with stem bark extract of
 358 *A. muricata*

359 M – Group 13 rats treated concomitantly with 10 mg/kg NaF and 500 mg/kg of stem bark
 360 extract showed mild regeneration with moderate spermatogenic arrest and mild apoptosis of
 361 the interstitial cells of the Leydig. There was mild regeneration with moderate spermatogenic
 362 arrest and mild apoptosis of the interstitial cells of the Leydig. N – Group 14 rats treated
 363 concomitantly with 10 mg/kg NaF and 1000 mg/kg of stem bark extract showed mild
 364 regeneration with moderate spermatogenic arrest and distortion of seminiferous tubules.
 365 There was mild regeneration with moderate spermatogenic arrest and distortion of
 366 seminiferous tubules. O - Group 15 rats treated concomitantly with 10 mg/kg NaF and 1500
 367 mg/kg of stem bark extract showed moderate enhanced spermatogenesis and mild
 368 spermatogenic arrest. There was moderate spermatogenic arrest, moderate enhanced
 369 spermatogenesis and seminiferous tubules lined by sertoli cells. P – Group 16 rats treated
 370 concomitantly with 10 mg/kg NaF and treated with 2000 mg/kg of stem bark extract showed
 371 well regeneration with normal spermatogenesis and seminiferous tubules lined by sertoli
 372 cells. Q – Group 17rats treated concomitantly with 10 mg/kg NaF and 2500 mg/kg of stem
 373 bark extract showed well regeneration with mild apoptosis of sertoli cell.

374 Discussion

375 The recent findings that fluoride exposure leads to biochemical/histological alterations in
 376 male reproductive system through multiple pathways indicates that both
 377 assessment/prophylaxis of chronic fluoride exposures in human populations is urgently
 378 required. Observation from this research work also indicates that sodium fluoride at a dose
 379 of 10 mg/kg caused non-significant decrease in haemoglobin concentration, platelet count,
 380 packed cell volume, and non-significant increase in neutrophil count, total white blood cell,
 381 and lymphocytes count. However, combined administration of NaF + the fruit juice and
 382 ethanol extracts of stem bark and leaf produced non-significant increase in the haemoglobin,
 383 packed cell volume and lymphocytes. The fruit juice at the concentrations of 1000 - 2500

384 mg/kg, and the groups treated with 500 and 1000 mg/kg of leaf extract, and 1000 and 2500
385 mg/kg of stem bark extract exhibited significant increase in platelet count. Reduction in
386 haemoglobin and packed cell volume is an indication of either the destruction of red blood
387 cells or the decreased production, which may lead to anaemia. On the contrary an increase
388 in the count of red blood cell, haemoglobin and packed cell volume is suggestive of
389 polycythaemia and positive erythropoiesis [22, 23]. Hence a non-significant increase or
390 activation on haemoglobin and packed cell volume in fruit juice, stem bark and leaf extracts
391 treated animals in comparison with the normal control is indicative of the ameliorative
392 potential of these extracts against NaF induced toxicity. Therefore, an increased count of
393 white blood cells and lymphocytes in NaF treated group, as observed in the present study,
394 suggests that NaF might have compromised the immune system. This report is in agreement
395 with [24], who reported a non-significant decrease in haemoglobin concentration of rats
396 treated with NaF alone in comparison with the control group.

397 [25, 26] reported that reduced blood platelets affect the viscosity of blood, which is correlated
398 positively to blood pressure. Concomitant administration of NaF and *A. muricata* extracts for
399 30 days adversely affected the count of blood platelets which may produce a positive effect
400 on the viscosity of blood. Probably prolonged duration of the treatment may ameliorate the
401 toxic effect of NaF [27]. Reduction in platelet count in experimental animals has been
402 reported to indicate an adverse effect on the oxygen carrying capacity of the blood as well as
403 thrombopoietin. Both significant and non-significant increase in platelets counts observed
404 from the results of this study suggests that the administration of *A. muricata* fruit juice, leaf
405 and stem bark extracts may ameliorate the disruption in the oxygen-carrying capacity of the
406 blood caused by NaF.

407 The most important biochemical mechanism by which fluoride decreases the level of
408 testosterone is its interference with steroidogenesis in Leydig cells. According to earlier
409 research, this interference has been demonstrated, in which activity levels of testicular
410 steroidogenic marker enzymes 3β -hydroxysteroid dehydrogenase (3β -HSD) and 17β -
411 hydroxysteroid dehydrogenase (17β -HSD) decreased significantly in NaF-treated rats [5,
412 28]. Since testicular steroidogenesis is controlled by these two rate-limiting enzymes, a
413 decline in their activities in Leydig cells significantly decreases the production and therefore
414 the level of testosterone. Known mechanisms by which fluoride decrease testosterone levels
415 are; inducing changes in both structures and enzyme activities in Leydig cells and interfering
416 with hypothalamus-hypophysis-testis axis [29]. Leydig cells require normal expression and
417 function of epidermal growth factor receptor (EGFR), androgen receptor (AR) and G-proteins
418 in order to synthesize testosterone. Fluoride exposure has been shown to reduce both
419 EGFR and AR expression [30] and to interfere with G-proteins in Leydig cells. However,
420 fluoride has been found to interfere with hypothalamus-hypophysis-testis axis [31]. The non-
421 significant decrease in testosterone level in NaF treated group in relation to the control group
422 reported in this study is consistent with so many previous research works which had
423 demonstrated that the NaF toxicity leads to a decrease in testosterone, a key hormone in
424 spermatogenesis [2, 32]. The result further reveals that concomitant administration of 10
425 mg/kg of NaF and extracts on testosterone levels of all the stem bark and leaf extracts
426 treated groups exhibited concentration-dependent significant increases while groups treated
427 with NaF and fruit juices exhibited no obvious changes. This observed increase could be
428 attributed to the interference of their phytochemical constituent(s) on the inhibitory action of
429 fluoride ion on steroidogenesis in Leydig cells or their antioxidant effect (properties) on free
430 radical generation by fluoride.

431 A lower concentration of 500 mg/kg of stem bark extract and 1000 mg/kg of fruit juice
432 produced a significant increase in FSH and LH concentrations respectively. This suggests
433 that stem bark extract and fruit juice at lower doses, with its antioxidant properties
434 ameliorated the toxicity effects of NaF on Gonadotropin hormones. Gonadotropins are
435 luteinizing hormone and follicle stimulating hormone from the pituitary gland. Testosterone in
436 males secreted by Leydig interstitial cells is increased under the influence of luteinizing
437 hormone. FSH regulates the development, growth, pubertal maturation and reproductive
438 processes of the body. Diminished secretion of FSH can result in hypogonadism. This
439 condition is typically manifested in males as a failure in the production of normal numbers of
440 sperm. Serum levels of FSH are decreased in anterior pituitary hypofunction, hypothalamic
441 disorders. Serum levels of LH are decreased in pituitary hypothalamic impairment.
442 Gonadotropin-releasing hormone stimulates the production and release of follicle stimulate
443 hormone (FSH) and luteinizing hormone (LH) from the pituitary gland [33].

444 Studies have reported that fluoride affects the synthesis of thyroid hormones, which
445 inversely impair the normal function of the male fecundity. Fluoride has been shown to
446 increase thyroid stimulating hormone (TSH) and reduce triiodothyronine (T_3) and thyroxine
447 (T_4) [34]. Fluoride is considered to interfere with thyroid hormone levels mainly through three
448 mechanisms; impairing normal structures of the thyroid gland, disrupting iodine metabolism
449 in thyroid glands and interfering with the tissue-specific metabolism of thyroid hormones.
450 Clinch in her review pointed out that fluoride interferes with the activity of Na/K-ATPase and
451 the sodium-iodide symporter. Since iodide uptake is facilitated by the combined actions of
452 the Na/k-ATPase and the sodium/iodide symporter [35], a decrease in the activities of these
453 enzymes caused by fluoride would reduce the uptake of iodide in the thyroid gland and
454 subsequent production of thyroid hormones. High fluoride intake has also been shown to
455 inhibit the activity of thyroid peroxidase [36]. Since thyroid peroxidase is an enzyme which is
456 essential for the production of thyroid hormones, decreased activity of thyroid peroxidase
457 caused by fluoride would also lead to reduced thyroid hormone synthesis (hypothyroidism)
458 [35] and is associated with impotence and decreased libido. Thyroid hormone affect brain
459 chemistry involved in sexual arousal, which in turn necessary stimulates the autonomic
460 nervous system and affects many other hormones **necessary for metabolism** [37]. There is a
461 correlation between hypothyroidism and low serum testosterone concentration. Also, type 2
462 iodothyronine deiodinase which regulates the tissue-specific conversion of T_4 to the
463 genomically active T_3 is predominantly expressed in elongated spermatids, suggesting that
464 thyroid hormone might have a direct effect on spermatogenesis [38, 39]. It is an established
465 fact that T_3 regulates the maturation and growth of testis, controlling Sertoli cell and Leydig
466 cell proliferation and differentiation during testicular development in rats and other mammal
467 species [40]. Our observations on the effect of NaF on thyroid hormone agree with the
468 previous research that indicated that fluoride increases TSH but reduces T_3 and T_4 [41].
469 **However, higher concentrations of the leaf extract significantly increased ($p<0.05$) the**
470 **concentration of T_3 hormones in the animals.** Fluoride is considered to interfere with thyroid
471 hormone levels mainly through three mechanisms; impairing normal structures of the thyroid
472 gland, disruptive iodine metabolism in thyroid glands and interfering with the tissue-specific
473 metabolism of thyroid hormones [42]. Several studies reveal that fluoride can directly
474 damage the structures of thyroid follicles, resulting in the following abnormalities; flattened
475 follicle epithelial cells, reduced cytoplasm [43]. These structural disruptions by fluoride will
476 disrupt the synthesis of thyroid hormones in the thyroid follicles [44].

477 Once fluoride crosses blood-testis membrane barriers that protect spermatogenesis, after a
478 prolonged exposure, it causes lack of maturation and differentiation of spermatocytes,
479 fragmentation of spermatozoa in the epididymis, and even cessation of spermatogenesis
480 [45]. The present investigation was carried out to explore the effects of fluoride (10mg/kg
481 NaF) and the possible ameliorative role of concomitant administration of fruit juice, leaf and
482 stem bark ethanol extract on the seminal characteristic of adult male Wistar rats. The sodium
483 fluoride treatment caused a substantial significant decrease in epididymal sperm motility,
484 progressive sperm motility, sperm concentration and live spermatozoa (%) along with a
485 simultaneous increase in dead spermatozoa (%) as compared to the rats of the control
486 group. Findings from this research work agree with [46, 47] who reported that exposure to
487 high concentrations of NaF leads to decreased sperm count, sperm motility, sperm survival
488 and increase in sperm abnormalities. The most important consequence of these fluoride
489 exposures is changes in the structure and functional behaviour of spermatozoa, disruption of
490 spermatogenesis and disturbance of multiple hormone systems that impact male fecundity.

491 **Conclusion**

492 Haematological evaluation indicates that doses above 500 mg/kg body weight of fruit juice,
493 stem bark and leaf extracts of *A. muricata* produced significant increases ($p<0.05$) in Hb
494 concentration and PCV while 2000 mg/kg of fruit juice and 1000 mg/kg of stem bark, and
495 leaf extracts produced a significant increase ($p<0.05$) in platelet count in comparison with
496 both the group treated with NaF alone and control group. The histopathologic findings in the
497 present study corroborates with the report from cauda epididymal spermatozoa analysis. It
498 might be concluded that NaF at 10 mg/kg caused potential reproductive cytotoxicities
499 leading to significant alterations in testicular tissue, altered semen characteristics, various
500 morphological abnormalities in spermatozoa and haematological parameters. Concomitant
501 administration of the fruit juice, ethanol stem bark and leaf extracts of *A. muricata* for a
502 period of 6 weeks resulted in significant prophylactic amelioration in all parameters altered.
503 Therefore, fruit juice, ethanol extracts of stem bark and leaf of *A. muricata* therapy could be
504 beneficial for the amelioration of fluoride-induced toxicity in male reproductive system and
505 fertility in genera at the tested dosages.

506 **Ethical Approval:**

507

508 As per international standard or university standard ethical approval has been collected and
509 preserved by the authors.

510

511 **References**

512 [1] Elbeticha A, Darmiani H, Al-Hiyasat AS. Fertility effects of sodium fluoride in male mice.
513 *Fluoride*. 2000;33(3): 128-134.

514 [2] Ortiz-Perez D, Rodriguez-Martinez F, Borja-Aburto VH, Castelo J, Grimaaldo JI, de La
515 Cruz E, Carrizaes L, Diaz-Barriga F. Fluoride-induced disruption of reproductive hormones in
516 men. *Environmental Research*. 2003;93(1): 20-30.

517 [3] Long H, Jin Y, Lin M, Sun Y, Zhang L, Clinch C. Fluoride toxicity in the male reproductive
518 system. *Fluoride*. 2009; 42(4): 260–276.

- 519 [4] Barbier O, Areola-Mendoza L, Dei Raco LM. Molecular mechanisms of fluoride toxicity.
520 Chemico-Biological Interactions. 2010;188(2): 319-333.
- 521 [5] Pushpalatha T, Srinivas M, Sreenivasula RP. Exposure to high fluoride concentration in
522 drinking water will affect spermatogenesis and steroidogenesis in male albino rats.
523 Biometals. 2005;18(3): 207-212.
- 524 [6] Patel S, Patel JK. A review of a miracle fruit of *Annona muricata*. Journal of
525 Pharmacognosy and phytochemistry. 2016;5(1): 137-148.
- 526 [7] Sharimila MA, Sathya RI. Green synthesis of *A. muricata* (L) ZNO NPS UV absorber on
527 textile. International Journal of Advanced Research. 2017; 5(5): 1721-1732.
- 528 [8] Sawat R, Dongre TP. Compositional analysis of *A. muricata* fruits. International Journal of
529 Universal Pharmacy and Bio Sciences. 2014; 3: 82-104.
- 530 [9] Adeyemi DO, Komolafe OA, Adewole SO, Obuotoi EM. Effects of *Annona muricata* (Linn)
531 on the morphology of pancreatic islet cells of experimentally induced diabetic wistar rats.
532 International Journal of Alternative Medicine. 2007;5: 1-8.
- 533 [10] Castell A, Bayes I, Ard A. Bioactivity of flavonoids in cells. Comprehensive Reviews in
534 Food Science and Food Safety. 2007;7: 299-398.
- 535 [11] Sandhar P, Humar HK, Prasha B, Tiwari T, Salhan P, Sharma M. A review of
536 phytochemistry of flavonoids. International Journal of Pharmaceutical Sciences. 2011;1: 25-
537 41.
- 538 [12] Sunarni T, Leviana F, Fiderianny I, Iwo MI, Wirasutisna KR. The antiperuricemic rats'
539 model and enzyme assay. Asian Journal of Pharmaceutical and Clinical Research. 2015; 8:
540 200-253.
- 541 [13] Gbeassor M., Kedjagni A.Y., Koumaglo K., De Souza C., Agbo K. , Aklikokou K.,
542 Amegbo K.A. (2006) In vitro antimalarial activity of six medicinal plants. *Phytotherapy*
543 *Research*, 4: 115-117.
- 544 [14] Gleye C, Duret P, Laurens A, Hocquenniller R, Laprovote O, Cava A. Journal of Natural
545 products.1996; 61: 576-579.
- 546 [15] Rodriguez-Fragoso L, Reyes-Esparza J, Burchiel S, Herreza-Ruiz D, Torres E. Risks
547 and benefits of commonly used herbal medicine in Mexico: Toxicology of Applied
548 Pharmacology. 2008;227(1): 125-135.
- 549 [16] Kedari TS, Khan AA. Gubayano (*Annona muricata*): A review of its traditional uses,
550 phytochemistry and pharmacology. American Journal of Research Communications.
551 2014;2(10): 247-258.
- 552 [17] Abdullah M, Syam AF, Maulahela H, Prabu OG, Laksono B, Meilany S, et al. The value
553 of caspase-3 after the application of soursop leaves extract (*Annona muricata*) in COLO205
554 colorectal cancer cell line. *J Gastroenterol Hepatol* 2016;21:199
555 <https://doi.org/10.1155/2017/4357165>
- 556 [18] Kiernan JA. Histological and histochemical methods: Theory and practice. 4th Edn.
557 Scion, Bloxham, UK; 2008.
- 558 [19] Dacie JV, Lewis S. M. Investigation of haematological disorders. Practical Haematology,
559 Churchill Livingstone Edinburgh, United Kingdom, 2006;177-180.
- 560 [20] Bjorndahl L, Soderlund I, Kvist U. Evaluation of the one-step eosin-nigrosin staining
561 technique for human sperm vitality assessment. Human Reproduction. 2003;8(4): 813-816.

- 562 [21] Ekaluo UB, Udokpoh AE, Ikpeme EV, Peter EU. Effect of Chloroquine treatment on
563 sperm count and weight of testes in male rats. *Global Journal of pure and Applied Science*.
564 2008;14: 175-177.
- 565 [22] Mansi K, Lahham J. Effects of *Artemisia sieberi* Besser (A/herba-alba) on heart rate and
566 some hematological values in normal and alloxan induced diabetes rats. *Journal of Basic*
567 *and Applied Sciences*. 2008;4: 57-62.
- 568 [23] Kuppast IJ, Vasudev NJ, Ravi MC, Birada SS. Studies on the haematological effects of
569 the extracts of *Cordiachotoma forst.* F. fruits. *Res. Journal of Pharmacol.*
570 *Pharmacodynamics*. 2009;1: 117-119.
- 571 [24] Zabulyte D, Uleckiene S, Kalibatas J, Paltanaviciene A, Jascaniniene N, Stosik M.
572 *Experimental studies on effect of sodium fluoride and nitrate on biochemical parameters in*
573 *rats. Bulletin of Veterinary Institute Pulawy*. 2007; 61: 79-82.
- 574 [25] Adedapo AA, Sofidiya MO, Masika PJ, Afolayan AJ. Safety evaluations of the aqueous
575 extract of *Acacia Karroo* stem bark in rats and mice. *Records of Natural Products*.
576 2008;2:128-134.
- 577 [26] Adeniyi TT, Ajayi GO, Akinsanya MA, Jaiyeola TM. Biochemical changes induced in rats
578 by aqueous and ethanoic corm extract of *Zyotritonia croceae*. *Scientific Research and*
579 *Essay*. 2010;5: 71-76.
- 580 [27] Mishra N, Tandon VL. Haematological effects of aqueous extract of ornamental plants in
581 male Swiss albino mice. *Vet World*. 2012;5(1): 19-23.
- 582 [28] Sarkar SD, Maiti R, Ghosh D. Management of fluoride induced testicular disorders by
583 calcium and vitamin-E co-administration in the albino rats. *Reproductive Toxicology*.
584 2006;22(4): 606-612.
- 585 [29] Hu L, Ying J, Mu L, Yu S, Liang Z, Carole C. Fluoride toxicity in the male reproductive
586 system. *Fluoride*. 2009;42(4): 17-32.
- 587 [30] Huang C, Yang H, Niu R, Sun Z, Wang J. Effect of sodium fluoride on androgen
588 receptor expression in male mice. *Fluoride*. 2008;41: 10-17.
- 589 [31] Ma X, Cheng X, Li F, Guo J. Experimental research on endocrine disturbing effect of
590 fluorine on hypothalamus-hypophysis-testis axis in male rats. *Wei Sheng Yen Jiu*.
591 2008;37(6): 733-775.
- 592 [32] Ghosh D, Das SS, Maiti, R, Jana D, Das UB. Testicular toxicity in sodium fluoride
593 treated rats: association with oxidative stress. *Reproductive Toxicology*. 2002;16(4): 385-
594 390.
- 595 [33] Schalli AV, Kastin AV, Arimura A. Hypothalamic follicle-stimulating hormone (FSH) and
596 luteinizing hormone (LH)-regulating hormone: structure, physiology, and clinical studies.
597 *Fertility and Sterility*. 1971;22(11): 703-721.
- 598 [34] Wang H, Yang Z, Zhou B, Gao H, Yan X, Wang J. Fluoride-induced thyroid dysfunction
599 in rats: roles of dietary protein and calcium level. *Toxicology and Industrial Health*.
600 2009a;25(1): 49-57.
- 601 [35] Clinch C. Fluoride interactions with iodine and iodide: implication for breast health.
602 *Fluoride*. 2009; 42(2): 75-87.
- 603 [36] Zhan XA, Li JX, Wang M, Xu ZR. Effects of fluoride on growth and thyroid function in
604 young pigs. *Fluoride*. 2006;39(2): 95-100.
- 605 [37] Arem R. *The Thyroid Solution*. New York. Ballantine Books. 2000;137-151

- 606 [38] Donnelly P, White C. Testicular dysfunction in men with primary hypothyroidism; Reversal
607 of hypogonadotropic hypogonadism with replacement thyroxine. *Clinical Endocrinology*.
608 2000;52: 197-201.
- 609 [39] Wajner SM, dos Santos Wagner M, Melo RC, Parreira GG, Chiarini-Garcia H, Bianco
610 AC, Fekete C, Sanchez E, Lechan RM, Maia AL. Type 2 iodothyronone deiodinase is highly
611 expressed in germ cells of adult rat testis. *Journal of Endocrinology*. 2007;194(1): 47-54.
- 612 [40] Holsberger DR, Cooke PS. Understanding the role of thyroid hormone in Sertoli cell
613 development: a mechanistic hypothesis. *Cell and Tissue Research*. 2005; 322: 133-140.
- 614 [41] Bachinskii PP, Gutsalenko OA, Naryzhniuk ND, Sidora VD, Shliakhta AL. Action of the
615 body fluorine of healthy persons and thyroidopathy patients on the function of hypophyseal-
616 thyroid the system. *Prob Endokrinol (Mosk)*. 1985;31(6): 25-29.
- 617 [42] Hu L, Ying J, Mu L, Yu S, Liang Z, Carole C. Fluoride toxicity in the male reproductive
618 system. *Fluoride*. 2009;42(4): 17-32.
- 619 [43] Wang J, Ge Y, Ning H, Nui R. DNA damage in brain and thyroid gland cells due to high
620 fluoride and low iodine In: Preedy, V.R., Burrow, G.N., Watso, R., editors. *Comprehensive
621 handbook of iodine: nutritional, biochemical pathological and therapeutic aspects*. San
622 Diego: Academic Press. 2009b;pp. 643-649.
- 623 [44] Bouaziz H, Soussia L, Guermzi F, Zeghal N. Fluoride-induced thyroid proliferative
624 changes and their reversal in female mice and their pups. *Fluoride*. 2005;38(2): 185-192.
- 625 [45] Nair N, Bedwal S, Prasad S, Saini MR, Bedwal RS. Short-term zinc deficiency in diet
626 induces increased oxidative stress in testes and epididymis of rats. *Indian Journal of
627 Experimental Biology*. 2005;43(9): 786-794.
- 628 [46] Huang C, Yang HB, Niu RY, Sun ZL, Wang RS, Chen YT. Infertility with defective
629 spermatogenesis and steroidogenesis in male mice lacking androgen receptor in leydig
630 cells. *Endocrine*. 2007;32(1): 96-106.
- 631 [47] Giri DK, Ghosh RC, Kashyap DK, Govina D. Evaluation of subacute fluoride toxicity on
632 spermatozoa and testicular tissue of male wistar rats. *Bioscan*. 2013;8(3): 983-987.