

# An Overview of Development of Quantitative Neurotoxicity Testing *In Vitro*

## ABSTRACT

*In vitro* neurotoxicity testing has been hampered by the fact that the brain architecture is complex. However, a series of innovation of neuroculturing broke through the barrier. The establishment of culturing for the primary neuron and the immobilized neuroblastoma cells enables neurotoxicity testing *in vitro*. Following to necrotic cell death, extensive morphological changes as seen during neuronal differentiation was used for the endpoints of neurotoxicity. Two-dimensional imaging techniques facilitated quantitative analyses of toxicity of many neurotoxins. Three-dimensional culturing of neurospheres *in vitro* has been expected to investigate the neurodevelopmental toxicity. The neurosphere assay *in vitro* also improved the sensitivity to estimate the neurotoxicity. The present study highlights an overview about the *in vitro* neurotoxicity testing for rotenone, a dopaminergic pesticide as an environmental toxicant.

**Keywords:** PC12 cells; NB-1 cells; neural stem cells; neurospheres; rotenone; quantitative neurotoxicity

## 1. INTRODUCTION

In light of the large number of chemicals, there is a demand to develop rapid screening techniques. Historically, PC12 cells have been used for testing neurotoxicity. The PC 12 cell line was derived from rat pheochromocytoma, a tumor arising from chromaffin cells of the adrenal medulla and developed to study cell differentiation [1]. Neuronal differentiation is a complex process that induces both morphological and biochemical changes. The most obvious things are a decrease in cell proliferation, and the emergence of extending processes. During neuronal differentiation, cells also acquire excitability and start to express some chemical coding genes that provide their functional identity. Upon exposure to neuronal stimulation, PC12 cells gradually exit the mitotic cycle and begin to differentiate, developing axonal projections, electrical excitability, and the characteristics of cholinergic and catecholaminergic neurons. Therefore, the PC12 model enables the detection of environmental toxicants.


Following to rat PC12 model cells, human cell lines such as NB-1 and SH-SY5Y were used to investigate potential species-specific differences, rather than non-human cell origins. It was expected to extrapolate human toxicity.

Recent evidence points to important contributions of exposure to environmental neurotoxins in the marked increase in neurodevelopmental disorders [2-4]. In response to the need for more efficient methods to identify potential developmental neurotoxins, neurosphere assay has recently been established. Neural stem cells play an essential role

in the development of central nervous system, having self-renewal potency and being multipotential. In 1992, it was demonstrated that cells from central nervous system of adult and embryonic mice can be isolated and propagated in culture [5]. In the presence of epidermal growth factor, cell agglomerations, termed neurospheres were formed. They proliferate in culture and have the ability to migrate and differentiate into neurons, astrocytes, and oligodendrocytes. Neurosphere culturing is three-dimensional cell systems and there a valuable *in vitro* model that mimics basic processes of brain development. Therefore, neurospheres are a useful tool for testing chemicals for their abilities to interfere with these processes: proliferation, migration, differentiation, and apoptosis.

In this paper, we compare a variety of cell-based neurotoxicity testing with several endpoints using rotenone, a dopaminergic pesticide (Fig.1, ref 6-8).

**Figure 1. The structure of rotenone and its derived plant**



## **2. MATERIAL AND METHODS**

### **2.1 Culture of PC12 cells**

PC12 cells (RCB 0009; RIKEN, Tsukuba, Japan) were grown in Dulbecco's modified Eagle's Medium (Sigma-Aldrich) supplemented with 10% fetal bovine serum (FBS; Life Technologies, Inc., Rockville, MD), 4.5 mg/ml glucose, penicillin (100 U/ml), and streptomycin (100 µg/ml) in a humidified atmosphere of 95% air, 5% CO<sub>2</sub> at 37 °C. The cells were subcultured (1:3) 2 to 3 times per week. Cell viability was determined by trypan blue exclusion method.

### **2.2 Culture of NB-1 cells**

Human neuroblastoma NB-1 cells were cultured in 45% RPMI 1640 and 45% Eagle's minimum essential medium containing 10% FBS (Life Technologies, Inc.), sodium pyruvate, penicillin (100 U/ml), and streptomycin (100 µg/ml) in a humidified atmosphere of 95% air, 5% CO<sub>2</sub> at 37 °C. The cells were subcultured (1:6) once a week. The viable cell number of the NB-1 cells was estimated by crystal violet staining. Fixed and dried cells in a plate were rehydrated with distilled water and photographed under a phase-contrast microscope (DMIRB, Leica Microsystems, Tokyo, Japan) equipped with digital camera. The digital images obtained were then analyzed using image analysis software by counting the cell number and total neurite length in the image field. The degree of neurite extension is represented as the total length of neuritis in micrometer per cell in randomly chosen phase-contrast microscope fields.

### **2.3 Culture of neurosphere**

Pregnant Wistar rats at embryonic day 14 (E14) were obtained from Clea (Tokyo, Japan). The animals were maintained in home cages at 22°C with a 12-h light-dark cycle. They received the MF diet (Oriental Yeast Corp., Tokyo, Japan) and distilled water ad libitum. All animal care procedures were in accordance with National Institute for Environmental Studies guidelines. The rats were sacrificed by diethyl ether overdose on E16. The embryos were removed and transferred to minimal essential medium (MEM; Sigma-Aldrich). Subsequently, the mesencephalons were dissected from the embryos, and were enzymatically digested with 50 U deoxyribonuclease I (Takara Corp., Kyoto, Japan) and 0.8 U papain (Sigma-Aldrich) at 32°C for 12 min. After stirring, the digestion mixture was passed through a 70-µm cell strainer (BD Biosciences). The run-through containing the neural stem cells was centrifuged at 800 x g for 10 min. It was then resuspended in Dulbecco's Modified Eagle's Medium (DMEM) and F12 medium (1:1; Invitrogen, Tokyo, Japan) supplemented with B27 (Invitrogen), 20 ng/ml basic fibroblast growth factor (bFGF; R&D Systems, Inc., MN) and 10 ng/ml epidermal growth factor (EGF; Roche Applied Science, Tokyo, Japan), and cultured in uncoated dishes without serum. Fresh culture medium containing EGF and bFGF was added after 3-4 days.

The neurospheres were seeded in an uncoated glass-bottomed dish (D110300; MATSUNAMI, Tokyo, Japan) in the presence of bFGF and EGF for 3 h, allow cells to adhere. The migrating distance of the cells was statistically measured from the edge of the sphere, using National Institute of Health ImageJ 1.38x software (public domain software).


TUNEL staining was carried out, as described previously [26, 27]. The cells were fixed in 4% paraformaldehyde, washed twice with PBS, and permeabilized in 0.5% Triton X-100 for 5 min on ice. TUNEL labelling was done with fluorescein dUTP (Roche Applied Science, Mannheim, Germany) in the presence of terminal deoxynucleotidyl transferase for 1 h at 37°C. Following labelling, the cells were washed with PBS twice and then directly surveyed under a fluorescence microscope. Images were captured using Viewfinder Lite ver.1.0 camera software through DP-50 digital camera (Olympus, Tokyo, Japan). For quantification of TUNEL-labeled cells, every field containing TUNEL-positive signals was photographed at 100x optical magnification. Then, TUNEL-positive cells were counted.

### 3. RESULTS

#### 3.1. Cell viability of PC12 cells exposed to rotenone

Rotenone neurotoxicity in PC12 cells was examined by trypan blue exclusion method. PC12 cells were exposed to variety of concentration of rotenone for 3 days. Following fixing the treated cells, a number of cells were counted. Cell viability was decreased in a semilogarithmic-linear, dose-dependent manner. IC<sub>50</sub> was about 1 µM (Fig.2).

**Figure 2. Rotenone neurotoxicity in rat pheochromocytoma PC12 cells.**




119

### 120 3.1. 1. Inhibition of neurite outgrowth of NB-1 cells exposed to rotenone

121 Rotenone neurotoxicity in NB-1 cells was examined as an endpoint of neurite  
 122 outgrowth. NB-1 cells were seeded on a culture plate and rotenone (0~50 μM) was  
 123 added for 24h. The treated cells were fixed and the length of neurite outgrowth was  
 124 measured by imaging analyses. The length was decreased in a semilogarithmic-  
 125 linear, dose-dependent manner. IC<sub>50</sub> was about 1 μM (Fig.3).

126 **Figure 3. Neurite outgrowth in rotenone-exposed human neuroblastoma**  
 127 **NB1 cells.**



138


139 3.1.2. Neurosphere assay for neurodevelopmental toxicity of rotenone

140 We isolated neural stem cells from E15 rat embryos (Fig.4a), using pooled mesencephalons  
141 from 12 fetuses. After 2~3 weeks in culture, neurospheres appeared (Fig. 4c), suggesting  
142 self-renewal occurred. Neurospheres of about 200  $\mu\text{m}$  in diameter consisted of about  $10^3$   
143 cells.

144

145 **Figure 4.** (a) An E15 rat fetus showing the mesencephalon. (b) Primary neurospheres after  
146 7 days *in vitro* (c) Primary neurospheres after 2~3 days *in vitro*. Adapted from ref. [9].


147



148

149 **Figure 5. Identification of cultured mesencephalic neurosphere.** Neurospheres were  
150 immunostained with (a) anti-nestin antibody, (b) anti-MAPs antibody, or (c) anti-GFAP  
151 antibody . Adapted from ref. [9].

152



153

154

155

156

157

158


159

160 To identify neural stem cells, we stained the neurospheres with an anti-nestin antibody, as  
161 shown in Figure 5a. The nestin-positive cells were localized both at the edge and within the  
162 spheres. Since neural stem cells are multipotent for neural differentiation, we also  
163 immunostained the neurospheres for MAPs, which were located in cells at the edge of the  
164 spheres (Fig.5b). Since on E15, when we isolated the neuronal stem cells, rat embryos are  
165 undergoing gliogenesis, we stained the neurospheres with anti-GFAP antibody, which mainly



166 stained cells at the sides of the spheres (Fig.5c). Our results suggested that heterogeneous  
167 cell populations were present in neurospheres, at late embryonic stages.

168 During the culture, cells emerged from the plated neurospheres and migrated along the  
169 radial axis (Fig. 6 a and b). After 3 h, the plated cells were treated with various  
170 concentrations of rotenone (0-10  $\mu$ M) for 24 h (Figs.7 b-f). The migration distance of the  
171 cells was measured from the edge of the neurospheres using NIH ImageJ 1.38x public  
172 domain software. Rotenone prevented the cells from migrating from the neurospheres in a  
173 linear, dose-dependent manner (Fig. 7g). The half-maximal inhibitory concentration ( $IC_{50}$ )  
174 was 0.32  $\mu$ M.

175  
176 **Figure 6. Neural stem cells radically migrated from neurosphere.** (a) The early  
177 stage , or (b) the later stage of the culturing  
178 (a) (b)



189  
190  
191  
192  
193  
194  
195 **Figure 7. Rotenone inhibition of cell migration from neurospheres *in vitro* at**  
196 **various concentrations .** (a) No migrating cells during the initial 3h, (b) 0  $\mu$ M, (c) 0.1  $\mu$ M ,  
197 (d) 1  $\mu$ M, (e) 5  $\mu$ M, (f) 10  $\mu$ M. The migration distance was quantitatively measured with NIH  
198 ImageJ 1.38x software (g). Adapted from ref. [10].  
199




217 To compare the toxicity of 2 chemicals, neurosphere assay for bisphenol A, an endocrine  
 218 disruptor, was carried out under the condition where rotenone was did (Fig.8). The percent  
 219 inhibition of migration by bisphenol A and rotenone at 1  $\mu$ M was 35% and 70%, respectively.  
 220 Thus, the rank order of potency of chemicals was: bisphenol A< rotenone. The value of  
 221 other endpoints was summarized in Table 1.

222

223 **Figure 8. Comparison of neurotoxicity of 2 chemicals by neurosphere assay *in vitro*.**

224

Modified from ref. 11.



225

226

227

228 **Table 1. Multiple endpoints used in chemical neurosphere testing *In Vitro***

229

Modified from refs. 10 and 11.

230

Endpoint	Chemical	
	Rotenone	Bisphenol A
1. Migration ( $IC_{50}$ )	0.32 $\mu$ M	↓
2. Proliferation ( $IC_{50}$ )	1.9 $\mu$ M	5.0 $\mu$ M
3. Apoptosis	↑	ND
4. Log $K_{ow}$	4.10	3.32

231 ↑:induction; ↓:inhibition; ND: not detected; values of Log  $K_{ow}$  were reported in International  
 232 Chemical Safety Cards (ICSC, Japanese Ver.).

233

234

**Table 2. Sensitivity to rotenone neurotoxicity in various *In Vitro* testing**

Source	Half maximal effects	Period s	Endpoint	Reference
Rat PC12 cells	1 $\mu$ M <	24h	Cell viability	20
	1 $\mu$ M	72h	Cell viability	Unpublished data



Human NB-1 cells	1 $\mu$ M	24h	Neurite outgrowth	Unpublished data
Human neurosphere	4 $\mu$ M	24h	Cell viability	21
Rat E16 neurosphere	0.32 $\mu$ M	24h	Migration	10
	1.9 $\mu$ M	24h	Proliferation	10
	1.4 $\mu$ M	24h	Apoptosis	10

#### 4. DISCUSSION

Rotenone is a botanical pesticide [6-8]. Neurotoxic nature of rotenone has been used to produce adult Parkinson model animals due to nigrostriatal dopaminergic lesions [12-14]. Upon the working hypothesis of the developmental origins of health and disease (DOHaD; ref.15), it is important to examine the neurodevelopmental toxicity of rotenone in neurosphere assays, comparing other cell lines. The hypothesis suggests that the environmental origin of human sporadic Parkinson disease occur early in life. One possible explanation for this phenomenon is that early exposure to neurotoxic chemicals reduces the number of dopaminergic neurons in the substantia nigra to levels below those needed to sustain normal function during the course of the neuronal attrition associated with aging.

There are many endpoints to evaluate neurotoxicity [Table 2]. Ultimate endpoint for neurotoxicity is cell death. Therefore, neuronal cell viability was used for evaluating its toxicity. Catastrophic cell death by neurotoxins is observed as perturbation of energy producing systems, cellular membrane defects, and increased influx of calcium ions. This is concomitant with necrotic action of the toxin. For more sensitive detection of the toxicity, a new biochemical marker is needed.

About 20 years ago, apoptotic nature of environmental toxins were discovered in several types of cells, including renal and neuronal cell [16,17]. Apoptosis is a programmed form of cell death mediating precisely controlled deletions of 'unwanted' cells. This phenomenon is initiated not only by physiological stimuli but also by an extensive array of nonphysiological agents. The characteristics of apoptosis are DNA fragmentation and chromatin condensation, in which are endpoints of toxicants as earlier phase than necrotic phase of toxins.

Development of techniques for Image acquisition enabled to open a new way to evaluate neurotoxicity: neurite outgrowth was also used for endpoint of neurotoxicity [18,19]. The growth of axonal and dendritic processes during brain development is a critical determinant of neural connectivity, and disruption of this process could lead to neuronal dysfunction. Neurite outgrowth can be recapitulated *in vitro* using a variety of cell models. These models have been become valuable tools for investigating the mechanism for known developmental neurotoxicants.

The developing human brain can be more susceptible to injury caused by toxic agents than the brain of an adult. Probably all potential neurotoxic compounds would also cause damage to the developing brain and at much lower doses. Indeed, neurodevelopmental disorders in children such as attention deficit disorder or autism have been associated with the exposure to chemicals in the environment during early fetal development [20].



272 Particularly, it has been suggested that one of possibility of autism-spectrum disorders are  
273 initiated in the embryonic neural stem cells [21]. Neural stem cells play an essential role in  
274 the development of central nervous system, having self-renewal potency and being  
275 multipotential: they are able to differentiate to neurons, astrocytes and oligodendrocytes to  
276 form neuronal architecture. Indeed, in the culture of neural stem cells, they form free-  
277 floating three-dimensional structures. Therefore, application of neurosphere for  
278 neurodevelopmental toxicity testing is reasonable.

279 Quantitative analyses in this study revealed the linearity in function as migration inhibition  
280 versus chemical concentration. Owing to the linearity of the functional relationship between  
281 the migration inhibition and the concentration of test chemicals, this approach could be  
282 employed as a reliable quantitative assay system, excluding the issue of nonlinearity in low  
283 dose of an endocrine disruptor such as bisphenol A [22,23].  
284

## 285 **5. CONCLUSION**

286 This paper highlights an overview about neurotoxicity of rotenone, investigating with rat  
287 PC12 cell, human NB-1 cells and rat embryonic neural stem cells. Furthermore, quantitative  
288 analysis revealed a linear function between the cellular endpoints and the rotenone  
289 concentration. This could be employed as a simple and rapid screening for neurotoxicity of  
290 environmental chemicals. Particularly, neurosphere assay would be hoped to develop the  
291 risk assessment methods for chemicals based on infant physiology.  
292

### 293 **Ethical Disclaimer:**

294  
295 As per international standard or university standard written ethical permission has been collected and preserved by  
296 the authors.  
297  
298  
299  
300  
301

## 302 **REFERENCES**

- 303  
304 1. Greene, L.A. Tischler, A.S. Establishment of a noradrenergic clonal line of rat  
305 adrenal pheochromocytoma cells which respond to nerve growth factor. Proc Natl Acad Sci  
306 U.S.A.1976, 73, 2424-2428.
- 307 2. Jacobson, J.L. Jacobson, S.W. Intellectual impairment in children exposed to  
308 polychlorinated bisphenyls in utero. N. Engl. J. Med. 1996, 335, 783-789.
- 309 3. Patandin, S., Lanting, C.I., Mulder, P.G., Boersma, E.R., Sauer, P.J., Weisglas-  
310 Kuperus, N. J. Pediatr. 1999, 134, 33-41.
- 311 4. Rice, D.C. Parallels between attention deficit hyperactivity disorder and behavioral  
312 deficits produced by neurotoxic exposure in monkeys. Environ. Health Perspect. 2000,  
313 108(Suppl.3), 511-533.
- 314 5. Reynolds, B.A., Tetzlaff, W., Weiss, S. A multipotent EGF-responsive striatal  
315 embryonic progenitor cell produces neurons and astrocytes. J. Neurosci. 1992, 12, 4565-  
316 4574.

- 317 6. Ray, D.E. Pesticides derived from plants and living organisms. In Handbook of  
318 Pesticide Toxicology, Hayes, W.J. and Laws E.R. Eds.; Academic Press, San Diego, US,  
319 1991; pp585-636.
- 320 7. Ujvary, I. Pest control agents from natural products. In Handbook of Pesticide  
321 Toxicology, 2nd ed.; Krieger, R. Ed.; Academic Press, California, US, 2001; pp109-180.
- 322 8. Ott, K.C. Rotenone. A brief review of its chemistry, environmental fate, and the  
323 toxicity of rotenone formulations. New Mexico Council of Trout Unlimited. 2008; Available at:  
324 <http://www.newmexicotu.org/Rotenone%20summary.pdf>.
- 325 9. Ishido M. Effects of *p*-nitrotoluene on cultured mesencephalic neural stem cells. J.  
326 Health Sci (Tokyo) 2009; 55, 114-118.
- 327 10. Ishido, M., Suzuki J. Inhibition by rotenone of mesencephalic neural stem-cell  
328 migration in a neurosphere assay in vitro. Toxicol in Vitro 2010; 24, 552-557.
- 329 11. Ishido, M., Suzuki, J. Quantitative analyses of inhibitory effects of bisphenol A on  
330 neural stem-cell migration using a neurosphere assay in vitro. J. Health Sci.(Tokyo) 2010;  
331 56, 175-181.
- 332 12. Betarbet, R., Sherer, T.B., MacKenzie, G., Garcia-Osuna, M., Panov, A.V.,  
333 Greenamyre, J.T. Chronic systemic pesticide exposure reproduces features of Parkinson's  
334 disease. Nat. Neurosci. 2000, 3, 1301-1306.
- 335 13. Dauer, W., Przedborski, S. Parkinson's disease: mechanisms and models. Neuron  
336 2003, 39; 889-909.
- 337 14. Blesa, J., Phani, S., Jackson-Lewis, V., Przedborski, S. Classic and new animal  
338 models of Parkinson's disease. J. Biomed. Biotech. 2012, doi:10.1155/2012/845618
- 339 15. Gilman, M.W., Barker, D., Bier, D., Cagampang, F., Challs, J., Fall, C., Godfrey, K.,  
340 Gluckman, P., Hanson, M., Kuh, D., Nathanielsz, P., Nestel, P., Thornburg, K.L. Meeting  
341 report on the 3rd international congress on developmental origins of health and disease  
342 (DOHaD). Pediatr. Res. 2007; 61, 625-629.
- 343 16. Ishido, M. et al. Cadmium-induced DNA fragmentation is inhibitable by zinc in  
344 porcine kidney LLC-PK<sub>1</sub> cells. Life Sci. 1995, 17, 351-356.
- 345 17. Kunimoto, M. Methylmercury induces apoptosis of rat cerebellar neurons in primary  
346 culture. Biochem. Biophys. Res. Commun. 1994, 204, 310-317.
- 347 18. Pramanik, R., Ishido, M., and Kunimoto, M. Methylmercury-mediated down-  
348 regulation of mtHSP70 and phospholipase A2 mRNA expression in human neuroblastoma  
349 NB-1 cells identified by cDNA macroarray analysis. J. Health Sci.(Tokyo) 2002, 48, 381-  
350 384.
- 351 19. Pramanik, R., Ishido, M., and Kunimoto, M. Effects of cadmium chloride on neurite  
352 outgrowth and gene expression in human neuroblastoma NB-1 cells. J. Health Sci.(Tokyo)  
353 2001, 47; 478-482.

- 354 20. Sai, Y., Wu, Q., Le, W., Ye, F., Dong, Z. Rotenone-induced PC12 cell toxicity is  
355 caused by oxidative stress resulting from altered dopamine metabolism. *Toxicology in Vitro*  
356 2008, 22; 1461-1468.
- 357 21. Li, J., Spletter, M.L., Johnson, D.A., Wright, L.S., Svendsen, C.N., Johnson, J.A.  
358 Rotenone-induced caspase 9/3-independent and -dependent cell death in undifferentiated  
359 and differentiated human neural stem cells. *J. Neurochem.* 2005; 92, 462-476.
- 360 22. Colborn, T. Neurodevelopment and endocrine disruption. *Environ Health Perspect.*  
361 2004, 115, 924-931.
- 362 23. Li, H., Radford, J.C., Ragusa, M.J., Shea, K.L., McKercher, S.R., Zaremba, J.D.,  
363 Soussou, W., Nie, Z., Kang, Y. J., Nakanishi, N., Okamoto, S., Roberts, A. J., Schwarz, J. J.,  
364 Lipton, S. A. Transcription factor MEF2C influences neural stem/progenitor cell  
365 differentiation and maturation in vivo. *Proc. Natl. Acad. Sci. U.S.A* 2008, 105, 9397-9402.
- 366 24. Welshons, W.V., Thayer, K.A., Judt, B.M., Taylor, J.A., Curran, E.M., vom Saal, F.S.  
367 Large effects from small exposure. *Environ. Health Perspect.* 2003, 111; 994-1006.
- 368 25. vom Saal, F.S., Hughes, C. An extensive new literature concerning low-dose effects  
369 of bisphenol A shows the needs for a new risk assessment. *Environ. Health Perspect.* 2005,  
370 113; 926-933.
- 371 26. Ishido, M. Melatonin inhibits maneb-induced aggregation of  $\alpha$ -synuclein in rat  
372 pheochromocytoma cells. *J. Pineal Res.* 2007, 42:125-130.
- 373 27. Ishido, M. Activation of STAT3 by pituitary adenylate cyclase-Activating polypeptide  
374 (PACAP) during PACAP-promoted neurite outgrowth of PC12 cells. *J. Mol. Neurosci.* 2010,  
375 42; 349-358.

376  
377  
378  
379  
380  
381  
382

## DEFINITIONS, ACRONYMS, ABBREVIATIONS

Here is the Definitions section. This is an optional section.

**Term:** Definition for the term