

Asymptotical behaviour of second-order rational difference equations system

Abstract

This paper deals with dynamics of the solution to second-order rational difference equations system

$$x_{n+1} = \frac{x_n}{A + y_n y_{n-1}}, \quad y_{n+1} = \frac{y_n}{B + x_n x_{n-1}}, \quad n = 0, 1, \dots,$$

where $A \in (0, \infty)$, $x_{-i} \in (0, \infty)$, $y_{-i} \in (0, \infty)$, $i = 0, 1$. Moreover the rate of convergence of a solution that converges to the equilibrium of system is discussed. Finally, some numerical examples are considered to show the results obtained.

MSC: 39A10

Keywords: Difference equation, Stability, Rate of convergence

1. Introduction

Difference equations or discrete dynamical systems are diverse field which impacts almost every branch of pure and applied mathematics. One of the reasons is a necessity for some techniques which can be used in investigating equations arising in mathematical models describing real life situations in population biology, economic, probability theory, genetics psychology, etc. The theory of difference equations occupies a central position in applicable analysis. There is no doubt that the theory of difference equations will continue to play an important role in mathematics as a whole. Rational difference equations of order greater than one are of paramount importance in applications. Such equations also appear naturally as discrete analogues and as numerical solutions of differential and delay differential equations. It is very interesting to investigate the behavior of solutions to rational difference equations and to discuss the local asymptotic stability of their equilibrium points. Recently there has been published quite a lot of works concerning the behavior of positive solutions of systems of difference equations [1-10]. These results are not only valuable in their own right, but they can provide insight into their differential counterparts.

Kurbanli[3] studied a three-dimensional system of rational difference equations

$$x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1} - 1}, \quad y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1} - 1}, \quad z_{n+1} = \frac{z_{n-1}}{y_n z_{n-1} - 1},$$

where the initial conditions are arbitrary real numbers.

Cinar et al. [4] has obtained the positive solution of the difference equation system

$$x_{n+1} = \frac{m}{y_n}, \quad y_{n+1} = \frac{p y_n}{x_{n-1} y_{n-1}}.$$

Cinar [5] has obtained the positive solution of the difference equation system

$$x_{n+1} = \frac{1}{y_n}, \quad y_{n+1} = \frac{y_n}{x_{n-1} y_{n-1}}.$$

Clark and Kulenovic [6] investigated the system of rational difference equations

$$x_{n+1} = \frac{x_n}{a + cy_n}, \quad y_{n+1} = \frac{y_n}{b + dx_n}, \quad n = 0, 1, \dots,$$

where $a, b, c, d \in (0, \infty)$ and the initial conditions x_0 and y_0 are arbitrary nonnegative numbers.

Zhang, Yang and Liu [7] investigated the global behavior for a system of the following third order nonlinear difference equations.

$$x_{n+1} = \frac{x_{n-2}}{B + y_{n-2}y_{n-1}y_n}, \quad y_{n+1} = \frac{y_{n-2}}{A + x_{n-2}x_{n-1}x_n},$$

where $A, B \in (0, \infty)$, and initial values $x_{-i}, y_{-i} \in (0, \infty), i = 0, 1, 2$.

Ibrahim [9] has obtained the positive solution of the difference equation system in the modeling competitive populations.

$$x_{n+1} = \frac{x_{n-1}}{x_{n-1}y_n + \alpha}, \quad y_{n+1} = \frac{y_{n-1}}{y_{n-1}x_n + \beta}.$$

Din et al. [10] studied the global behavior of positive solution to the fourth-order rational difference equations

$$x_{n+1} = \frac{\alpha x_{n-3}}{\beta + \gamma y_n y_{n-1} y_{n-2} y_{n-3}}, \quad y_{n+1} = \frac{\alpha_1 y_{n-3}}{\beta_1 + \gamma_1 x_n x_{n-1} x_{n-2} x_{n-3}}.$$

where the parameters $\alpha, \beta, \gamma, \alpha_1, \beta_1, \gamma_1$ and the initial conditions $x_{-i}, y_{-i}, i = 0, 1, 2, 3$ are positive real numbers.

Although difference equations are sometimes very simple in their forms, they are extremely difficult to understand thoroughly the behavior of their solutions. In book [11], Kocic and Ladas have studied global behavior of nonlinear difference equations of higher order. Similar nonlinear systems of difference equations were investigated (see [12-20]).

Our aim in this paper is to investigate the solutions, stability character and asymptotic behavior of the system of difference equations

$$x_{n+1} = \frac{x_n}{A + y_n y_{n-1}}, \quad y_{n+1} = \frac{y_n}{B + x_n x_{n-1}}, \quad n = 0, 1, \dots, \quad (1.1)$$

where $A, B \in (0, \infty)$ and initial conditions $x_i, y_i \in (0, \infty), i = -1, 0$.

2. Main Results

Let I_x, I_y be some intervals of real number and $f : I_x^2 \times I_y^2 \rightarrow I_x, g : I_x^2 \times I_y^2 \rightarrow I_y$ be continuously differentiable functions. Then for every initial conditions $(x_i, y_i) \in I_x \times I_y (i = -1, 0)$, the system of difference equations

$$\begin{cases} x_{n+1} = f(x_n, x_{n-1}, y_n, y_{n-1}), \\ y_{n+1} = g(x_n, x_{n-1}, y_n, y_{n-1}), \end{cases} \quad n = 0, 1, 2, \dots, \quad (2.1)$$

has a unique solution $\{(x_n, y_n)\}_{n=-1}^\infty$. A point $(\bar{x}, \bar{y}) \in I_x \times I_y$ is called an equilibrium point of (2.1) if $\bar{x} = f(\bar{x}, \bar{x}, \bar{y}, \bar{y}), \bar{y} = g(\bar{x}, \bar{x}, \bar{y}, \bar{y})$, i. e., $(x_n, y_n) = (\bar{x}, \bar{y})$ for all $n \geq 0$.

Definition 2.1 Assume that $(\bar{x}, \bar{y})$ be a fixed point of (2.1). Then

- (i) $(\bar{x}, \bar{y})$ is said to be stable relative to $I_x \times I_y$ if for every $\varepsilon > 0$, there exists $\delta > 0$ such that for any initial conditions $(x_i, y_i) \in I_x \times I_y (i = -1, 0)$, with $\sum_{i=-1}^0 |x_i - \bar{x}| < \delta, \sum_{i=-1}^0 |y_i - \bar{y}| < \delta$, implies $|x_n - \bar{x}| < \varepsilon, |y_n - \bar{y}| < \varepsilon$.
- (ii) $(\bar{x}, \bar{y})$ is called an attractor relative to $I_x \times I_y$ if for all $(x_i, y_i) \in I_x \times I_y (i = -1, 0)$, $\lim_{n \rightarrow \infty} x_n = \bar{x}, \lim_{n \rightarrow \infty} y_n = \bar{y}$.
- (iii) $(\bar{x}, \bar{y})$ is called asymptotically stable relative to $I_x \times I_y$ if it is stable and an attractor.
- (iv) Unstable if it is not stable.

Theorem 2.1[11] Assume that $X(n+1) = F(X(n)), n = 0, 1, \dots$, is a system of difference equations and $\bar{X}$ is the equilibrium point of this system i.e., $F(\bar{X}) = \bar{X}$. If all eigenvalues of the Jacobian matrix J_F , evaluated at $\bar{X}$ lie inside the open unit disk $|\lambda| < 1$, then $\bar{X}$ is locally asymptotically stable. If one of them has modulus greater than one then $\bar{X}$ is unstable.

Clearly, system (1.1) has always trivial equilibrium $(0, 0)$. If $0 < A < 1, 0 < B < 1$, system (1.1) has a unique positive equilibrium point $(\sqrt{1-B}, \sqrt{1-A})$.

Theorem 2.2. Consider system (1.1), then the following statements are true:

- (i) If $0 < A < 1, 0 < B < 1$, then the trivial equilibrium point $(0, 0)$ is local unstable.
- (ii) If $A > 1, B > 1$, then the trivial equilibrium point $(0, 0)$ is locally asymptotically stable.

Proof. We can obtain easily the linearized system of (1.1) about the positive equilibrium $(0, 0)$ is

$$\Phi_{n+1} = B\Phi_n \quad (2.2)$$

where

$$\Phi_n = \begin{pmatrix} x_n \\ x_{n-1} \\ y_n \\ y_{n-1} \end{pmatrix}, \quad B = \begin{pmatrix} \frac{1}{A} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{B} & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

The characteristic equation of (2.2) is

$$\lambda^2(\lambda - \frac{1}{A})(\lambda - \frac{1}{B}) = 0 \quad (2.3)$$

(i) If $0 < A < 1, 0 < B < 1$, This shows that two roots of characteristic equation lie outside unit disk. So the trivial equilibrium point $(0, 0)$ is a repel, i.e. it is local unstable.

(ii) If $A > 1, B > 1$, The all root of characteristic equation lie inside unit disk. So the trivial equilibrium point $(0, 0)$ is a source, i.e. it is locally asymptotically stable.

Theorem 2.3 Suppose that $A > 1, B > 1$ holds, then the trivial equilibrium point $(0, 0)$ is globally asymptotically stable.

Proof. For $A > 1, B > 1$, from (ii) of Theorem 2.2, the equilibrium point $(0, 0)$ is locally asymptotically stable. From (1.1), it is easy to see every positive solution (x_n, y_n) is bounded. i.e. $0 < x_n < x_0, 0 < y_n < y_0$. Now it is sufficient to prove (x_n, y_n) is decreasing. From (1.1), we have

$$\frac{x_{n+1}}{x_n} = \frac{1}{A + y_n y_{n-1}} \leq \frac{1}{A} < 1, \quad \frac{y_{n+1}}{y_n} = \frac{1}{B + x_n x_{n-1}} \leq \frac{1}{B} < 1.$$

This implies that the sequences $\{x_n\}$ and $\{y_n\}$ are decreasing. Hence $\lim_{n \rightarrow \infty} x_n = 0, \lim_{n \rightarrow \infty} y_n = 0$. Therefore the trivial equilibrium point $(0, 0)$ is globally asymptotically stable.

Theorem 2.4 Assume that $0 < A < 1, 0 < B < 1$. Then the positive equilibrium point $(\bar{x}, \bar{y}) = (\sqrt{1-B}, \sqrt{1-A})$ is locally unstable.

Proof. We can obtain easily the linearized system of (1.1) about the positive equilibrium $(\bar{x}, \bar{y})$ is

$$\Phi_{n+1} = G\Phi_n \quad (2.4)$$

where $\Phi_n = (x_n, x_{n-1}, y_n, y_{n-1})^T$,

$$G = \begin{pmatrix} \frac{1}{A} & 0 & -\bar{x}\bar{y} & -\bar{x}\bar{y} \\ 1 & 0 & 0 & 0 \\ -\bar{x}\bar{y} & -\bar{x}\bar{y} & \frac{1}{B} & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

Let $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ denote the 4 eigenvalues of Matrix G . Let $D = \text{diag}(d_1, d_2, d_3, d_4), d_i \neq 0 (i = 1, 2, 3, 4)$ be a diagonal matrix,

Clearly D is invertible. Computing DGD^{-1} , we obtained

$$DGD^{-1} = \begin{pmatrix} \frac{1}{A} & 0 & -\frac{d_1}{d_3}\bar{x}\bar{y} & -\frac{d_1}{d_4}\bar{x}\bar{y} \\ \frac{d_2}{d_1} & 0 & 0 & 0 \\ -\frac{d_3}{d_1}\bar{x}\bar{y} & -\frac{d_3}{d_2}\bar{x}\bar{y} & \frac{1}{B} & 0 \\ 0 & 0 & \frac{d_4}{d_3} & 0 \end{pmatrix}$$

It is well known that G has the same eigenvalues as DGD^{-1} , we obtain that

$$\begin{aligned} \max_{1 \leq k \leq 4} |\lambda_k| &= \|DGD^{-1}\| \\ &= \max \left\{ d_2 d_1^{-1}, d_4 d_3^{-1}, \frac{1}{A} + \frac{d_1}{d_3}\bar{x}\bar{y} + \frac{d_1}{d_4}\bar{x}\bar{y}, \right. \\ &\quad \left. \frac{1}{B} + \frac{d_3}{d_1}\bar{x}\bar{y} + \frac{d_3}{d_2}\bar{x}\bar{y} \right\} \\ &> 1 \end{aligned}$$

It follows from Theorem 2.1 [11] that the positive equilibrium points $(\bar{x}, \bar{y})$ is locally unstable.

3. Rate of convergence

In this section we will determine the rate of convergence of a solution that converges to the equilibrium point $(0,0)$ of the system (1.1). The following result gives the rate of convergence of solution of a system of difference equations

$$X_{n+1} = [P + Q(n)]X_n \quad (3.1)$$

where X_n is a four dimensional vector, $P \in C^{4 \times 4}$ is a constant matrix, $Q : Z^+ \rightarrow C^{4 \times 4}$ is a matrix function satisfying

$$\|Q(n)\| \rightarrow 0, \text{ when } n \rightarrow \infty. \quad (3.2)$$

where $\|\cdot\|$ denotes any matrix norm which is associated with the vector norm.

Theorem 3.1.[21] *Assume that condition (3.2) hold, if X_n is a solution of (3.1), then either $X_n = 0$ for all large n or*

$$\rho = \lim_{n \rightarrow \infty} \sqrt[n]{\|X_n\|} \quad (3.3)$$

or

$$\rho = \lim_{n \rightarrow \infty} \frac{\|X_{n+1}\|}{\|X_n\|} \quad (3.4)$$

exists and is equal to the moduls of one the eigenvalues of the matrix P .

Assume that $\lim_{n \rightarrow \infty} x_n = \bar{x}$, $\lim_{n \rightarrow \infty} y_n = \bar{y}$, we will find a system of limiting equations for the system (1.1). The error terms are given as

$$\begin{cases} x_{n+1} - \bar{x} = \sum_{i=0}^1 P_i(x_{n-i} - \bar{x}) + \sum_{i=0}^1 Q_i(y_{n-i} - \bar{y}) \\ y_{n+1} - \bar{y} = \sum_{i=0}^1 R_i(x_{n-i} - \bar{x}) + \sum_{i=0}^1 S_i(y_{n-i} - \bar{y}) \end{cases}$$

Set $e_n^1 = x_n - \bar{x}$, $e_n^2 = y_n - \bar{y}$, therefore it follows that

$$\begin{cases} e_{n+1}^1 = \sum_{i=0}^1 P_i e_{n-i}^1 + \sum_{i=0}^1 Q_i e_{n-i}^2 \\ e_{n+1}^2 = \sum_{i=0}^1 R_i e_{n-i}^1 + \sum_{i=0}^1 S_i e_{n-i}^2 \end{cases}$$

where

$$P_0 = \frac{1}{A + y_n y_{n-1}}, P_1 = 0, Q_0 = -\frac{x_n y_{n-1}}{(A + y_n y_{n-1})^2}, Q_1 = -\frac{x_n y_n}{(A + y_n y_{n-1})^2}.$$

$$R_0 = -\frac{x_{n-1}y_n}{(B + x_nx_{n-1})^2}, R_1 = -\frac{x_ny_n}{(B + x_nx_{n-1})^2}, S_0 = \frac{1}{B + x_nx_{n-1}}, S_1 = 0.$$

Now it is clear that

$$\lim_{n \rightarrow \infty} P_0 = \frac{1}{A}, \lim_{n \rightarrow \infty} Q_0 = \lim_{n \rightarrow \infty} Q_1 = 0, \lim_{n \rightarrow \infty} R_0 = \lim_{n \rightarrow \infty} R_1 = 0, \lim_{n \rightarrow \infty} S_0 = \frac{1}{B}.$$

Hence, the limiting system of error terms at $(0, 0)$ can be written as

$$E_{n+1} = GE_n \quad (3.5)$$

where $E_n = (e_n^1, e_{n-1}^1, e_n^2, e_{n-1}^2)^T$, and

$$G = J_F(0, 0) = (d_{ij})_{4 \times 4} = \begin{pmatrix} \frac{1}{A} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{B} & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

Using Theorem 3.1, we have the following result

Theorem 3.2. *Assume that $A > 1, B > 1$, and $\{(x_n, y_n)\}$ be a positive solution of the system (1.1). Then, the error vector E_n of every solution of (1.1) satisfies both of the following asymptotic relations*

$$\lim_{n \rightarrow \infty} \sqrt[n]{\|E_n\|} = |\lambda J_F(0, 0)|, \lim_{n \rightarrow \infty} \frac{\|E_{n+1}\|}{\|E_n\|} = |\lambda J_F(0, 0)|.$$

where $\lambda J_F(0, 0)$ is equal to the moduls of one the eigenvalues of the Jacobian matrix evaluted at the equilibrium $(0, 0)$.

4. Numerical examples

In order to illustrate the results of the previous sections and to support our theoretical discussions, we consider an interesting numerical example in this section.

Example 4.1. *Consider the system (1.1) with initial conditions $x_{-1} = 0.8, x_0 = 1.2, y_{-1} = 1.8, y_0 = 2.2$. Moreover, choosing the parameters $A = 1.7, B = 1.5$. Then system (1.1) can be written as*

$$x_{n+1} = \frac{x_n}{1.7 + y_n y_{n-1}}, \quad y_{n+1} = \frac{y_n}{1.5 + x_n x_{n-1}} \quad (4.1)$$

The plot of system (4.1) is shown in Fig. 1.




Fig.1. The plot of system (4.1).

Example 4.2. Consider the system (1.1) with initial conditions $x_{-1} = 4.8, x_0 = 4.6, y_{-1} = 5.8, y_0 = 5.2$, Moreover, choosing the parameters $A = 0.7, B = 0.8$. Then system (1.1) can be written as

$$x_{n+1} = \frac{x_n}{0.7 + y_n y_{n-1}}, \quad y_{n+1} = \frac{y_n}{0.8 + x_n x_{n-1}} \quad (4.2)$$

The plot of system (4.2) is shown in Fig. 2.




Fig.2. The plot of system (4.2).

5. Conclusions

In this paper, the dynamical behavior of second-order discrete system is studied. It conclude that: (i) the positive equilibrium point $(\sqrt{1-B}, \sqrt{1-A})$ is locally unstable if $0 < A < 1, 0 < B < 1$. (ii) the trivial

equilibrium point $(0,0)$ is globally asymptotically stable if $A > 1, B > 1$. Some numerical examples are provided to support theoretical results.

Competing interests

The author declares that she has no competing interests exist.

References

- [1] G. Papaschinopoulos and C. J. Schinas, On a system of two nonlinear difference equations, *J. Math. Anal. Appl.* 219(1998), 415–426.
- [2] D. Clark, M. R. S. Kulenovic and J. F. Selgrade, Global asymptotic behavior of a two-dimensional difference equation modelling competition, *Nonlinear Analysis* 52 (2003), 1765–1776.
- [3] A. S. Kurbanli, On the behavior of solutions of the system of rational difference equations: $x_{n+1} = x_{n-1}/(y_n x_{n-1} - 1)$, $Y_{n+1} = y_{n-1}/(x_n y_{n-1} - 1)$ and $z_{n+1} = z_{n-1}/(y_n z_{n-1} - 1)$, *Discrete Dynamics in Nature and Society*, vol. 2011, Article ID 932632, 12 pages, 2011.
- [4] C. Cinar, I. Yalcinkaya and R. Karatas, On the positive solutions of the difference equation system $x_{n+1} = \frac{m}{y_n}$, $y_{n+1} = \frac{py_n}{x_{n-1}y_{n-1}}$, *J. Inst. Math. Comp. Sci.*, 18(2005), 135–136,
- [5] C. Cinar, On the positive solutions of the difference equation system $x_{n+1} = \frac{1}{y_n}$, $y_{n+1} = \frac{y_n}{x_{n-1}y_{n-1}}$, *Applied Mathematics and Computation*, 158(2004) 303–305.
- [6] D. Clark and M. R. S. Kulenovic, A coupled system of rational difference equations, *Computers and Mathematics with Applications* 43 (2002), 849–867.
- [7] Q. Zhang, L. Yang and J. Liu, Dynamics of a system of rational third-order difference equation, *Advances in Difference Equations* 2012, 2012:136, 1–8.
- [8] Q. Zhang, J. Liu, Z. Luo, Dynamical behavior of a system of third-order rational difference equation, Hindawi Publishing Corporation, *Discrete Dynamics in Nature and Society* Article ID 530453, 1–6.
- [9] T. F. Ibrahim, Two-dimensional fractional system of nonlinear difference equations in the modeling competitive populations, *International Journal of Basic and Applied Sciences* 12(2012), No: 05, 103–121.
- [10] Q. Din, MN Qureshi and A. Q. Khan, Dynamics of a fourth-order system of rational difference equations, *Advances in Difference Equations* 2012(2012), 215, pp. 1–15.
- [11] V. L. Kocic and G. Ladas, Global behavior of nonlinear difference equations of higher order with application, Kluwer Academic Publishers, Dordrecht, 1993.
- [12] K. Liu, Z. Zhao, X. Li and P. Li, More on three-dimensional systems of rational difference equations, *Discrete Dynamics in Nature and Society*, Volume 2011, Article ID 178483, 2011.
- [13] T. F. Ibrahim and Q. Zhang, Stability of an anti-competitive system of rational difference equations, *Archives Des Sciences*, 66 (2013), No. 5, 44–58.
- [14] E. M. E. Zayed and M. A. El-Moneam, On the global attractivity of two nonlinear difference equations, *J. Math. Sci.*, 177 (2011), 487–499.
- [15] N. Touafek and E. M. Elsayed, On the periodicity of some systems of nonlinear difference equations, *Bull. Math. Soc. Sci. Math. Roumanie*, 2(2012), 217–224.
- [16] N. Touafek and E. M. Elsayed, On the solutions of systems of rational difference equations, *Mathematical and Computer Modelling*, 55(2012), 1987–1997.
- [17] S. Kalabusic, M. R. S. Kulenovic and E. Pilav, Dynamics of a two-dimensional system of rational difference equations of Leslie–Gower type, *Advances in Difference Equations*, 2011, doi:10.1186/1687-1847-2011-29.

- [18] T. F. Ibrahim, Boundedness and stability of a rational difference equation with delay, *Rev. Roumaine Math.Pures Appl.*, Vol. 57 (2012), no.3, 215–224.
- [19] T. F. Ibrahim and N. Touafek, On a third-order rational difference equation with variable coefficients, *DCDIS Series B: Applications & Algorithms*, Vol. 20 (2013), no. 2, 251–264.
- [20] T. F. Ibrahim, Oscillation, non-oscillation, and asymptotic behavior for third order nonlinear giffer-
ence equations, *Dynamics of Continuous, Discrete and Impulsive Systems, Series A: Mathematical
Analysis*, Vol. 20 (2013), no. 4, 523–532.
- [21] M. Pituk, More on Poincare’s and Peron’s theorems for difference equations, *Journal Difference
Equations and Applications*, 8 (2002), 201–216.