Original Research Article

There is no standard model of ZFC

Abstract: Main results are:(i) $\neg Con(ZFC + \exists M_{st}^{ZFC})$,

(ii) let k be an inaccessible cardinal then $\neg Con(ZFC + \exists \kappa)$ [10]-[11].

Keywords: Gödel encoding,Russell's paradox, standard model, Henkin semantics, strongly inaccessible cardinal.

I.Introduction.

1.1.Main results.

Let us remind that accordingly to naive set theory, any definable collection is a set. Let R be the set of all sets that are not members of themselves. If R qualifies as a member of itself, it would contradict its own definition as a set containing all sets that are not members of themselves. On the other hand, if such a set is not a member of itself, it would qualify as a member of itself by the same definition. This contradiction is Russell's paradox. In 1908, two ways of avoiding the paradox were proposed, Russell's type theory and Zermelo set theory, the first constructed axiomatic set theory. Zermelo's axioms went well beyond Frege's axioms of extensionality and unlimited set abstraction, and evolved into the now-canonical Zermelo–Fraenkel set theory *ZFC*.*"But how do we know that ZFC is a consistent theory, free of contradictions? The short answer is that we don't; it is a matter of faith (or of skepticism)"*— E.Nelson wrote in his paper [1]. However, it is deemed unlikely that even ZFC_2 which is significantly stronger than ZFC harbors an unsuspected contradiction; it is widely believed that if ZFC and ZFC_2 were inconsistent, that fact would have been uncovered by now. This much is certain —ZFC and ZFC_2 is immune to the classic paradoxes of naive set theory: Russell's paradox, the Burali-Forti paradox, and Cantor's paradox.

Remark 1.1.1.Note that in this paper we view (i) the first order set theory ZFC under the

canonical first order semantics (ii) the second order set theory ZFC_2 under the Henkin

semantics [2],[3],[4],[5],[6].

Remark 1.1.2. Second-order logic essantially differs from the usual first-order predicate calculus in that it has variables and quantifiers not only for individuals but also for subsets of the universe and variables for n-ary relations as well [2],[6].The deductive calculus $\mathbf{DED}_2$ of second order logic is based on rules and axioms which guarantee that the quantifiers range at least over definable subsets [6]. As to the semantics, there are two tipes of models: (i) Suppose $\mathbf{U}$ is an ordinary first-order structure and $\mathbf{S}$ is a set of subsets of the domain A of $\mathbf{U}$. The main idea is that the set-variables range over $\mathbf{S}$,i.e. $\langle \mathbf{U}, \mathbf{S} \rangle \models \exists X \Phi(X) \Leftrightarrow \exists S(S \in \mathbf{S})[\langle \mathbf{U}, \mathbf{S} \rangle \models \Phi(S)]$.

We call $\langle \mathbf{U}, \mathbf{S} \rangle$ a Henkin model, if $\langle \mathbf{U}, \mathbf{S} \rangle$ satisfies the axioms of $\mathbf{DED}_2$ and truth in $\langle \mathbf{U}, \mathbf{S} \rangle$ is preserved by the rules of $\mathbf{DED}_2$. We call this semantics of second-order logic the Henkin semantics and second-order logic with the Henkin semantics the Henkin second-order logic. There is a special class of Henkin models, namely those $\langle \mathbf{U}, \mathbf{S} \rangle$ where $\mathbf{S}$ is the set of all subsets of A. We call these full models. We call this semantics of second-order logic the full semantics and second-order logic with the full semantics the full second-order logic.

Remark 1.1.3. We emphasize that the following facts are the main features of second-order logic:

1.The Completeness Theorem: A sentence is provable in $\mathbf{DED}_2$ if and only if it holds in all Henkin models [2],[6].

2.The Löwenheim-Skolem Theorem: A sentence with an infinite Henkin model has a countable Henkin model.

3.The Compactness Theorem: A set of sentences, every finite subset of which has a Henkin model, has itself a Henkin model.

4.The Incompleteness Theorem: Neither $\mathbf{DED}_2$ nor any other effectively given deductive calculus is complete for full models, that is, there are always sentences which are true in all full models but which are unprovable.

5.Failure of the Compactness Theorem for full models.

6.Failure of the Löwenheim-Skolem Theorem for full models.

7.There is a finite second-order axiom system $\mathbb{Z}_2$ such that the semiring $\mathbb{N}$ of natural numbers is the only full model (up to isomorphism) of $\mathbb{Z}_2$.

8. There is a finite second-order axiom system RCF_2 such that the field $\mathbb{R}$ of real numbers is the only (up to isomorphism) full model of RCF_2.

Remark 1.1.4. For let second-order ZFC be, as usual, the theory that results obtained from ZFC when the axiom schema of replacement is replaced by its second-order universal closure,i.e.

$$\forall X[Func(X) \Rightarrow \forall u \exists v \forall r[r \in v \Leftrightarrow \exists s(s \in u \wedge (s,r) \in X)]], \qquad (1.1.1)$$

where X is a second-order variable, and where $Func(X)$ abbreviates " X is a functional relation",see [7].

Designation 1.1.1. We will denote (i) by ZFC_2^{Hs} set theory ZFC_2 with the Henkin semantics, (ii) by $\overline{ZFC}_2^{Hs}$ set theory $ZFC_2^{Hs} + \exists M_{st}^{ZFC_2^{Hs}}$ and (iii) by ZFC_{st} set theory $ZFC + \exists M_{st}^{ZFC}$, where M_{st}^{Th} is a standard model of the theory Th.

Axiom $\exists M^{ZFC}$.[8]. There is a set M^{ZFC} and a binary relation $\epsilon \subseteq M^{ZFC} \times M^{ZFC}$ which makes M^{ZFC} a model for ZFC.

Remark 1.1.3.(i)We emphasize that it is well known that axiom $\exists M^{ZFC}$ a single statement in ZFC see [8],Ch.II,section 7.We denote this statement throught all this paper by symbol $Con(ZFC; M^{ZFC})$.The completness theorem says that $\exists M^{ZFC} \Leftrightarrow Con(ZFC)$.

(ii) Obviously there exists a single statement in ZFC_2^{Hs} such that $\exists M^{ZFC_2^{Hs}} \Leftrightarrow Con(ZFC_2^{Hs})$.

We denote this statement throught all this paper by symbol $Con\left(ZFC_2^{Hs}; M^{ZFC_2^{Hs}}\right)$ and there exists a single statement $\exists M^{Z_2^{Hs}}$ in Z_2^{Hs}. We denote this statement throught all this paper by symbol $Con\left(Z_2^{Hs}; M^{Z_2^{Hs}}\right)$.

Axiom $\exists M_{st}^{ZFC}$.[8].There is a set M_{st}^{ZFC} such that if R is $\{\langle x,y\rangle | x \in y \wedge x \in M_{st}^{ZFC} \wedge y \in M_{st}^{ZFC}\}$ then M_{st}^{ZFC} is a model for ZFC under the relation R.

Definition 1.1.1.[8].The model M_{st}^{ZFC} and $M_{st}^{Z_2^{Hs}}$ is called a standard model since the relation $\in$ used is merely the standard $\in$- relation.

Remark 1.1.4.[8].Note that axiom $\exists M^{ZFC}$ doesn't imply axiom $\exists M_{st}^{ZFC}$.

Remark 1.1.6.Note that in order to deduce: (i) $\sim Con(ZFC_2^{Hs})$ from $Con(ZFC_2^{Hs})$, and (ii) $\sim Con(ZFC)$ from $Con(ZFC)$, by using Gödel encoding, one needs something more than the consistency of ZFC_2^{Hs}, e.g., that ZFC_2^{Hs} has an omega-model $M_{\omega}^{ZFC_2^{Hs}}$ or an standard model $M_{st}^{ZFC_2^{Hs}}$ i.e., a model in which the *integers are the standard integers*.To put it another way, why should we believe a statement just because there's a ZFC_2^{Hs}-proof of it? It's clear that if ZFC_2^{Hs} is inconsistent, then we won't believe ZFC_2^{Hs}-proofs. What's slightly more subtle is that the mere consistency of ZFC_2 isn't quite enough to get us to believe arithmetical theorems of ZFC_2^{Hs}; we must also believe that these arithmetical theorems are asserting something about the standard naturals. It is "conceivable" that ZFC_2^{Hs} might be consistent but that the only nonstandard models $M_{Nst}^{ZFC_2^{Hs}}$ it has are those in which the integers are nonstandard, in which case we might not "believe" an arithmetical statement such as "ZFC_2^{Hs} is

inconsistent" even if there is a ZFC_2^{Hs}-proof of it.

2.Derivation of the inconsistent definable set in set theory $\overline{ZFC}_2^{Hs}$ and in set theory ZFC_{st}.

2.1.Derivation of the inconsistent definable set in set theory $\overline{ZFC}_2^{Hs}$.

We assume now that $Con\left(Z_2^{Hs}; M_{st}^{Z_2^{Hs}}\right)$.

Designation 2.1.1.Let Γ_X be the collection of the all 1-place open wff of the set theory
$\overline{ZFC}_2^{Hs}$.

Definition 2.1.1.Let $\Psi_1(X), \Psi_2(X)$ be 1-place open wff's of the set theory $\overline{ZFC}_2^{Hs}$.

(i) We define now the equivalence relation $(\bullet \sim_X \bullet) \subset \Gamma_X \times \Gamma_X$ by

$$\Psi_1(X) \sim \Psi_2(X) \Leftrightarrow \forall X[\Psi_1(X) \Leftrightarrow \Psi_2(X)] \tag{2.1.1}$$

(ii) A subset Λ_X^{Hs} of $\Gamma(X)$ such that $\Psi_1(X) \sim \Psi_2(X)$ holds for all $\Psi_1(X)$ and $\Psi_2(X)$ in $\Lambda(X)$,
and never for $\Psi_1(X)$ in $\Lambda(X)$ and $\Psi_2(X)$ outside $\Lambda(X)$, is called an equivalence class of
$\Gamma(X)$.

(iii)The collection of all possible equivalence classes of $\Gamma(X)$ by $\sim$, denoted $\Gamma(X)/\sim_X$

$$\Gamma_X/\sim_X \triangleq \{[\Psi(X)]_{Hs} | \Psi(X) \in \Gamma(X)\}. \tag{2.1.2}$$

(iv) For any $\Psi(X) \in \Gamma(X)$ let $[\Psi(X)] \triangleq \{\Phi(X) \in \Gamma(X) | \Psi(X) \sim \Phi(X)\}$ denote the
equivalence class to which $\Psi(X)$ belongs. All elements of $\Gamma(X)$ equivalent to each other
are also elements of the same equivalence class.

Definition 2.1.2.[9].Let Th be any theory in the recursive language $\mathcal{L}_{Th} \supset \mathcal{L}_{PA}$,where $\mathcal{L}_{PA}$
is a language of Peano arithmetic.We say that a number-theoretic relation $R(x_1,\ldots,x_n)$ of
n arguments is expressible in Th if and only if there is a wff $\widehat{R}(x_1,\ldots,x_n)$ of Th with the free
variables $x_1,\ldots,x_n$ such that,for any natural numbers $k_1,\ldots,k_n$, the following hold:

(i) If $R(k_1,\ldots,k_n)$ is true, then $\vdash_{Th} \widehat{R}(\bar{k}_1,\ldots,\bar{k}_n)$.

(ii) If $R(k_1,\ldots,k_n)$ is false, then $\vdash_{Th} \neg\widehat{R}(\bar{k}_1,\ldots,\bar{k}_n)$.

Designation 2.1.2.(i) Let $g_{ZFC_2^{Hs}}(u)$ be a Gödel number of given an expression u of
the set theory $\overline{ZFC}_2^{Hs} \triangleq \quad ZFC_2^{Hs} + \exists M_{st}^{ZFC_2^{Hs}}$.

(ii) Let $\mathbf{Fr}_2^{Hs}(y,v)$ be the relation : y is the Gödel number of a wff of the set theory$\overline{ZFC}_2^{Hs}$

that contains free occurrences of the variable X with Gödel number v [8]-[9].

(iii) Note that the relation $\mathbf{Fr}_2^{Hs}(y,v)$ is expressible in $\overline{ZFC}_2^{Hs}$ by a wff $\widehat{\mathbf{Fr}_2^{Hs}}(y,v)$

(iv) Note that for any $y,v \in \mathbb{N}$ by definition of the relation $\mathbf{Fr}_2^{Hs}(y,v)$ follows that

$$\widehat{\mathbf{Fr}_2^{Hs}}(y,v) \Leftrightarrow \exists!\Psi(X)\Big[\Big(g_{\overline{ZFC}_2^{Hs}}(\Psi(X)) = y\Big) \wedge \Big(g_{\overline{ZFC}_2^{Hs}}(X) = v\Big)\Big], \qquad (2.1.3)$$

where $\Psi(X)$ is a unique wff of $\overline{ZFC}_2^{Hs}$ which contains free occurrences of the variable X

with Gödel number v. We denote a unique wff $\Psi(X)$ defined by using equivalence (1.2.3)

by symbol $\Psi_{y,v}(X)$, i.e.

$$\widehat{\mathbf{Fr}_2^{Hs}}(y,v) \Leftrightarrow \exists!\Psi_{y,v}(X)\Big[\Big(g_{\overline{ZFC}_2^{Hs}}(\Psi_{y,v}(X)) = y\Big) \wedge \Big(g_{\overline{ZFC}_2^{Hs}}(X) = v\Big)\Big], \qquad (2.1.4)$$

(v) Let $\wp_2^{Hs}(y,v,v_1)$ be a Gödel number of the following wff: $\exists!X[\Psi(X) \wedge Y = X]$, where

$g_{\overline{ZFC}_2^{Hs}}(\Psi(X)) = y, g_{\overline{ZFC}_2^{Hs}}(X) = v,\ g_{\overline{ZFC}_2^{Hs}}(Y) = v_1.$

(vi) Let $\mathrm{Pr}_{ZFC_2^{Hs}}(z)$ be a predicate asserting provability in $\overline{ZFC}_2^{Hs}$, which defined by formula

(2.6) in section 2, see Remark 2.2 and Designation 2.3,(see also [9]-[10]).

Definition 2.1.3. Let Γ_X^{Hs} be the countable collection of the all 1-place open wff's of the set theory$\overline{ZFC}_2^{Hs}$ that contains free occurrences of the variable X.

Definition 2.1.4. Let $g_{\overline{ZFC}_2^{Hs}}(X) = v$. Let Γ_v^{Hs} be a set of the all Gödel numbers of the 1-place open wff's of the set theory$\overline{ZFC}_2^{Hs}$ that contains free occurrences of the variable X

with Gödel number v, i.e.

$$\Gamma_v^{Hs} = \{y \in \mathbb{N} | \langle y,v\rangle \in \mathbf{Fr}_2^{Hs}(y,v)\}, \qquad (2.1.5)$$

or in the following equivalent form:

$$\forall y(y \in \mathbb{N})\Big[\, y \in \Gamma_v \Leftrightarrow (y \in \mathbb{N}) \wedge \widehat{\mathbf{Fr}_2^{Hs}}(y,v)\,\Big].$$

Remark 2.1.1.Note that from the axiom of separation it follows directly that Γ_v^{Hs} is a set

in the sense of the set theory $\overline{ZFC}_2^{Hs}$.

Definition 2.1.5.(i)We define now the equivalence relation $(\bullet \sim_X \bullet) \subset \Gamma_X^{Hs} \times \Gamma_X^{Hs}$ by

$$\Psi_1(X) \sim_X \Psi_2(X) \Leftrightarrow (\forall X[\Psi_1(X) \Leftrightarrow \Psi_2(X)]) \qquad (2.1.6)$$

(ii) A subcollection Λ_X^{Hs} of Γ_X^{Hs} such that $\Psi_1(X) \sim_X \Psi_2(X)$ holds for all $\Psi_1(X)$ and $\Psi_2(X)$ in

Λ_X^{Hs}, and never for $\Psi_1(X)$ in Λ_X^{Hs} and $\Psi_2(X)$ outside Λ_X^{Hs}, is an equivalence class of

Γ_X^{Hs}.

(iii) For any $\Psi(X) \in \Gamma_X^{Hs}$ let $[\Psi(X)]_{Hs} \triangleq \{\Phi(X) \in \Gamma_X^{Hs} | \Psi(X) \sim_X \Phi(X)\}$ denote the equivalence class to which $\Psi(X)$ belongs. All elements of Γ_X^{Hs} equivalent to each other are also elements of the same equivalence class.

(iv) The collection of all possible equivalence classes of Γ_X^{Hs} by $\sim_X$, denoted $\Gamma_X^{Hs}/\sim_X$

$$\Gamma_X^{Hs}/\sim_X \triangleq \{[\Psi(X)]_{Hs} | \Psi(X) \in \Gamma_X^{Hs}\}. \tag{2.1.7}$$

Definition 2.1.6.(i)We define now the equivalence relation $(\bullet \sim_v \bullet) \subset \Gamma_v^{Hs} \times \Gamma_v^{Hs}$ in the sense of the set theory $\overline{ZFC}_2^{Hs}$ by

$$y_1 \sim_v y_2 \Leftrightarrow (\forall X[\Psi_{y_1,v}(X) \Leftrightarrow \Psi_{y_2,v}(X)]) \tag{2.1.8}$$

Note that from the axiom of separation it follows directly that the equivalence relation $(\bullet \sim_v \bullet)$ is a relation in the sense of the set theory $\overline{ZFC}_2^{Hs}$.

(ii) A subset Λ_v^{Hs} of Γ_v^{Hs} such that $y_1 \sim_v y_2$ holds for all y_1 and y_1 in Λ_v^{Hs},and never for y_1 in Λ_v^{Hs} and y_2 outside Λ_v^{Hs}, is an equivalence class of Γ_v^{Hs}.

(iii) For any $y \in \Gamma_v^{Hs}$ let $[y]_{Hs} \triangleq \{z \in \Gamma_v^{Hs} | y \sim_v z\}$ denote the equivalence class to which y belongs. All elements of Γ_v^{Hs} equivalent to each other are also elements of the same equivalence class.

(iv)The collection of all possible equivalence classes of Γ_v^{Hs} by $\sim_v$, denoted $\Gamma_v^{Hs}/\sim_v$

$$\Gamma_v^{Hs}/\sim_v \triangleq \{[y]_{Hs} | y \in \Gamma_v^{Hs}\}. \tag{2.1.9}$$

Remark 2.1.2. Note that from the axiom of separation it follows directly that $\Gamma_v^{Hs}/\sim_v$ is a set in the sense of the set theory $\overline{ZFC}_2^{Hs}$.

Definition 2.1.7.Let $\Im_2^{Hs}$ be the countable collection of the all sets definable by 1-place open wff of the set theory $\overline{ZFC}_2^{Hs}$, i.e.

$$\forall Y\{Y \in \Im_2^{Hs} \Leftrightarrow \exists \Psi(X)[([\Psi(X)]_{Hs} \in \Gamma_X^{Hs}/\sim_X) \wedge [\exists! X[\Psi(X) \wedge Y = X]]]\}. \tag{2.1.10}$$

Definition 2.1.8.We rewrite now (2.1.10) in the following equivalent form

$$\forall Y\{Y \in \Im_2^{Hs} \Leftrightarrow \exists \Psi(X)[([\Psi(X)]_{Hs} \in \Gamma_X^{*Hs}/\sim_X) \wedge (Y = X)]\}, \tag{2.1.11}$$

where the countable collection $\Gamma_X^{*Hs}/\sim_X$ is defined by

$$\forall \Psi(X)\{[\Psi(X)] \in \Gamma_X^{*Hs}/\sim_X \Leftrightarrow [([\Psi(X)] \in \Gamma_X^{Hs}/\sim_X) \wedge \exists! X\Psi(X)]\} \tag{2.1.12}$$

Definition 2.1.9. Let $\Re_2^{Hs}$ be the countable collection of the all sets such that

$$\forall X(X \in \Im_2^{Hs})[X \in \Re_2^{Hs} \Leftrightarrow X \notin X]. \tag{2.1.13}$$

Remark 2.1.3. Note that $\Re_2^{Hs} \in \Im_2^{Hs}$ since $\Re_2^{Hs}$ is a collection definable by 1-place

open wff

$$\Psi(Z,\mathfrak{I}_2^{Hs}) \triangleq \forall X(X \in \mathfrak{I}_2^{Hs})[X \in Z \Leftrightarrow X \notin X].$$

From (2.1.13) one obtains

$$\mathfrak{R}_2^{Hs} \in \mathfrak{R}_2^{Hs} \Leftrightarrow \mathfrak{R}_2^{Hs} \notin \mathfrak{R}_2^{Hs}. \tag{2.1.14}$$

But (2.1.14) gives a contradiction

$$(\mathfrak{R}_2^{Hs} \in \mathfrak{R}_2^{Hs}) \wedge (\mathfrak{R}_2^{Hs} \notin \mathfrak{R}_2^{Hs}). \tag{2.1.15}$$

However contradiction (2.1.15) it is not a contradiction inside $\overline{ZFC}_2^{Hs}$ for the reason that

the countable collection $\mathfrak{I}_2^{Hs}$ is not a set in the sense of the set theory $\overline{ZFC}_2^{Hs}$.

In order to obtain a contradiction inside $\overline{ZFC}_2^{Hs}$ we introduce the following definitions.

Definition 2.1.10.We define now the countable set $\Gamma_v^{*Hs}/\sim_v$ by

$$\forall y\left\{[y]_{Hs} \in \Gamma_v^{*Hs}/\sim_v \Leftrightarrow ([y]_{Hs} \in \Gamma_v^{Hs}/\sim_v) \wedge \widehat{\mathbf{Fr}_2^{Hs}}(y,v) \wedge [\exists! X \Psi_{y,v}(X)]\right\}. \tag{2.1.16}$$

Remark 2.1.4. Note that from the axiom of separation it follows directly that $\Gamma_v^{*}/$ is a set in the sense of the set theory $\overline{ZFC}_2^{Hs}$.

Definition 2.1.11.We define now the countable set $\mathfrak{I}_2^{*Hs}$ by formula

$$\forall Y\left\{Y \in \mathfrak{I}_2^{*Hs} \Leftrightarrow \exists y\left[([y] \in \Gamma_v^{*Hs}/\sim_v) \wedge \left(g_{\overline{ZFC}_2^{Hs}}(X) = v\right) \wedge Y = X\right]\right\}. \tag{2.1.17}$$

Note that from the axiom schema of replacement (1.1.1) it follows directly that $\mathfrak{I}_2^{*Hs}$ is a

set in the sense of the set theory $\overline{ZFC}_2^{Hs}$.

Definition 2.1.12.We define now the countable set $\mathfrak{R}_2^{*Hs}$ by formula

$$\forall X(X \in \mathfrak{I}_2^{*Hs})[X \in \mathfrak{R}_2^{*Hs} \Leftrightarrow X \notin X]. \tag{2.1.18}$$

Note that from the axiom schema of separation it follows directly that $\mathfrak{R}_2^{*Hs}$ is a set in the

sense of the set theory $\overline{ZFC}_2^{Hs}$.

Remark 2.1.5.Note that $\mathfrak{R}_2^{*Hs} \in \mathfrak{I}_2^{*Hs}$ since $\mathfrak{R}_2^{*Hs}$ is a definable by the following formula

$$\Psi^*(Z) \triangleq \forall X(X \in \mathfrak{I}_2^{*Hs})[X \in Z \Leftrightarrow X \notin X]. \tag{2.1.19}$$

Theorem 2.1.1.Set theory $\overline{ZFC}_2^{Hs}$ is inconsistent.

Proof. From (2.1.18) and Remark 2.1.5 we obtain $\mathfrak{R}_2^{*Hs} \in \mathfrak{R}_2^{*Hs} \Leftrightarrow \mathfrak{R}_2^{*Hs} \notin \mathfrak{R}_2^{*Hs}$ from which immediately one obtains a contradiction $(\mathfrak{R}_2^{*Hs} \in \mathfrak{R}_2^{*Hs}) \wedge (\mathfrak{R}_2^{*Hs} \notin \mathfrak{R}_2^{*Hs})$.

2.2.Derivation of the inconsistent definable set in set

theory ZFC_{st}.

Designation 2.2.1.(i) Let $g_{ZFC_{st}}(u)$ be a Gödel number of given an expression u of the set theory $ZFC_{st} \triangleq ZFC + \exists M_{st}^{ZFC}$.

(ii) Let $\mathbf{Fr}_{st}(y,v)$ be the relation : y is the Gödel number of a wff of the set theoryZFC_{st} that contains free occurrences of the variable X with Gödel number v [9].

(iii) Note that the relation $\mathbf{Fr}_{st}(y,v)$ is expressible in ZFC_{st} by a wff $\widehat{\mathbf{Fr}}_{st}(y,v)$

(iv) Note that for any $y,v \in \mathbb{N}$ by definition of the relation $\mathbf{Fr}_{st}(y,v)$ follows that

$$\widehat{\mathbf{Fr}}_{st}(y,v) \Leftrightarrow \exists!\Psi(X)[(g_{ZFC_{st}}(\Psi(X)) = y) \wedge (g_{ZFC_{st}}(X) = v)], \qquad (2.2.1)$$

where $\Psi(X)$ is a unique wff of ZFC_{st} which contains free occurrences of the variable X with Gödel number v. We denote a unique wff $\Psi(X)$ defined by using equivalence (2.2.1) by symbol $\Psi_{y,v}(X)$, i.e.

$$\widehat{\mathbf{Fr}}_{st}(y,v) \Leftrightarrow \exists!\Psi_{y,v}(X)[(g_{ZFC_{st}}(\Psi_{y,v}(X)) = y) \wedge (g_{ZFC_{st}}(X) = v)], \qquad (2.2.2)$$

(v) Let $\wp_{st}(y,v,v_1)$ be a Gödel number of the following wff: $\exists!X[\Psi(X) \wedge Y = X]$, where $g_{ZFC_{st}}(\Psi(X)) = y, g_{ZFC_{st}}(X) = v,\ g_{ZFC_{st}}(Y) = v_1$.

(vi) Let $\mathrm{Pr}_{ZFC_{st}}(z)$ be a predicate asserting provability in ZFC_{st}, which defined by formula (2.6) in section 2, see Remark 2.2 and Designation 2.3,(see also [8]-[9]).

Definition 2.2.1. Let Γ_X^{st} be the countable collection of the all 1-place open wff's of the set theory ZFC_{st} that contains free occurrences of the variable X.

Definition 2.2.2. Let $g_{ZFC_{st}}(X) = v$. Let Γ_v^{st} be a set of the all Gödel numbers of the 1-place open wff's of the set theory ZFC_{st} that contains free occurrences of the variable X with Gödel number v, i.e.

$$\Gamma_v^{st} = \{y \in \mathbb{N} | \langle y,v \rangle \in \mathbf{Fr}_{st}(y,v)\}, \qquad (2.2.3)$$

or in the following equivalent form:

$$\forall y(y \in \mathbb{N})\Big[y \in \Gamma_v^{st} \Leftrightarrow (y \in \mathbb{N}) \wedge \widehat{\mathbf{Fr}}_{st}(y,v)\Big].$$

Remark 2.2.1.Note that from the axiom of separation it follows directly that Γ_v^{st} is a set in the sense of the set theory ZFC_{st}.

Definition 2.2.3.(i)We define now the equivalence relation $(\bullet \sim_X \bullet) \subset \Gamma_X^{st} \times \Gamma_X^{st}$ by

$$\Psi_1(X) \sim_X \Psi_2(X) \Leftrightarrow (\forall X[\Psi_1(X) \Leftrightarrow \Psi_2(X)]) \qquad (2.2.4)$$

(ii) A subcollection Λ_X^{st} of Γ_X^{st} such that $\Psi_1(X) \sim_X \Psi_2(X)$ holds for all $\Psi_1(X)$ and $\Psi_2(X)$ in

Λ_X^{st}, and never for $\Psi_1(X)$ in Λ_X^{st} and $\Psi_2(X)$ outside Λ_X^{st}, is an equivalence class of Γ_X^{st}.

(iii) For any $\Psi(X) \in \Gamma_X^{st}$ let $[\Psi(X)]_{st} \triangleq \{\Phi(X) \in \Gamma_X^{st} | \Psi(X) \sim_X \Phi(X)\}$ denote the equivalence class to which $\Psi(X)$ belongs. All elements of Γ_X^{st} equivalent to each other are also elements of the same equivalence class.

(iv) The collection of all possible equivalence classes of Γ_X^{st} by $\sim_X$, denoted $\Gamma_X^{st}/\sim_X$

$$\Gamma_X^{st}/\sim_X \triangleq \{[\Psi(X)]_{st} | \Psi(X) \in \Gamma_X^{st}\}. \tag{2.2.5}$$

Definition 2.2.4.(i)We define now the equivalence relation $(\bullet \sim_v \bullet) \subset \Gamma_v^{st} \times \Gamma_v^{st}$ in the sense of the set theory ZFC_{st} by

$$y_1 \sim_v y_2 \Leftrightarrow (\forall X[\Psi_{y_1,v}(X) \Leftrightarrow \Psi_{y_2,v}(X)]) \tag{2.2.6}$$

Note that from the axiom of separation it follows directly that the equivalence relation $(\bullet \sim_v \bullet)$ is a relation in the sense of the set theory ZFC_{st}.

(ii) A subset Λ_v^{st} of Γ_v^{st} such that $y_1 \sim_v y_2$ holds for all y_1 and y_1 in Λ_v^{st},and never for y_1 in Λ_v^{st} and y_2 outside Λ_v^{st}, is an equivalence class of Γ_v^{st}.

(iii) For any $y \in \Gamma_v^{st}$ let $[y]_{st} \triangleq \{z \in \Gamma_v^{st} | y \sim_v z\}$ denote the equivalence class to which y belongs. All elements of Γ_v^{st} equivalent to each other are also elements of the same equivalence class.

(iv)The collection of all possible equivalence classes of Γ_v^{st} by $\sim_v$, denoted $\Gamma_v^{st}/\sim_v$

$$\Gamma_v^{st}/\sim_v \triangleq \{[y]_{st} | y \in \Gamma_v^{st}\}. \tag{2.2.7}$$

Remark 2.2.2. Note that from the axiom of separation it follows directly that $\Gamma_v^{st}/\sim_v$ is a set in the sense of the set theory ZFC_{st}.

Definition 2.2.5.Let $\mathfrak{I}_{st}$ be the countable collection of the all sets definable by 1-place open wff of the set theory ZFC_{st}, i.e.

$$\forall Y\{Y \in \mathfrak{I}_{st} \Leftrightarrow \exists \Psi(X)[([\Psi(X)]_{st} \in \Gamma_X^{st}/\sim_X) \wedge [\exists! X[\Psi(X) \wedge Y = X]]]\}. \tag{2.2.8}$$

Definition 2.2.6.We rewrite now (2.2.8) in the following equivalent form

$$\forall Y\{Y \in \mathfrak{I}_{st} \Leftrightarrow \exists \Psi(X)[([\Psi(X)]_{st} \in \Gamma_X^{*st}/\sim_X) \wedge (Y = X)]\}, \tag{2.2.9}$$

where the countable collection $\Gamma_X^{*st}/\sim_X$ is defined by

$$\forall \Psi(X)\{[\Psi(X)]_{st} \in \Gamma_X^{*st}/\sim_X \Leftrightarrow [([\Psi(X)]_{st} \in \Gamma_X^{st}/\sim_X) \wedge \exists! X\Psi(X)]\} \tag{2.2.10}$$

Definition 2.2.7. Let $\mathfrak{R}_{st}$ be the countable collection of the all sets such that

$$\forall X(X \in \mathfrak{I}_{st})[X \in \mathfrak{R}_{st} \Leftrightarrow X \notin X]. \tag{2.2.11}$$

Remark 2.2.3. Note that $\mathfrak{R}_{st} \in \mathfrak{I}_{st}$ since $\mathfrak{R}_{st}$ is a collection definable by 1-place open wff

$$\Psi(Z,\mathfrak{I}_{st}) \triangleq \forall X(X \in \mathfrak{I}_{st})[X \in Z \Leftrightarrow X \notin X].$$

From (2.2.11) and Remark 2.2.3 one obtains directly

$$\mathfrak{R}_{st} \in \mathfrak{R}_{st} \Leftrightarrow \mathfrak{R}_{st} \notin \mathfrak{R}_{st}. \tag{2.2.12}$$

But (2.2.12) immediately gives a contradiction

$$(\mathfrak{R}_{st} \in \mathfrak{R}_{st}) \wedge (\mathfrak{R}_{st} \notin \mathfrak{R}_{st}). \tag{2.2.13}$$

However contradiction (2.2.13) it is not a true contradiction inside ZFC_{st} for the reason

that the countable collection $\mathfrak{I}_{st}$ is not a set in the sense of the set theory ZFC_{st}.

In order to obtain a true contradiction inside ZFC_{st} we introduce the following definitions.

Definition 2.2.8.We define now the countable set $\Gamma_v^{*st}/\sim_v$ by formula

$$\forall y\left\{[y]_{st} \in \Gamma_v^{*st}/\sim_v \Leftrightarrow ([y]_{st} \in \Gamma_v^{st}/\sim_v) \wedge \widehat{\mathbf{Fr}}_{st}(y,v) \wedge [\exists! X \Psi_{y,v}(X)]\right\}. \tag{2.2.14}$$

Remark 2.2.4. Note that from the axiom of separation it follows directly that $\Gamma_v^{*st}/\sim_v$ is a

set in the sense of the set theory ZFC_{st}.

Definition 2.2.9.We define now the countable set $\mathfrak{I}_{st}^*$ by formula

$$\forall Y\{Y \in \mathfrak{I}_{st}^* \Leftrightarrow \exists y[([y]_{st} \in \Gamma_v^{*st}/\sim_v) \wedge (g_{ZFC_{st}}(X) = v) \wedge Y = X]\}. \tag{2.2.15}$$

Note that from the axiom schema of replacement it follows directly that $\mathfrak{I}_{st}^*$ is a set in the

sense of the set theory ZFC_{st}.

Definition 2.2.10.We define now the countable set $\mathfrak{R}_{st}^*$ by formula

$$\forall X(X \in \mathfrak{I}_{st}^*)[X \in \mathfrak{R}_{st}^* \Leftrightarrow X \notin X]. \tag{2.2.16}$$

Note that from the axiom schema of separation it follows directly that $\mathfrak{R}_{st}^*$ is a set in the

sense of the set theory ZFC_{st}.

Remark 2.2.5.Note that $\mathfrak{R}_{st}^* \in \mathfrak{I}_{st}^*$ since $\mathfrak{R}_{st}^*$ is a definable by the following formula

$$\Psi^*(Z) \triangleq \forall X(X \in \mathfrak{I}_{st}^*)[X \in Z \Leftrightarrow X \notin X]. \tag{2.2.17}$$

Theorem 2.2.1.Set theory ZFC_{st} is inconsistent.

Proof. From (2.2.17) and Remark 2.2.5 we obtain $\mathfrak{R}_{st}^* \in \mathfrak{R}_{st}^* \Leftrightarrow \mathfrak{R}_{st}^* \notin \mathfrak{R}_{st}^*$ from which

immediately one obtains a contradiction $(\mathfrak{R}_{st}^* \in \mathfrak{R}_{st}^*) \wedge (\mathfrak{R}_{st}^* \notin \mathfrak{R}_{st}^*)$.

2.3.Derivation of the inconsistent definable set in ZFC_{Nst}.

Definition 2.3.1.Let $\overline{PA}$ be a first order theory which contain usual postulates of Peano

arithmetic [9] and recursive defining equations for every primitive recursive function as desired.So for any $(n+1)$-place function f defined by primitive recursion over any n-place base function g and $(n+2)$-place iteration function h there would be the defining equations:

(i) $f(0,y_1,\ldots,y_n) = g(y_1,\ldots,y_n)$, (ii) $f(x+1,y_1,\ldots,y_n) = h(x,f(x,y_1,\ldots,y_n),y_1,\ldots,y_n)$.

Designation 2.3.1.(i) Let M_{Nst}^{ZFC} be a nonstandard model of ZFC and let $M_{st}^{\overline{PA}}$ be a standard model of $\overline{PA}$. We assume now that $M_{st}^{\overline{PA}} \subset M_{Nst}^{ZFC}$ and denote such nonstandard model of the set theory ZFC by $M_{Nst}^{ZFC}[\overline{PA}]$. (ii) Let ZFC_{Nst} be the theory

$$ZFC_{Nst} = ZFC + M_{Nst}^{ZFC}[\overline{PA}].$$

Designation 2.3.2.(i) Let $g_{ZFC_{Nst}}(u)$ be a Gödel number of given an expression u of the set theory $ZFC_{Nst} \triangleq ZFC + \exists M_{Nst}^{ZFC}[\overline{PA}]$.

(ii) Let $\mathbf{Fr}_{Nst}(y,v)$ be the relation : y is the Gödel number of a wff of the set theory ZFC_{Nst} that contains free occurrences of the variable X with Gödel number v [9].

(iii) Note that the relation $\mathbf{Fr}_{Nst}(y,v)$ is expressible in ZFC_{Nst} by a wff $\widehat{\mathbf{Fr}}_{Nst}(y,v)$

(iv) Note that for any $y,v \in \mathbb{N}$ by definition of the relation $\mathbf{Fr}_{Nst}(y,v)$ follows that

$$\widehat{\mathbf{Fr}}_{Nst}(y,v) \Leftrightarrow \exists!\Psi(X)[(g_{ZFC_{Nst}}(\Psi(X)) = y) \wedge (g_{ZFC_{Nst}}(X) = v)], \qquad (2.3.1)$$

where $\Psi(X)$ is a unique wff of ZFC_{st} which contains free occurrences of the variable X with Gödel number v. We denote a unique wff $\Psi(X)$ defined by using equivalence (2.3.1) by symbol $\Psi_{y,v}(X)$, i.e.

$$\widehat{\mathbf{Fr}}_{Nst}(y,v) \Leftrightarrow \exists!\Psi_{y,v}(X)[(g_{ZFC_{Nst}}(\Psi_{y,v}(X)) = y) \wedge (g_{ZFC_{Nst}}(X) = v)], \qquad (2.3.2)$$

(v) Let $\wp_{Nst}(y,v,v_1)$ be a Gödel number of the following wff: $\exists!X[\Psi(X) \wedge Y = X]$, where

$g_{ZFC_{Nst}}(\Psi(X)) = y, g_{ZFC_{Nst}}(X) = v,\ g_{ZFC_{Nst}}(Y) = v_1$.

(vi) Let $\mathrm{Pr}_{ZFC_{Nst}}(z)$ be a predicate asserting provability in ZFC_{Nst}, which defined by formula (2.6) in section 2, see Remark 2.2 and Designation 2.3,(see also [9]-[10]).

Definition 2.3.2. Let Γ_X^{Nst} be the countable collection of the all 1-place open wff's of the set theory ZFC_{Nst} that contains free occurrences of the variable X.

Definition 2.3.3. Let $g_{ZFC_{Nst}}(X) = v$. Let Γ_v^{Nst} be a set of the all Gödel numbers of the 1-place open wff's of the set theory ZFC_{Nst} that contains free occurrences of the variable X with Gödel number v, i.e.

$$\Gamma_v^{Nst} = \{y \in \mathbb{N} | \langle y, v \rangle \in \mathbf{Fr}_{Nst}(y, v)\}, \tag{2.3.3}$$

or in the following equivalent form:

$$\forall y (y \in \mathbb{N}) \Big[y \in \Gamma_v^{Nst} \Leftrightarrow (y \in \mathbb{N}) \wedge \widehat{\mathbf{Fr}}_{Nst}(y, v) \Big].$$

Remark 2.3.1.Note that from the axiom of separation it follows directly that Γ_v^{st} is a set
in the sense of the set theory ZFC_{Nst}.

Definition 2.3.3.(i)We define now the equivalence relation $(\bullet \sim_X \bullet) \subset \Gamma_X^{Nst} \times \Gamma_X^{Nst}$ by

$$\Psi_1(X) \sim_X \Psi_2(X) \Leftrightarrow (\forall X [\Psi_1(X) \Leftrightarrow \Psi_2(X)]) \tag{2.3.4}$$

(ii) A subcollection Λ_X^{st} of Γ_X^{st} such that $\Psi_1(X) \sim_X \Psi_2(X)$ holds for all $\Psi_1(X)$ and $\Psi_2(X)$ in
Λ_X^{st}, and never for $\Psi_1(X)$ in Λ_X^{Nst} and $\Psi_2(X)$ outside Λ_X^{Nst}, is an equivalence class of Γ_X^{Nst}.

(iii) For any $\Psi(X) \in \Gamma_X^{Nst}$ let $[\Psi(X)]_{Nst} \triangleq \{\Phi(X) \in \Gamma_X^{Nst} | \Psi(X) \sim_X \Phi(X)\}$ denote the equivalence class to which $\Psi(X)$ belongs. All elements of Γ_X^{st} equivalent to each other
are also elements of the same equivalence class.

(iv) The collection of all possible equivalence classes of Γ_X^{Nst} by $\sim_X$, denoted $\Gamma_X^{Nst}/\sim_X$

$$\Gamma_X^{Nst}/\sim_X \triangleq \{[\Psi(X)]_{Nst} | \Psi(X) \in \Gamma_X^{Nst}\}. \tag{2.3.5}$$

Definition 2.3.4.(i)We define now the equivalence relation $(\bullet \sim_v \bullet) \subset \Gamma_v^{Nst} \times \Gamma_v^{Nst}$ in the
sense of the set theory ZFC_{Nst} by

$$y_1 \sim_v y_2 \Leftrightarrow (\forall X [\Psi_{y_1,v}(X) \Leftrightarrow \Psi_{y_2,v}(X)]) \tag{2.3.6}$$

Note that from the axiom of separation it follows directly that the equivalence relation
$(\bullet \sim_v \bullet)$ is a relation in the sense of the set theory ZFC_{Nst}.

(ii) A subset Λ_v^{Nst} of Γ_v^{Nst} such that $y_1 \sim_v y_2$ holds for all y_1 and y_1 in Λ_v^{Nst},and never for y_1 in
Λ_v^{Nst} and y_2 outside Λ_v^{Nst}, is an equivalence class of Γ_v^{Nst}.

(iii) For any $y \in \Gamma_v^{Nst}$ let $[y]_{Nst} \triangleq \{z \in \Gamma_v^{Nst} | y \sim_v z\}$ denote the equivalence class to which y
belongs. All elements of Γ_v^{Nst} equivalent to each other are also elements of the same equivalence class.

(iv)The collection of all possible equivalence classes of Γ_v^{Nst} by $\sim_v$, denoted $\Gamma_v^{Nst}/\sim_v$

$$\Gamma_v^{Nst}/\sim_v \triangleq \{[y]_{Nst} | y \in \Gamma_v^{Nst}\}. \tag{2.3.7}$$

Remark 2.3.2. Note that from the axiom of separation it follows directly that $\Gamma_v^{Nst}/\sim_v$ is a
set in the sense of the set theory ZFC_{Nst}.

Definition 2.3.5.Let $\Im_{Nst}$ be the countable collection of the all sets definable by

1-place
open wff of the set theory ZFC_{Nst}, i.e.

$$\forall Y\{Y \in \mathfrak{I}_{Nst} \Leftrightarrow \exists \Psi(X)[([\Psi(X)]_{Nst} \in \Gamma_X^{Nst}/\sim_X\ ) \wedge [\exists! X[\Psi(X) \wedge Y = X]]]\}. \quad (2.3.8)$$

Definition 2.3.6.We rewrite now (2.3.8) in the following equivalent form

$$\forall Y\{Y \in \mathfrak{I}_{Nst} \Leftrightarrow \exists \Psi(X)[([\Psi(X)]_{Nst} \in \Gamma_X^{*Nst}/\sim_X\ ) \wedge (Y = X)]\}, \quad (2.3.9)$$

where the countable collection $\Gamma_X^{*Nst}/\sim_X$ is defined by

$$\forall \Psi(X)\{[\Psi(X)]_{Nst} \in \Gamma_X^{*Nst}/\sim_X \Leftrightarrow [([\Psi(X)]_{Nst} \in \Gamma_X^{Nst}/\sim_X\ ) \wedge \exists! X\Psi(X)]\} \quad (2.3.10)$$

Definition 2.3.7. Let $\mathfrak{R}_{Nst}$ be the countable collection of the all sets such that

$$\forall X(X \in \mathfrak{I}_{Nst})[X \in \mathfrak{R}_{Nst} \Leftrightarrow X \notin X]. \quad (2.3.11)$$

Remark 2.3.3.Note that $\mathfrak{R}_{Nst} \in \mathfrak{I}_{Nst}$ since $\mathfrak{R}_{Nst}$ is a collection definable by 1-place open
wff

$$\Psi(Z, \mathfrak{I}_{Nst}) \triangleq \forall X(X \in \mathfrak{I}_{Nst})[X \in Z \Leftrightarrow X \notin X].$$

From (2.3.11) one obtains

$$\mathfrak{R}_{Nst} \in \mathfrak{R}_{Nst} \Leftrightarrow \mathfrak{R}_{Nst} \notin \mathfrak{R}_{Nst}. \quad (2.3.12)$$

But (2.3.12) gives a contradiction

$$(\mathfrak{R}_{Nst} \in \mathfrak{R}_{Nst}) \wedge (\mathfrak{R}_{Nst} \notin \mathfrak{R}_{Nst}). \quad (2.3.13)$$

However a contradiction (2.3.13) it is not a true contradiction inside ZFC_{Nst} for the reason
that the countable collection $\mathfrak{I}_{Nst}$ is not a set in the sense of the set theory ZFC_{Nst}.
In order to obtain a true contradiction inside ZFC_{Nst} we introduce the following definitions.

Definition 2.3.8.We define now the countable set $\Gamma_v^{*Nst}/\sim_v$ by formula

$$\forall y\left\{[y]_{Nst} \in \Gamma_v^{*Nst}/\sim_v \Leftrightarrow ([y]_{Nst} \in \Gamma_v^{Nst}/\sim_v\ ) \wedge \widehat{\mathbf{Fr}}_{Nst}(y,v) \wedge [\exists! X\Psi_{y,v}(X)]\right\}. \quad (2.3.14)$$

Remark 2.3.4. Note that from the axiom of separation it follows directly that $\Gamma_v^{*Nst}/\sim_v$ is
a set in the sense of the set theory ZFC_{st}.
Definition 2.3.9.We define now the countable set $\mathfrak{I}_{Nst}^*$ by formula

$$\forall Y\{Y \in \mathfrak{I}_{Nst}^* \Leftrightarrow \exists y[([y]_{Nst} \in \Gamma_v^{*Nst}/\sim_v\ ) \wedge (g_{ZFC_{Nst}}(X) = v) \wedge Y = X]\}. \quad (2.3.15)$$

Note that from the axiom schema of replacement it follows directly that $\mathfrak{I}_{st}^*$ is a set in the
sense of the set theory ZFC_{Nst}.
Definition 2.3.10.We define now the countable set $\mathfrak{R}_{Nst}^*$ by formula

$$\forall X(X \in \mathfrak{I}^*_{Nst})[X \in \mathfrak{R}^*_{Nst} \Leftrightarrow X \notin X]. \tag{2.3.16}$$

Note that from the axiom schema of separation it follows directly that $\mathfrak{R}^*_{Nst}$ is a set in the

sense of the set theory ZFC_{Nst}.

Remark 2.3.5.Note that $\mathfrak{R}^*_{Nst} \in \mathfrak{I}^*_{Nst}$ since $\mathfrak{R}^*_{Nst}$ is a definable by the following formula

$$\Psi^*(Z) \triangleq \forall X(X \in \mathfrak{I}^*_{Nst})[X \in Z \Leftrightarrow X \notin X]. \tag{2.3.17}$$

Theorem 2.3.1.Set theory ZFC_{Nst} is inconsistent.

Proof. From (2.3.16) and Remark 2.3.5 we obtain $\mathfrak{R}^*_{Nst} \in \mathfrak{R}^*_{Nst} \Leftrightarrow \mathfrak{R}^*_{Nst} \notin \mathfrak{R}^*_{Nst}$ from which one obtains a contradiction $(\mathfrak{R}^*_{Nst} \in \mathfrak{R}^*_{Nst}) \wedge (\mathfrak{R}^*_{Nst} \notin \mathfrak{R}^*_{Nst})$.

References.

[1] E. Nelson.Warning Signs of a Possible Collapse of Contemporary Mathematics. https://web.math.princeton.edu/~nelson/papers/warn.pdf In Infinity: New Research Frontiers, by Michael Heller (Editor), W. Hugh Woodin Pages 75–85, 2011. Published February 14th 2013 by Cambridge University Press Hardcover, 311 pages. ISBN: 1107003873 (ISBN13: 9781107003873)

[2] L. Henkin, "Completeness in the theory of types". Journal of Symbolic Logic 15 (2): 81–91. doi:10.2307/2266967. JSTOR 2266967

[3] M. Rossberg, "First-Order Logic, Second-Order Logic, and Completeness". In V.Hendricks et al., eds. First-order logic revisited. Berlin: Logos-Verlag.

[4] S. Shapiro, Foundations without Foundationalism: A Case for Second-order Logic. Oxford University Press. ISBN 0-19-825029-0

[5] A. Rayo and G. Uzquiano,Toward a Theory of Second-Order Consequence, Notre Dame Journal of Formal Logic Volume 40, Number 3, Summer 1999.

[6] J. Vaananen, Second-Order Logic and Foundations of Mathematics, The Bulletin of Symbolic Logic, Vol.7, No. 4 (Dec., 2001), pp. 504-520.

[7] G. Uzquiano, Quantification without a domain. New Waves in Philosophy of Mathematics. Springer, 29 Sep 2009 - Philosophy - 327 pp. ISBN 0230245196, 9780230245198

[8] P. Cohen, Set Theory and the continuum hypothesis.Reprint of the W. A. Benjamin,Inc.,New York,1966 edition. ISBN-13: 978-0486469218

[9] E. Mendelson,Introduction to mathematical logic.1997. ISBN-10: 0412808307. ISBN-13: 978-0412808302

[10] 2016 EUROPEAN SUMMER MEETING OF THE ASSOCIATION FOR SYMBOLIC LOGIC, LOGIC COLLOQUIUM ’16, Leeds, UK, July 31-August 6, 2016,

The Bulletin of Symbolic Logic Vol. 23, No. 2 (JUNE 2017), pp. 213-266
J.Foukzon, Inconsistent countable set in second order ZFC and unexistence of the strongly inaccessible cardinals, pp.240.

[11] J. Foukzon, Inconsistent Countable Set in Second Order ZFC and Nonexistence of the Strongly Inaccessible Cardinals,British Journal of Mathematics & Computer Science, ISSN: 2231-0851,Vol.: 9, Issue.: 5
http://www.sciencedomain.org/abstract/9622

[12] J. Foukzon,Generalized Lob's Theorem.Strong Reflection Principles and Large Cardinal Axioms. Consistency Results in Topology.
http://arxiv.org/abs/1301.5340v10

[13] J.Foukzon,E. R. Men'kova,Generalized Löb's Theorem. Strong Reflection Principles and Large Cardinal Axioms, Advances in Pure Mathematics, Vol.3 No.3, 2013.
http://dx.doi.org/10.4236/apm.2013.33053