

Recursively-Defined Combinatorial Functions: The Case of Binomial and Multinomial Coefficients and Probabilities

Abstract

This paper studies a prominent class of recursively-defined combinatorial functions, namely, the binomial and multinomial coefficients and probabilities. The paper reviews the basic notions and mathematical definitions of these four functions. Subsequently, it characterizes each of these functions *via* a recursive relation that is valid over a certain two-dimensional or multi-dimensional region and is supplemented with certain boundary conditions. Visual interpretations of these characterizations are given in terms of regular acyclic signal flow graphs. The graph for the binomial coefficients resembles a Pascal Triangle, while that for trinomial or multinomial coefficients looks like a Pascal Pyramid, Tetrahedron, or Hyper-Pyramid. Each of the four functions is computed using both its conventional and recursive definitions. Moreover, the recursive structures of the binomial coefficient and probability are utilized in an iterative evaluation of each of them, which is substantially more efficient than the conventional or recursive evaluation. Analogous iterative evaluations of the multinomial coefficient and probability can be constructed similarly. Applications to the reliability evaluation for two-valued and multi-valued k-out-of-n systems are also pointed out.

Keywords: Binomial, multinomial, recursion, boundary condition, signal flow graph, iteration, k-out-of-n.

1. Introduction

Many combinatorial functions can be characterized by a general framework based on simple two-dimensional or multi-dimensional recursion [1-19]. Four prominent cases among these combinatorial functions are the functions of binomial coefficients [6, 10, 15, 20-22] and binomial probabilities [1, 6, 10, 23,

24], as well as their extensions to multinomial coefficients [21, 22, 25-32] and multinomial probabilities [21, 22, 33, 34].

This paper reviews the basic notions and concepts of the aforementioned four combinatorial functions. It points out that computing any of them (*via* conventional methods involving factorials) has exponential temporal complexity. The paper explores the recursive structure for each of these functions with an eye of constructing *more efficient iterative* computational methods, which could possibly be of a polynomial (sub-exponential) rather than exponential temporal complexity.

We follow the technique introduced by Rushdi [1, 6, 10] of characterizing a recursively-defined function *via* three entities, *viz.*, (a) a recursive relation, (b) a region of validity for the recurrence, and (c) boundary conditions. We provide pictorial insight for these entities through the use of (Mason) linear signal flow graphs (SFGs) [35-37], which turn out to be regular and acyclic or loopless. Linear signal flow graphs are very useful in representing and manipulating linear relations in a wide variety of scientific and engineering applications [6, 10, 37-42]. The SFG for the binomial coefficient is simply a Pascal Triangle (also known as al-Karkhi Triangle or al-Khayyam Triangle) [21, 22, 43-48], while the SFG for the multinomial coefficients is a Pascal Pyramid or a Pascal Hyper-pyramid [49-56].

The topics of binomial coefficients and binomial probabilities, and their associated topics of k-out-of-n system reliability and unreliability have been extensively studied before [1, 6, 10, 15, 20-24]. They are included herein to make the paper self-contained and to facilitate the extension to the topics of multinomial coefficients and multinomial probabilities, and subsequently to the associated topics of multi-valued k-out-of-n system reliability and unreliability.

The organization of the remainder of this paper is as follows. Section 2 introduces binomial coefficients and probabilities and shows that their recurrences are represented by signal flow graphs that resemble Pascal Triangles. Section 3 exposes the concept and conventional computational methods for the multinomial coefficients and probabilities. Subsequently, Sections 4 and 5 introduce recurrences for multinomial coefficients and multinomial probabilities, respectively, and show how such recurrences are represented *via* Pascal Pyramids or Hyper-pyramids. Section 6 makes a quick comparison of iteration versus recursion, with a stress on the utility of both

techniques in the computation of the four functions considered herein. Section 7 concludes the paper.

2. Binomial Coefficients and Binomial Probabilities

In a (conventional) Bernoulli trial, one of two distinct outcomes might result. These two outcomes are typically called success and failure, and their probabilities are assumed to be constant, and denoted by p and q , respectively. The number of k successes in n independent trials is called the binomial (combinatorial) coefficient, and is given by [21-24]

$$B(k, n) = n! / (k! (n - k)!) = \prod_{m=1}^{\min(k, n-k)} ((n - m + 1)/m) \quad (1)$$

The binomial coefficient $B(k, n)$ also expresses the number of ways of selecting k objects out of n objects, when repetition (replacement) is not allowed and order does not matter. In particular, $B(k, n)$ denotes the number of subsets with k elements among the subsets of a set on n elements. The binomial coefficient satisfies the recursive relation [6, 15, 21, 22].

$$B(k, n) = B(k, n - 1) + B(k - 1, n - 1), \quad 0 < k < n \quad (2)$$

subject to the boundary conditions

$$B(k, n) = 1 \quad \text{for } (k = 0 \text{ or } k = n) \quad \text{and } n \geq 0 \quad (3)$$

Figure 1 is a Signal Flow Graph (SFG) representing the computation of $B(k, n)$ via (2) and (3). To facilitate comparison with the multinomial case, we use a Cartesian grid (k_1, k_2) where $(k_1 \geq 0, k_2 \geq 0)$ to represent $B(k, n)$ as $B(k_1, k_1 + k_2)$ where $k_1 = k$ and $k_2 = n - k$. Figure 1 is traditionally referred to as Pascal Triangle (albeit it is also known by many other names such as al-Karkhi Triangle or al-Khayyam Triangle [57, 58]). This triangle occupies an octant of the $k - n$ plane or the $k_1 - k_2$ plane. It extends without bound as n goes to infinity, but it is otherwise perfectly bounded by the two straight arrows given in (3), viz., $(k = 0, n \geq 0)$ and $(k = n, n \geq 0)$.

For convenience, and to maintain symmetry, equations (1)-(3) are rewritten as:

$$B(k_1, k_1 + k_2) = (k_1 + k_2)! / (k_1! k_2!) \quad (1a)$$

$$\begin{aligned} B(k_1, k_1 + k_2) &= B(k_1, k_1 + k_2 - 1) + B(k_1 - 1, k_1 + k_2 - 1), \quad k_1 > 0, k_2 > 0 \\ &= B((k_1, k_1 + k_2) | k_2 \leftarrow k_2 - 1) + B((k_1, k_1 + k_2) | k_1 \leftarrow k_1 - 1), \\ &\quad k_1 > 0, k_2 > 0 \end{aligned} \quad (2a)$$

$$B(k_1, k_1 + k_2) = 1 \quad \text{for} \quad (k_1 = 0, k_2 \geq 0) \text{ or } (k_2 = 0, k_1 \geq 0) \quad (2b)$$

Note that equations (2a) & (3a) exhibit symmetric behavior towards the two dimensions set by k_1 and k_2 . This symmetric behavior can be easily maintained when these two dimensions are replaced by r dimensions ($r > 2$) in the study of multinomial coefficients.

The probability of obtaining k successes in n trials is the probability mass function (pmf) of the binomial distribution [23, 24], and is given by

$$E(k, n, p) = B(k, n) p^k (1 - p)^{n-k}, \quad 0 \leq k \leq n \quad (4)$$

And the corresponding Cumulative Distribution Function (CDF), and Complementary Cumulative Distribution Function (CCDF) are

$$U(k, n, p) = \sum_{m=0}^{k-1} E(m, n, p) \quad (5)$$

$$R(k, n, p) = \sum_{m=k}^n E(m, n, p) \quad (6)$$

The aforementioned $U(k, n, p)$ and $R(k, n, p)$ represent, respectively, the unreliability and reliability of a (two-valued) k -out-of- n : G system, which is a coherent system that is good if and only if *at least* k out of its n components are good [1, 6, 10].

Let us now generalize our notion of a Bernoulli trial by relaxing the condition that the probabilities p and q of trial success and failure be constant (equivalently, assuming that component reliabilities are not necessarily equal). This generalization does not affect $B(k, n)$, but leads us to replace $E(k, n, p)$ by $E(k, n, \mathbf{p})$, where $\mathbf{p} = [p_1 \ p_1 \ p_2 \ \dots \dots \dots p_n]^T$. Likewise, we replace $U(k, n, p)$ and $R(k, n, p)$ by $U(k, n, \mathbf{p})$ and $R(k, n, \mathbf{p})$. Note that equations (5) and (6) continue to hold with p replaced by $\mathbf{p}$, i.e.,

$$U(k, n, \mathbf{p}) = \sum_{m=0}^{k-1} E(m, n, \mathbf{p}) \quad (5a)$$

$$R(k, n, \mathbf{p}) = \sum_{m=k}^n E(m, n, \mathbf{p}) \quad (6a)$$

The probability $E(k, n, \mathbf{p})$ satisfies the recursive relation [1-3, 6, 10, 13]

$$E(k, n, \mathbf{p}) = q_n * E(k, n-1, \mathbf{p}/p_n) + p_n * E(k-1, n-1, \mathbf{p}/p_n),$$

$$0 \leq k \leq n, \\ \{k, n\} \neq \{0, 0\} \quad (7)$$

where $\mathbf{p}/p_n = [p_1 \ p_1 \ p_2 \ \dots \dots \dots p_{n-1}]^T$. Equations (7) are subject to the boundary conditions

$$E(k, n, \mathbf{p}) = 1.0 \text{ for } k = n = 0 \quad (8)$$

$$E(k, n, \mathbf{p}) = 0.0 \text{ for } (k = -1 \text{ or } k = n+1) \text{ and } (n \geq 0) \quad (9)$$

Again, to facilitate extension to the multinomial case, we use the grid (k_1, k_2) instead of the grid (k, n) , where $k_1 = k$ and $k_2 = n - k$. We also rewrite p_n as p_{1n} and rewrite q_n as p_{2n} where p_{1n} and p_{2n} are the probabilities of outcome 1 and outcome 2 in the n th trial, respectively. With this change the vectors $\mathbf{p}$ and $\mathbf{p}/p_n$ are replaced by two column matrices of the form

$$\mathbf{p} = \begin{bmatrix} p_{11} & p_{12} & p_{13} & \dots & p_{1(n-1)} & p_{1n} \\ p_{21} & p_{22} & p_{23} & \dots & p_{2(n-1)} & p_{2n} \end{bmatrix}^T \quad (10)$$

$$\mathbf{p}/p_n = \begin{bmatrix} p_{11} & p_{12} & p_{13} & \dots & p_{1(n-1)} \\ p_{21} & p_{22} & p_{23} & \dots & p_{2(n-1)} \end{bmatrix}^T \quad (10)$$

a)

Equations (7) – (9) are now rewritten as

$$E(k_1, k_1 + k_2, \mathbf{p}) = p_{2n} * E(k_1, k_1 + k_2 - 1, \mathbf{p}/\mathbf{p}_n) + p_{1n} * E(k_1 - 1, k_1 + k_2 - 1, \mathbf{p}/\mathbf{p}_n) \\ = p_{2n} * E((k_1, k_1 + k_2, \mathbf{p}/\mathbf{p}_n) | k_2 \leftarrow k_2 - 1) +$$

$$p_{1n} * E((k_1, k_1 + k_2, \mathbf{p}/\mathbf{p}_n)) | k_1 \leftarrow k_1 - 1) \quad (7a)$$

$$E(k_1, k_1 + k_2, \mathbf{p}) = 1.0 \quad \text{for } k_1 = k_2 = 0 \quad (8a)$$

$$E(k_1, k_1 + k_2, \mathbf{p}) = 0.0 \quad \text{for } (k_1 = -1, k_2 \geq 1) \text{ or } (k_1 \geq 1, k_2 = -1) \quad (9a)$$

Again, equations (7a)–(9a) exhibit symmetric behavior towards the two dimensions of k_1 and k_2 , and can keep such a behavior when a vectorial $\mathbf{k}$ of r dimensions ($r > 2$) is involved.

Figure 2 is a Signal Flow Graph (SFG) representing the computation of the binomial probability $E(k, n, \mathbf{p})$. This graph appeared earlier in a (k, n) grid in [1-3, 6, 10, 13, 16] and in a (k_1, k_2) grid in [19]. The SFG in Fig. 2 has a striking similarity to the one in Fig. 1. Both SFGs are acyclic (have no loops), and hence are analyzed through path enumeration and multiplication of transmittances along paths. For both graphs, the Mason gain formula [6, 10, 35-38] can be easily applied with all deltas set to 1.

The SFG in Fig. 1 has constant transmittances of value 1 each, while the one in Fig. 2 has variable transmittances representing outcome probabilities in a certain trial. Both graphs have an infinite number of source nodes along the two Cartesian axes, which serve as a boundary for the region in which the pertinent recursion is valid. However, Fig. 2 has only a single real source of value 1.0 (highlighted in black). It has a doubly infinite number of fictitious sources (each of value 0.0 colored in white). These sources can be omitted, but they are retained herein to maintain symmetry of the recursion and to appropriately depict the limit for its region of validity.

In passing, we note that the binomial recursion introduced in this section is just one kind of binomial recursion, which might be conveniently called Pascal's recursion. This recursion has the distinct advantages of being extendible to the multinomial case and of its use of addition solely (rather than multiplication and division). Other types of binomial recursion (that possess certain merits of their own) also exist [20, 59, 60], but will not be pursued further herein.

3. Multinomial Coefficients and Multinomial Probabilities

Let us generalize the original Bernoulli trials of Section 2 in another direction by allowing the number of outcomes in a single trial to be $r > 2$. We use the vector

$$\mathbf{k} = [k_1 k_2 \dots \dots k_r]^T$$

to denote the numbers of occurrences of the various r outcomes in n independent trials, *i.e.*, k_i is the number of times outcome i ($1 \leq i \leq r$) occurs, so that

$$k_i \geq 0, \quad 1 \leq i \leq r \quad (11a)$$

$$\sum_{i=1}^r k_i = n \quad (11b)$$

The multinomial coefficient $M(\mathbf{k}, n, r)$ is the number of ways in which outcome i ($1 \leq i \leq r$) occurs k_i times in n independent trials. It is given in terms of the factorial of n divided by the product of factorials of the k_i 's, namely [21, 22]

$$\begin{aligned} M(\mathbf{k}, n, r) &= n! / (k_1! k_2! \dots k_r!) \\ &= (k_1 + k_2 + \dots + k_r)! / (k_1! k_2! \dots k_r!) \end{aligned} \quad (12)$$

Note that since $B(\mathbf{k}, n) = M(\mathbf{k}, n, 2)$, equation (12) reduces to (1a) when $r = 2$.

Let us further assume that the probability of occurrence of outcome i throughout the n independent trials is a constant p_i such that

$$p_i \geq 0, \quad 1 \leq i \leq r \quad (13a)$$

$$\sum_{i=1}^r p_i = 1.0 \quad (13b)$$

The multinomial probability (the probability that occurrences of the r outcomes in n independent trials follow the vector $\mathbf{k}$) is denoted by $E(\mathbf{k}, n, r, \mathbf{p})$ and given by

$$E(\mathbf{k}, n, r, \mathbf{p}) = M(\mathbf{k}, n, r) \prod_{i=1}^r p_i^{k_i} \quad (14)$$

We now further combine the two generalizations by allowing $r > 2$ outcomes and allowing variable probabilities for various outcomes in different trials. Let p_{ij} be the probability of occurrence of outcome i in trial j , so that

$$p_{ij} \geq 0, \quad 1 \leq i \leq r, \quad 1 \leq j \leq n \quad (15a)$$

$$\sum_{i=1}^r p_{ij} = 1.0 \quad 1 \leq j \leq n \quad (15b)$$

This change does not affect $M(\mathbf{k}, n, r)$, but it affects $E(\mathbf{k}, n, r, \mathbf{p})$ since $\mathbf{p}$ is no longer a column vector of length r , but is instead a matrix of r rows and n columns. Equation (14) is no longer valid, and is to be replaced by a recursive relation to be presented in Section 5.

4. Recurrence for Multinomial Coefficients

To obtain a recurrence for the multinomial coefficients, we note that the r events

$$V_i = \{\text{outcome } i \text{ occurs in the } n\text{th trial}\}, \quad 1 \leq i \leq r$$

are mutually exclusive and exhaustive. Hence, we can write (when $k_i > 0$ for $1 \leq i \leq r$)

$$M(\mathbf{k}, n, r) = \sum_{i=1}^r (M(\mathbf{k}, n, r) | V_i) = \sum_{i=1}^r M(\mathbf{k} | k_i \leftarrow k_i - 1, n - 1, r) \quad (16)$$

Note that the r -dimensional vector

$$\mathbf{k} = [k_1 \ k_2 \ \dots \ k_{i-1} \ \textcolor{red}{k_i} \ k_{i+1} \ \dots \ k_r]^T \quad (17)$$

is replaced in the RHS of (16) by another vector of the same size given by

$$(\mathbf{k} | k_i \leftarrow k_i - 1) = [k_1 \ k_2 \ \dots \ k_{i-1} \ \textcolor{red}{(k_i - 1)} \ k_{i+1} \ \dots \ k_r]^T \quad (18)$$

The above recurrence occurs subject to the following boundary conditions, which are valid for $n > 0$

$$M(\mathbf{k}, n, r) = 1 \quad \text{when} \quad r = 1 \quad (19a)$$

$$M(\mathbf{k}, n, r) = M(\mathbf{k}/k_i, n, r - 1) \quad \text{when} \quad k_i = 0, \quad 1 \leq i \leq r \quad (19b)$$

The condition in (19a) occurs when one particular $k_i = n$ so that all other k_i 's are 0's. The condition (19b) means that if the number of occurrences of a particular outcome becomes zero, this outcome is expelled from the set of possible outcomes, so that their number reduces from r to $(r - 1)$ and the r -dimensional vector of occurrences $\mathbf{k}$ in (17) is replaced by the $(r - 1)$ -dimensional vector

$$\mathbf{k}/k_i = [k_1 \ k_2 \ \dots \ k_{i-1} \ k_{i+1} \ \dots \ k_r] \quad (20)$$

The recurrence in (16) and its boundary condition in (19) reduce for $r = 2$ to the recurrence in (2a) and its boundary conditions in (2b). Figure 3 generalizes Fig. 1 for the trinomial case ($r = 3$), and displays a SFG representing (16) and (19), in which the evaluation of $M([1 \ 1 \ 1]^T, 3, 3)$ is obtained recursively as the sum of three equal entities, namely

$$M([1 \ 1 \ 1]^T, 3, 3) = M([0 \ 1 \ 1]^T, 2, 3) + M([1 \ 0 \ 1]^T, 2, 3) + M([1 \ 1 \ 0]^T, 2, 3) \quad (21)$$

The first of the coefficients in the RHS of (21) is given as

$$M([0 \ 1 \ 1]^T, 2, 3) = M([1 \ 1]^T, 2, 2) = M([0 \ 1]^T, 1, 2) + M([1 \ 0]^T, 1, 2) \quad (22)$$

Now, the two coefficients in the RHS of (22) are given by

$$M([0 \ 1]^T, 1, 2) = M([1 \ 0]^T, 1, 2) = M([1]^T, 1, 1) = 1 \quad (23)$$

The above results mean that

$$M([0 \ 1 \ 1]^T, 2, 3) = 1 + 1 = 2 \quad (24)$$

$$M([0 \ 1 \ 1]^T, 2, 3) = M([1 \ 0 \ 1]^T, 2, 3) = M([1 \ 1 \ 0]^T, 2, 3) = 2 \quad (25)$$

$$M([1 \ 1 \ 1]^T, 3, 3) = 2 + 2 + 2 = 6 \quad (26)$$

Each of the values in (23)-(26) can be checked *via* the basic definition (12).

5. Recurrence for Multinomial Probabilities

The recursive relation for multinomial probability can be obtained *via* straightforward extension of (7) or (7a) (for $k_i \geq 0$, and $\mathbf{k} \neq \mathbf{0}$)

$$E(\mathbf{k}, n, \mathbf{p}) = \sum_{i=1}^r (P_{in} * E(\mathbf{k}|k_i \leftarrow k_i - 1, n - 1, \mathbf{p}/p_i)) \quad (27)$$

The above recurrence is subject to the following boundary conditions:

$$E(\mathbf{k}, n, \mathbf{p}) = 1.0 \text{ for } \mathbf{k} = \mathbf{0} \text{ (} n = 0 \text{ or } \mathbf{p} = \emptyset = \text{the empty matrix)} \quad (28)$$

$$E(\mathbf{k}, n, \mathbf{p}) = 0.0 \text{ when one particular } k_i = -1, \quad n \geq 0 \quad (29)$$

Figure 4 is a Signal Flow Graph (SFG) representing the computation of the trinomial probability $E(\mathbf{k}, n, \mathbf{p})$, where $\mathbf{p}$ is a 3-by- n matrix, whose columns (probabilities of outcomes in various trials) are distinct. Again, this SFG is acyclic (loopless). It has a striking similarity to the Pascal Pyramid in Fig. 3, and it is a 3-dimensional extension of Fig. 2. The SFG in Fig. 4 has a single real source of value 1.0 (highlighted as black) while the SFG in Fig.3 has a triply-infinite similar black nodes of value 1.0 each. In Fig. 4, a doubly-infinite number of fictitious zero nodes are assumed to exist at each of the three planes $k_1 = -1$, $k_2 = -1$ and $k_3 = -1$, However, these zero nodes are conveniently omitted in Fig. 4.

6. Iteration Versus Recursion

Direct use of recursion constitutes a “lazy” way (from programming point of view) to compute any of the four quantities discussed herein. A recursive program might be written as follows (based on the assumption that applying the recurrence at a point in its region of validity does not continue indefinitely, but ultimately comes to an end at the boundary in a finite number of steps)

If (Conditions of the boundary are satisfied)

Then Use Boundary conditions

Else Utilize validity of recursion by implementing it and reducing the problem to two sub-problems for which the boundary is to be reached in fewer steps than those needed for the parent problem;

For example, the multinomial coefficient might be computed *via* the Matlab code

```
function S = multm(K)
    S = 0;
    sort(K);
    K(K==0) = [];
    n = sum(K);
    r = length(K);
    if r > 1
        for i=1:r
            X = K(i)-1;
            Y = K;
            Y(i) = X;
            Y(Y==0) = [];
            S = S+multm(Y);
```

```

        end
    else
        S = 1;
    end
end

k = input('Enter the elements of k: ');
tic
R = multm(k);
toc
disp('The multinomial is')
disp(R)

```

Despite the elegant simplicity of the code above, it is a seriously inefficient algorithm for large values of n . This brings us to the heart of the problem of recursion versus iteration [1-3, 5-7, 10, 20, 23, 24, 61-72]. Figure 5 pinpoints the main trouble with recursion, with its unwarranted aggravation of both temporal and spatial complexities. In fact, use of recursion amounts to a need for constructing a recursion stack that needs a lot of time and memory as it continues to grow till each time the boundary is hit (at the leaves of the recursion tree [6]), upon which the stack growth is temporarily reversed, so that ultimately the stack diminishes with the required answer obtained. Figure 5 demonstrates the recursion stack needed for the computation of $M([1\ 3\ 2]^T, 6, 3)$ with arrows denoting its growth and diminishing directions.

The Signal Flow Graphs represented herein are truly insightful for recursion elimination or removal [6, 10, 73-80], *i.e.*, for replacing a recursive algorithm by a non-recursive or iterative one. Non-recursive algorithms already exist for the binomial coefficients [6, 20] and for the binomial probability [1-3, 6, 10, 23, 24]. In these iterative algorithms, one can minimize computer storage by using a strategy of row, column, or diagonal sweep in which the contents of a predecessor entity (row or column or diagonal) are safely overwritten by those of a corresponding successor entity [1, 6, 10]. One can also avoid or significantly reduce the danger of overflow encountered when computing binomial coefficients *via* factorials [24].

In passing, we note that while the non-recursive algorithm for the binomial coefficient seems to have been known for millennia (definitely, for several centuries before Pascal), the analogous non-recursive algorithm for the (generalized) binomial probability seems to have appeared only recently. This latter algorithm is called the BH1 algorithm [1, 6, 10], and is deduced *via* generating functions by Barlow and Heidtmann [81]. It was later re-obtained *via* recursion removal by Rushdi [1, 6, 10] through an insightful development, in which SFGs played a pivotal role. The BH1 algorithm combined simplicity and

efficiency (it has quadratic temporal complexity, and was believed for a while to be optimal). Later, Belfore [82] presented a more elaborate algorithm, which is currently believed to be the optimal non-recursive algorithm for the (generalized) binomial probability.

7. Conclusions

This paper offers a tutorial exposition of four recursively-defined combinatorial functions, *viz.*, the binomial and multinomial coefficients and probabilities. Each of these functions is characterized by a certain recurrence over a specific region of validity together with given boundary conditions. These boundary conditions are along straight lines in a two-dimensional space for the binomial functions. Analogously, the boundary conditions extend along hyper-planes in the n -dimensional space for the multinomial functions. The region of validity is infinite as it extends without bounds when n tends to infinity. Otherwise, this region is bounded by the hyper-planes at which the boundary conditions hold.

A notable contribution of the paper is the visualization of the aforementioned recurrences and boundary conditions *via* acyclic Signal Flow Graphs (SFGs) in the two-dimensional or n -dimensional space. Each of the four functions is amenable to evaluation *via* recursive algorithms, which are elegant (albeit temporally and spatially inefficient). The SFG interpretation proved really handy for the construction of efficient iterative solution for the two binomial functions. Similar efficient iterative algorithms for computing the two multinomial functions are warranted, and will be possibly constructed with the aid of the pertinent SFGs.

References

- [1] Rushdi, A. M., Utilization of symmetric switching functions in the computation of k -out-of- n system reliability, *Microelectronics and Reliability*, 26(5): 973-987, (1986).
- [2] Rushdi, A. M., Efficient computation of k -to- l -out-of- n system reliability, *Reliability Engineering*, 17(3): 157-163, (1987), Erratum: *ibid*, 19(4): 321, (1987).
- [3] Rushdi, A. M., and Dehlawi F., Optimal computation of k -to- l -out-of- n system reliability, *Microelectronics and Reliability*, 27(5): 875-896, (1987), Erratum : *ibid*, 28(4): 671, (1988).
- [4] Rushdi, A. M., Threshold systems and their reliability, *Microelectronics and Reliability*, 30(2): 299-312, (1990).

- [5] Rushdi, A. M., Comment on: An efficient non-recursive algorithm for computing the reliability of k-out-of-n systems, *IEEE Transactions on Reliability*, 40(1): 60-61, (1991).
- [6] Rushdi, A. M., *Reliability of k-out-of-n Systems*, Chapter 5 in Misra, K. B. (Editor), *New Trends in System Reliability Evaluation*, Vol. 16, Fundamental Studies in Engineering, Elsevier Science Publishers, Amsterdam, The Netherlands, 185-227, (1993).
- [7] Rushdi, A. M., & Al-Thubaity, A. O. (1993). Efficient computation of the sensitivity of k-out-of-n system reliability. *Microelectronics and Reliability*, 33(13), 1963-1979.
- [8] Haynes, S. M. (1995). Explaining recursion to the unsophisticated. *ACM SIGCSE Bulletin*, 27(3), 3-6.
- [9] Neuwirth, E. (2001). Recursively defined combinatorial functions: extending Galton's board. *Discrete Mathematics*, 239(1-3), 33-51.
- [10] Rushdi, A. M., Partially-redundant systems: Examples, reliability, and life expectancy, *International Magazine on Advances in Computer Science and Telecommunications*, 1(1): 1-13, (2010).
- [11] Spivey, M. Z. (2011). On solutions to a general combinatorial recurrence. *Journal of Integer Sequences*, 14(9), 1-19.
- [12] Mansour, T., & Shattuck, M. (2013). A combinatorial approach to a general two-term recurrence. *Discrete Applied Mathematics*, 161(13-14), 2084-2094.
- [13] Rushdi, M. A. M., Ba-Rukab, O. M. & Rushdi, A. M. (2016). Multidimensional recursive relation and mathematical induction techniques: The case of failure frequency of k-out-of-n systems. *Journal of king Abdulaziz University: Engineering Science*, 27(2): 15-31.
- [14] Rushdi, A. M. A., & Bjaili, H. A. (2016). An ROBDD algorithm for the reliability of double-threshold systems. *British Journal of Mathematics and Computer Science*, 19(6), 1-17.
- [15] Spivey, M. Z. (2016). The Binomial Recurrence. *Mathematics Magazine*, 89(3), 192-195.
- [16] Rushdi, A. M., & Rushdi, M. A. (2017). Switching-algebraic analysis of system reliability, Chapter 6 in M. Ram and P. Davim (Editors), *Advances in Reliability and System Engineering*. Springer International Publishing, Cham, Switzerland, pp. 139-161.
- [17] Rushdi, A. M. A., & Alturki, A. M. (2017). Computation of k-out-of-n System Reliability via Reduced Ordered Binary Decision Diagrams, *British Journal of Mathematics & Computer Science*, 22(3), 1-9.
- [18] Rushdi, A. M. A., & Alturki, A. M. (2018). Novel Representations for coherent threshold reliability system, *Turkish Journal of Electrical Engineering and Computer Sciences*, 26(1): 257-269.

- [19] Rushdi A. M. A., and Alturki, A. M., Unification of mathematical concepts and algorithms of k-out-of-n system reliability: A perspective of improved disjoint products, *Journal of Engineering Research*, 6(1), (2018).
- [20] Manolopoulos, Y. (2002). Binomial coefficient computation: recursion or iteration?. *ACM SIGCSE Bulletin*, 34(4), 65-67.
- [21] Trivedi, K. S., *Probability & Statistics with Reliability, Queuing, and Computer Science Applications*, 2nd Edition, Prentice-Hall, Englewood Cliffs, NJ, USA, (2002).
- [22] Paoletta, M. S. (2006). *Fundamental Probability: A Computational Approach*. Wiley, New York, NY. USA.
- [23] Rushdi, A. M., & Al-Qasimi, A. M. (1994). Efficient computation of the pmf and the CDF of the generalized binomial distribution. *Microelectronics Reliability*, 34(9), 1489-1499.
- [24] Al-Qasimi, A. M., & Rushdi, A. M. A. (2008). A tutorial on how to efficiently calculate and format tables of the binomial distribution. *Engineering Sciences*, 19(1), 3-17.
- [25] Tauber, S. (1963). On multinomial coefficients. *American Mathematical Monthly*, 1058-1063.
- [26] Abramson, M. (1968). Multinomial coefficients. *Mathematics Magazine*, 41(4), 199-205.
- [27] Staib, J., & Staib, L. (1978). The Pascal Pyramid. *The Mathematics Teacher*, 71(6), 505-510.
- [28] Schwandt, A., & Lind, J. (1979). Revisiting Pascal's pyramid. *The Mathematics Teacher*, 72(2), 84-86.
- [29] Walser, H. (2000). The Pascal pyramid. *The College Mathematics Journal*, 31(5), 383-392.
- [30] Hilton, P., Holton, D., & Pedersen, J. (2002). From Binomial to Trinomial Coefficients and Beyond. In *Mathematical Vistas* (pp. 159-198). Springer, New York, NY.
- [31] Horn, M. E. (2003). Pascal pyramids, Pascal hyper-pyramids and a bilateral multinomial theorem. *arXiv preprint math/0311035*.
- [32] Putz, J. F. (1986). The Pascal polytope: An extension of Pascal's triangle to N dimensions. *The College Mathematics Journal*, 17(2), 144-155.
- [33] Mo, Y., A multiple-valued decision-based approach to solve dynamic fault tree, *IEEE Transactions on Reliability*, 63(1): 81-93, (2014).
- [34] Mo, Y., Xing , L., Amari, S. V., and Dugan, J. B., Efficient analysis of multi-state k-out-of-n system, *Reliability Engineering & System Safety*, 133: 95-105, (2015).
- [35] Mason, S. J. (1953). Feedback theory-some properties of signal flow graphs. *Proceedings of the IRE*, 41(9), 1144-1156.
- [36] Mason, S. J. (1956). Feedback theory: Further properties of signal flow graphs. *Proceedings of the IRE*, 44(7), 920-926.

- [37] Golnaraghi, F., & Kuo, B. C. (2010). Automatic Control Systems. Ninth Edition, Wiley, New York, NY, USA.
- [38] Affandi, A. M., & Rushdi, A. M. (1995, July). An accurate six-port microstrip reflectometer. In Microwave and Optoelectronics Conference, 1995. Proceedings of the 1995 SBMO/IEEE MTT-S International (Vol. 2, pp. 563-568). IEEE.
- [39] Rushdi, A. A. (2010). A mathematical model of DNA replication. International Magazine on Advances in Computer Science and Telecommunications (IMACST), 1(1), 23-30.
- [40] Rushdi, R. A. & Rushdi, A. M. (2018). Karnaugh-map utility in medical studies: The case of Fetal Malnutrition. International Journal of Mathematical, Engineering and Management Sciences, 3(3), 220-244, Available at www.ijmems.in/ijmems—volumes.html.
- [41] Rushdi, A. M., & Talmees, F. A. (2018). An exposition of the eight basic measures in diagnostic testing using several pedagogical tools. Journal of Advances in Mathematics and Computer Science, 26(3), 1-17.
- [42] Rushdi, R. A., Rushdi, A. M. & Talmees, F. A. (2018). Novel pedagogical methods for conditional-probability computations in medical disciplines, Journal of Advances in Medicine and Medical Research, 25(10), 1-15.
- [43] Smith, K. J. (1973). Pascal's triangle. The Two-Year College Mathematics Journal, 4(1), 1-13.
- [44] Curley, W. (1974). Programming Pascal's Triangle. Mathematics Teacher, 67(8), 705-708.
- [45] Crouse, R. (1978). Binomial Coefficients and the Partitioning of n-dimensional Space. The Mathematics Teacher, 71(8), 698-701.
- [46] Hilton, P., & Pedersen, J. (1987). Looking into Pascal's triangle: Combinatorics, arithmetic, and geometry. Mathematics Magazine, 60(5), 305-316.
- [47] Stojmenovic, I. (2000). Recursive algorithms in computer science courses: Fibonacci numbers and binomial coefficients. IEEE Transactions on Education, 43(3), 273-276.
- [48] Edelman, A., & Strang, G. (2004). Pascal matrices. The American Mathematical Monthly, 111(3), 189-197.
- [49] Zeitler, H. (2002). Trinomial formula, Pascal-and Sierpinski-pyramid. International Journal of Mathematical Education in Science and Technology, 33(2), 256-266.
- [50] Baker, J. E. (2013). Pascal Pyramids: a mathematical exploration using spreadsheets. Spreadsheets in Education (eJSiE), 6(2), 1-7.
- [51] Hoggatt Jr, V. E., & Alexanderson, G. L. (1971). A property of multinomial coefficients. Fibonacci Quarterly, 9(4), 351-356.

- [52] Bondarenko, B. A. (1993). Generalized Pascal triangles and pyramids: their fractals, graphs, and applications. Santa Clara, CA: Fibonacci Association.
- [53] Mueller, S. (1969). Recursions associated with Pascal's pyramid. *Pi Mu Epsilon Journal*, 4(10), 417-422.
- [54] Lucas, J. F. (1983). Paths and Pascal numbers. *The Two-Year College Mathematics Journal*, 14(4), 329-341.
- [55] Vincenzi, G., & Siani, S. (2014). Fibonacci-like sequences and generalized Pascal's triangles. *International Journal of Mathematical Education in Science and Technology*, 45(4), 609-614.
- [56] Wolfram, S. (1984). Geometry of binomial coefficients. *The American Mathematical Monthly*, 91(9), 566-571.
- [57] Berggren, J. L. (2007) Mathematics in Medieval Islam. In Katz, V. (Ed.) *The Mathematics of Egypt, Mesopotamia, China, India, and Islam*. (pp.515-670). Princeton University Press, New Jersey.
- [58] Kaddoura, I., & Zeid, M. (2015). A Triangle Analog to Pascal's characterizing primes. *arXiv preprint arXiv:1510.02498*.
- [59] Rolfe, T. (2001). Binomial coefficient recursion: the good, and the bad and ugly. *SIGCSE Bulletin*, 33(2), 35-36.
- [60] Zehna, P. W. (1987). How electronics ended the Poisson approximation to the binomial distribution. *American Mathematical Monthly*, 94(10), 984-987.
- [61] Robinson, R. M. (1947). Primitive recursive functions. *Bulletin of the American Mathematical Society*, 53(10), 925-942.
- [62] Axt, P. (1965). Iteration of primitive recursion. *Mathematical Logic Quarterly*, 11(3), 253-255.
- [63] Burstall, R. M., & Darlington, J. (1977). A transformation system for developing recursive programs. *Journal of the ACM (JACM)*, 24(1), 44-67.
- [64] Bird, R. S. (1977). Improving programs by the introduction of recursion. *Communications of the ACM*, 20(11), 856-863.
- [65] Karp, R. M., & Tarjan, R. E. (1980). Linear expected-time algorithms for connectivity problems. *Journal of Algorithms*, 1(4), 374-393.
- [66] Kessler, C. M., & Anderson, J. R. (1986). Learning flow of control: Recursive and iterative procedures. *Human-Computer Interaction*, 2(2), 135-166.
- [67] Wiedenbeck, S. (1988, February). Learning recursion as a concept and as a programming technique. In *ACM SIGCSE Bulletin* (Vol. 20, No. 1, pp. 275-278). ACM.
- [68] Wiedenbeck, S. (1989). Learning iteration and recursion from examples. *International Journal of Man-Machine Studies*, 30(1), 1-22.
- [69] Shavlik, I. W. (1990). Acquiring recursive and iterative concepts with explanation-based learning. *Machine Learning*, 5(1), 39-70.

- [70] Quesada, A. (1999). Should iteration and recursion be part of the secondary student's mathematics toolbox?. *International Journal of Computer Algebra in Mathematics Education*, 6(2), 103-16.
- [71] Liu, Y. A., & Stoller, S. D. (1999). From recursion to iteration: what are the optimizations?. *ACM Sigplan Notices*, 34(11), 73-82.
- [72] Busi, N., Gabbrielli, M., & Zavattaro, G. (2004, July). Comparing recursion, replication, and iteration in process calculi. In *International Colloquium on Automata, Languages, and Programming* (pp. 307-319). Springer, Berlin, Heidelberg.
- [73] Bird, R. S. (1977). Notes on recursion elimination. *Communications of the ACM*, 20(6), 434-439.
- [74] Auslander, M. A., & Strong, H. R. (1978). Systematic recursion removal. *Communications of the ACM*, 21(2), 127-134.
- [75] Rohl, J. S. (1981). Eliminating recursion from combinatoric procedures. *Software: Practice and Experience*, 11(8), 803-817.
- [76] Cohen, N. H. (1983). Eliminating redundant recursive calls. *ACM Transactions on Programming Languages and Systems (TOPLAS)*, 5(3), 265-299.
- [77] Colussi, L. (1984). Recursion as an effective step in program development. *ACM Transactions on Programming Languages and Systems (TOPLAS)*, 6(1), 55-67.
- [78] Boiten, E. A. (1992). Improving recursive functions by inverting the order of evaluation. *Science of Computer Programming*, 18(2), 139-179.
- [79] Regehr, J., Reid, A., & Webb, K. (2003, October). Eliminating stack overflow by abstract interpretation. In *International Workshop on Embedded Software* (pp. 306-322). Springer, Berlin, Heidelberg.
- [80] Schaeckeler, S., & Shang, W. (2007, September). Stack size reduction of recursive programs. In *Proceedings of the 2007 international conference on Compilers, architecture, and synthesis for embedded systems* (pp. 48-52). ACM.
- [81] Barlow, R. E. & Heidtmann, K. D. (1984). Comment and Reply on 'Computing k-out-of-n system reliability,' *IEEE Transactions on Reliability*, R-33(4): 322-323.
- [82] Belfore, II, L. A. (1995). An $O(n (\log_2 n)^2)$ Algorithm for computing the reliability of k-out-of-n: G & k-to-1-out-of-n: G systems, *IEEE Transactions on Reliability*, 44 (1): 132-136.




Fig. 1. Two Dimensional Signal Flow Graph for computing the binomial coefficients $B(k_1, k_1 + k_2)$. If only the top left triangular half of the shown square is retained, the figure becomes a Pascal Triangle (also called al-Karkhi Triangle or al-Khayyam Triangle). The value of a non-source node at (k_1, k_2) is $B(k_1, k_1 + k_2)$ and equals the number of paths from source (black) nodes to this node. For example, the non-source nodes in the rightmost column shown are $B(1, 6) = 6$, $B(2, 7) = 21$, $B(3, 8) = 56$, $B(4, 9) = 126$, and $B(5, 10) = 252$.




Fig. 2. Two Dimensional Signal Flow Graph for the probability mass function (pmf) of the binomial distribution




Fig. 3. Multi-Dimensional Signal Flow Graph for computing the trinomial coefficients. This is Pascal Pyramid comprising Pascal Triangles in various planes




Fig. 4. Multi-dimensional Signal Flow Graph for the probability mass function of the multinomial distribution up to two generalized Bernoulli trials. A doubly-infinite number of fictitious zero nodes exist at each of the planes $k_1 = -1$, $k_2 = -1$ and $k_3 = -1$ but are not shown

$M([1\ 3\ 2]^T, 6, 3) = M([0\ 3\ 2]^T, 5, 3) + M([1\ 2\ 2]^T, 5, 3) + M([1\ 3\ 1]^T, 5, 3)$	$=10 + 30 + 20=60$
$M([0\ 3\ 2]^T, 5, 3) = M([3\ 2]^T, 5, 2)$	$=10$
$M([1\ 2\ 2]^T, 5, 3) = M([0\ 2\ 2]^T, 4, 3) + M([1\ 1\ 2]^T, 4, 3) + M([1\ 2\ 1]^T, 4, 3)$	$=6+ 12 +12=30$
$M([0\ 2\ 2]^T, 4, 3) = M([2\ 2]^T, 4, 2) = M([1\ 2]^T, 3, 2) + M([2\ 1]^T, 3, 2)$	$=3 + 3 =6$
$M([2\ 2]^T, 4, 2) = M([1\ 2]^T, 3, 2) + M([2\ 1]^T, 3, 2)$	$=3 + 3 =6$
$M([1\ 2]^T, 3, 2)$	$=3$
$M([2\ 1]^T, 3, 2) = M([1\ 1]^T, 2, 2) + M([2\ 0]^T, 2, 2)$	$=2+ 1 =3$
$M([1\ 1]^T, 2, 2) = M([0\ 1]^T, 1, 2) + M([1\ 0]^T, 1, 2)$	$=1 +1 =2$
$M([0\ 1]^T, 1, 2) = M([1]^T, 1, 1)$	$=1$
$M([1\ 0]^T, 1, 2) = M([1]^T, 1, 1)$	$=1$
$M([0\ 1\ 2]^T, 3, 3) = M([1\ 0\ 2]^T, 3, 3)$	$=3$
$M([1\ 1\ 1]^T, 3, 3) = M([0\ 1\ 1]^T, 2, 3) + M([1\ 0\ 1]^T, 2, 3) + M([1\ 1\ 0]^T, 2, 3)$	$=2+2+2=6$
$M([1\ 1\ 2]^T, 4, 3) = M([0\ 1\ 2]^T, 3, 3) + M([1\ 0\ 2]^T, 3, 3) + M([1\ 1\ 1]^T, 3, 3)$	$=3 +3+6=12$
$M([1\ 3\ 1]^T, 5, 3) = M([0\ 3\ 1]^T, 4, 3) + M([1\ 2\ 1]^T, 4, 3) + M([1\ 3\ 0]^T, 4, 3)$	$=4 +12+4=20$

Fig. 5. Recursion Stack used in the computation of the multinomial coefficient $M([1\ 3\ 2]^T, 6, 3)$. The downward arrow shows the direction of the initial growth of the stack, while the upward arrow indicates its subsequent decrease in size till ultimately it vanishes