

22887:Ms_JAMCS_38975.pdf

by Louis Asiedu

Submission date: 11-Jan-2018 05:32PM (UTC+0000)

Submission ID: 901876639

File name: ignments_aa5fd530-f276-4482-8589-2d40bfce862b_Ms_JAMCS_38975.pdf (387.64K)

Word count: 2207

Character count: 9784

28

A New Generalization of Weibull Distribution

Original research
paper

Abstract

2

The Lomax-Weibull distribution is introduced as a new generalization of Weibull distribution. Some properties of the new distribution are investigated. Also, maximum likelihood and Bayesian estimation of the four unknown parameters are discussed. The asymptotic variance-covariance matrix is obtained. Finally, a numerical example is provided.

1

Keywords: The Lomax-Weibull (Lomax-W) distribution; maximum likelihood estimates; asymptotic variance-covariance matrix; Bayesian estimation.

2017 Statistics Subject Classification: ;;

1 Introduction

Parametric modeling of data is in the main stream of statistical inference, and the Weibull distribution had been widely used for modeling many types of data, and reliability data in particular. As it has been noted and suggested by many authors, the Weibull distribution as such may not give adequate fits in many data. For this reason, among many others, many authors suggested new families of distributions in the hope of adding flexibility in the modeling process. Many of these families are modification, extension or combinations of existing one. Among these new-families: "the beta-G family([Eugene et al. (2002)], [Jones (2004)]); the gamma-G family(type 1)[Zografos and Balakrishnan (2009)]; the Kumaraswamy-G (Kw-G)family [Cordeiro and de Castro (2011)]; the McDonald-G(Mc-G)family [Alexander et al. (2012)]; exponentiated generalized-G family[Cordeiro et al. (2013)]; the transformed - transformer T-X family[Alzaatreh et al. (2013)]; the exponentiated T-X family[Alzaghal et al. (2013)]; the Lomax-Fréchet and the Lomax-Gumbel distributions[Gupta et al. (2015)] and [Gupta et al. (2016)], respectively"and more. In the following section we will concern with the last one. Gupta et al., 2015 and Gupta et al. (2016) suggested a new obtaining family of distributions from two other families F and G with the following cdf:

$$F_G(x) = \frac{F(G(x))}{F(1)} \quad (1.1)$$

20

and the corresponding pdf is

$$f_G(x) = \frac{F'(G(x))g(x)}{F(1)} \quad (1.2)$$

Let $F(x)$ and $f(x)$ be the cdf and pdf of the first distribution with support $[0, a]$, $1 \leq a < \infty$ and $G(x)$ and $g(x)$ be the cdf and pdf of the second distribution with support $[a, b]$, where $b, c \in R$ or $[0, \infty)$ or $(-\infty, \infty)$. For different choices of first and second distributions one can construct a large number of distributions such as Gupta et al., 2015 introduced the Lomax-Fréchet distribution; Gupta et al., 2016 introduced the Lomax-Gumbel distribution.

2 Lomax-Weibull Distribution

The cdf for Lomax distribution is $F(x) = 1 - (1 + \frac{x}{\lambda})^{-\alpha}$ and the cdf of Weibull distribution is $G(x) = 1 - e^{-\beta x^\gamma}$. It is suggested here to use the idea of Gupta et al. (2015) and Gupta et al. (2016) by putting $F(x)$ and $G(x)$ in formulas (1) and (2), to generate what may be termed as the Lomax-W distribution. The cdf of the Lomax-W distribution is

$$F_{LW}(x) = K[1 - (1 + \frac{1}{\lambda}(1 - e^{-\beta x^\gamma}))^{-\alpha}], \quad (2.1)$$

and

$$f_{LW}(x) = \frac{K\alpha\beta\gamma}{\lambda} x^{\gamma-1} e^{-\beta x^\gamma} [1 - (1 + \frac{1}{\lambda}(1 - e^{-\beta x^\gamma}))^{-\alpha}] \quad (2.2)$$

where the constant $K = [F(1)]^{-1} = [1 - (1 + \frac{1}{\lambda})^{-\alpha}]^{-1}$, and $\alpha, \lambda, \beta, \gamma > 0, x \geq 0$. using (3) and (4), one can obtain the following hazard function:

$$h_{LW}(x) = \frac{K\alpha\beta\gamma}{\lambda} \frac{x^{\gamma-1} e^{-\beta x^\gamma} [1 - (1 + \frac{1}{\lambda}(1 - e^{-\beta x^\gamma}))^{-\alpha}]}{1 - K[1 - (1 + \frac{1}{\lambda}(1 - e^{-\beta x^\gamma}))^{-\alpha}]} \quad (2.3)$$

For different parameter values, figures 1-2 show plots of the pdf and hazard function of the Lomax-W distribution as follows:




Figure 1: pdf of Lomax-W distribution for various values of the parameters




Figure 2: Hazard function of Lomax-W distribution for various values of the parameters

13

The quantile function of the Lomax-W distribution is given by:

$$x_q = \left[-\frac{1}{\beta} \log \left[1 - \lambda \left(\left(1 - \frac{q}{K} \right)^{-\frac{1}{\alpha}} - 1 \right) \right] \right]^{\frac{1}{\gamma}}. \quad (2.4)$$

9

This formula can be used to generate random numbers from the Lomax-W distribution and calculate L-moments.

The r^{th} raw moments for the Lomax-W distribution is

$$E(X^r) = \frac{K\alpha\beta}{\lambda} \sum_{i=0}^{\infty} \sum_{j=0}^i (-1)^j \lambda^{-i} C_i^{-(\alpha+1)} C_j^i (\beta(j+1))^{-(\frac{r}{\gamma}+1)} \Gamma\left(\frac{r}{\gamma}+1\right) \quad (2.5)$$

where $C_j^i = \frac{\Gamma(i+1)}{\Gamma(i-j+1)\Gamma(j+1)}$.

3 Parameter Estimation

1

Now, the maximum likelihood and Bayesian Estimation are discussed.

8

3.1 Maximum Likelihood Estimation

Let $x_1, x_2, \dots, x_n$ be a random sample of size n follows the Lomax-W distribution. To estimate the four unknown parameters $(\alpha, \lambda, \beta, \gamma)$ of the Lomax-W distribution we first obtain the likelihood function as follows:

$$L = \left(\frac{K}{\lambda} \right)^n \prod_{i=1}^n x_i^{\gamma-1} e^{-\beta \sum_{i=1}^n x_i^{\gamma}} \prod_{i=1}^n \left(1 + \frac{1}{\lambda} (1 - e^{-\beta x_i^{\gamma}}) \right)^{-(\alpha+1)}, \quad (3.1)$$

9 and the log-likelihood function is

$$\ln L = n \ln K + n \ln \alpha + n \ln \beta + n \ln \gamma - n \ln \lambda + (\gamma - 1) \sum_{i=1}^n \ln x_i - \beta \sum_{i=1}^n x_i^\gamma - (\alpha + 1) \sum_{i=1}^n \ln \left(1 + \frac{1}{\lambda} (1 - e^{-\beta x_i^\gamma}) \right). \quad (3.2)$$

The first order derivative of the log-likelihood function with respect to α, λ, β and γ are

$$\begin{aligned} \frac{\partial \ln L}{\partial \alpha} &= \frac{n(1 + \lambda^{-1})^{-\alpha} \ln(1 + \lambda^{-1})}{(1 + \frac{1}{\lambda})^{-\alpha}} + \frac{n}{\alpha} - \sum_{i=1}^n \ln \left(1 + \frac{1}{\lambda} (1 - e^{-\beta x_i^\gamma}) \right), \\ \frac{\partial \ln L}{\partial \lambda} &= -\frac{21}{\lambda^2} \frac{(1 + \lambda^{-1})^{-(\alpha+1)}}{(1 + \frac{1}{\lambda})^{-\alpha}} - \frac{n}{\lambda} + \frac{(\alpha + 1)(1 - e^{-\beta x_i^\gamma})}{\lambda^2 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))}, \\ \frac{\partial \ln L}{\partial \beta} &= \frac{n}{\beta} - \sum_{i=1}^n x_i^\gamma - \frac{(\alpha + 1)}{\lambda} \sum_{i=1}^n \frac{x_i^\gamma e^{-\beta x_i^\gamma}}{(1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))}, \\ \frac{\partial \ln L}{\partial \gamma} &= \frac{n}{\gamma} + \sum_{i=1}^n \ln x_i - \beta \sum_{i=1}^n x_i^\gamma \ln x_i - \frac{\beta(\alpha + 1)}{\lambda} \sum_{i=1}^n \frac{e^{-\beta x_i^\gamma} x_i^\gamma \ln x_i}{(1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))}. \end{aligned} \quad (3.3)$$

1 To obtain the estimates of the 1 four unknown parameters, one can equate equations in (10) to zero and solving them numerically. The second order derivative of log-likelihood function are:

$$\begin{aligned} I_{\alpha, \alpha} = \frac{\partial^2 \ln L}{\partial \alpha^2} &= -\frac{n}{\alpha^2} - \frac{n(1 + \lambda^{-1})^{-\alpha} (\ln(1 + \lambda^{-1}))^2}{[1 - (1 + \lambda^{-1})^{-\alpha}]} - \frac{n(1 + \lambda^{-1})^{-2\alpha} (\ln(1 + \lambda^{-1}))^2}{[1 - (1 + \lambda^{-1})^{-\alpha}]^2}, \\ I_{\alpha, \lambda} = \frac{\partial^2 \ln L}{\partial \alpha \partial \lambda} &= -\frac{n(1 + \lambda^{-1})^{-(\alpha+1)} [1 - \alpha \ln(1 + \lambda^{-1})]}{[1 - (1 + \lambda^{-1})^{-\alpha}]} + \frac{n\alpha(1 + \lambda^{-1})^{-(2\alpha+1)} \ln(1 + \lambda^{-1})}{\lambda^2 [1 - (1 + \lambda^{-1})^{-\alpha}]^2} \\ &\quad + \sum_{i=1}^n \frac{(1 - e^{-\beta x_i^\gamma})}{\lambda^2 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))}, \\ I_{\alpha, \beta} = \frac{\partial^2 \ln L}{\partial \alpha \partial \beta} &= -\sum_{i=1}^n \frac{x_i^\gamma e^{-\beta x_i^\gamma}}{\lambda (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))}, \\ I_{\alpha, \gamma} = \frac{\partial^2 \ln L}{\partial \alpha \partial \gamma} &= -\sum_{i=1}^n \frac{\beta x_i^\gamma e^{-\beta x_i^\gamma} \ln x_i}{\lambda (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))}, \\ I_{\lambda, \lambda} = \frac{\partial^2 \ln L}{\partial \lambda^2} &= \frac{n}{\lambda^2} - \sum_{i=1}^n \frac{2(\alpha + 1)(1 - e^{-\beta x_i^\gamma})}{\lambda^3 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))} + \frac{2n\alpha(1 + \lambda^{-1})^{-(\alpha+1)}}{\lambda^3 (1 - (1 + \lambda^{-1})^{-\alpha})} \\ &\quad + \sum_{i=1}^n \frac{(\alpha + 1)(1 - e^{-\beta x_i^\gamma})^2}{\lambda^4 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))^2} - \frac{\alpha(\alpha + 1)(1 + \lambda^{-1})^{-(\alpha+2)}}{\lambda^4 (1 - (1 + \lambda^{-1})^{-\alpha})} - \frac{n\alpha^2(1 + \lambda^{-1})^{-2(\alpha+1)}}{\lambda^4 (1 - (1 + \lambda^{-1})^{-\alpha})^2}, \\ I_{\lambda, \beta} = \frac{\partial^2 \ln L}{\partial \lambda \partial \beta} &= -\sum_{i=1}^n \frac{(\alpha + 1)x_i^\gamma e^{-\beta x_i^\gamma} (1 - e^{-\beta x_i^\gamma})}{\lambda^3 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))^2} + \sum_{i=1}^n \frac{(\alpha + 1)x_i^\gamma e^{-\beta x_i^\gamma}}{\lambda^2 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))}, \\ I_{\lambda, \gamma} = \frac{\partial^2 \ln L}{\partial \lambda \partial \gamma} &= \sum_{i=1}^n \frac{(\alpha + 1)\beta \ln x_i x_i^\gamma e^{-\beta x_i^\gamma}}{\lambda^2 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))} - \sum_{i=1}^n \frac{(\alpha + 1)\beta \ln x_i x_i^\gamma e^{-\beta x_i^\gamma} (1 - e^{-\beta x_i^\gamma})}{\lambda^3 (1 + \lambda^{-1} (1 - e^{-\beta x_i^\gamma}))^2}, \end{aligned} \quad (15)$$

$$\begin{aligned}
 I_{\beta,\beta} &= \frac{\partial^2 \ln L}{\partial \beta^2} = -\frac{n}{\beta^2} + \sum_{i=1}^n \frac{(\alpha+1)x_i^{2\gamma} e^{-\beta x_i^\gamma}}{\lambda(1+\lambda^{-1}(1-e^{-\beta x_i^\gamma}))} + \sum_{i=1}^n \frac{(\alpha+1)x_i^{2\gamma} e^{-2\beta x_i^\gamma}}{\lambda^2(1+\lambda^{-1}(1-e^{-\beta x_i^\gamma}))^2}, \\
 I_{\beta,\gamma} &= \frac{\partial^2 \ln L}{\partial \beta \partial \gamma} = -\sum_{i=1}^n x_i^\gamma \ln x_i + \sum_{i=1}^n \frac{(\alpha+1)\beta \ln x_i x_i^{2\gamma} e^{-\beta x_i^\gamma}}{\lambda^2(1+\lambda^{-1}(1-e^{-\beta x_i^\gamma}))^2} - \sum_{i=1}^n \frac{(\alpha+1)\ln x_i x_i^\gamma e^{-\beta x_i^\gamma} [1-\beta x_i^\gamma]}{\lambda(1+\lambda^{-1}(1-e^{-\beta x_i^\gamma}))}, \\
 I_{\gamma,\gamma} &= \frac{\partial^2 \ln L}{\partial \gamma^2} = -\frac{n}{\gamma^2} - \sum_{i=1}^n \beta x_i^\gamma (\ln x_i)^2 + \sum_{i=1}^n \frac{(\alpha+1)\beta^2 (\ln x_i)^2 x_i^{2\gamma} e^{-2\beta x_i^\gamma}}{\lambda^2(1+\lambda^{-1}(1-e^{-\beta x_i^\gamma}))^2} \\
 &\quad - \sum_{i=1}^n \frac{(\alpha+1)\beta (\ln x_i)^2 x_i^\gamma e^{-\beta x_i^\gamma} [1-\beta x_i^\gamma]}{\lambda(1+\lambda^{-1}(1-e^{-\beta x_i^\gamma}))}.
 \end{aligned}$$

and the observed information matrix is given by

$$J(\theta) = - \begin{bmatrix} I_{\alpha,\alpha} & I_{\alpha,\lambda} & I_{\alpha,\beta} & I_{\alpha,\gamma} \\ I_{\lambda,\alpha} & I_{\lambda,\lambda} & I_{\lambda,\beta} & I_{\lambda,\gamma} \\ I_{\beta,\alpha} & I_{\beta,\lambda} & I_{\beta,\beta} & I_{\beta,\gamma} \\ I_{\gamma,\alpha} & I_{\gamma,\lambda} & I_{\gamma,\beta} & I_{\gamma,\gamma} \end{bmatrix} \quad (3.4)$$

To obtain the asymptotic variance-covariance matrix of $(\hat{\alpha}, \hat{\lambda}, \hat{\beta}, \hat{\gamma})$, one can invert the information matrix and replace the unknown parameters by their mles. So 100(1 - ν)% approximate confidence intervals (CIs) for the parameters α, λ, β and γ are respectively:

$$\begin{aligned}
 &(\hat{\alpha} - z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\alpha})}, \hat{\alpha} + z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\alpha})}), \\
 &(\hat{\lambda} - z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\lambda})}, \hat{\lambda} + z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\lambda})}), \\
 &(\hat{\beta} - z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\beta})}, \hat{\beta} + z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\beta})}), \text{ and} \\
 &(\hat{\gamma} - z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\gamma})}, \hat{\gamma} + z_{\frac{\nu}{2}} \sqrt{\text{var}(\hat{\gamma})}).
 \end{aligned}$$

3.2 Bayesian Estimation

Let non-informative prior distribution take the form $\pi_0(\alpha, \lambda, \beta, \gamma) \propto \frac{1}{\alpha \lambda \beta \gamma}$ where α, λ, β and $\gamma > 0$. To compute approximate Bayes estimates, one can generate samples from posterior distributions using the Gibbs sampling procedure. The joint posterior distribution is

$\pi(\alpha, \lambda, \beta, \gamma | x) \propto \pi_0(\alpha, \lambda, \beta, \gamma) \exp l(x, \alpha, \lambda, \beta, \gamma)$, where $l(x, \alpha, \lambda, \beta, \gamma)$ is the logarithm of the likelihood function is given in (9). When we set $\rho_1 = \log(\beta)$, $\rho_2 = \log(\gamma)$, $\rho_3 = \log(\alpha)$ and $\rho_4 = \log(\lambda)$, the joint prior distribution will be $\pi(\rho_1, \rho_2, \rho_3, \rho_4) \propto \text{constant}$, $-\infty < \rho_1, \rho_2, \rho_3$, and $\rho_4 < \infty$ and the joint posterior distribution take the form

$$\begin{aligned}
 \pi(\rho_1, \rho_2, \rho_3, \rho_4 | x) &\propto \pi(\rho_1, \rho_2, \rho_3, \rho_4) \log[1 - (1 + \exp(-\rho_4))^{-\exp(\rho_3)}] + \rho_1 \\
 &\quad + \rho_2 - \rho_4 + \sum_{i=1}^n (\exp(\rho_2) - 1) \log[x_i] - \sum_{i=1}^n \exp(\rho_1) x_i^{\exp(\rho_2)} \\
 &\quad - (\exp(\rho_3) + 1) \log[1 + \exp(-\rho_4)(1 + \exp^{-\exp(\rho_1) x_i^{\exp(\rho_2)}})],
 \end{aligned}$$

Posterior summaries such as: the mean; the standard deviation; credible interval and others can be calculated using the WinBUGS software.

4 A Numerical Example

Now, we generate a sample of size $n = 100$ from Lomax-W distribution to estimate the four unknown parameters α, λ, β , and γ . For our sample, let $\alpha_0 = 1.5$, $\lambda_0 = 0.5$, $\beta_0 = 1.5$, and $\gamma_0 = 2.5$. The MLEs and 95 % confidence intervals for the four parameters α, λ, β and γ are listed in table(1):

Table 1: the MLEs and CIs for the four parameters α, λ, β and γ of the Lomax-W model

Parameter	Estimate	Standard Deviation	95% Confidence Interval
α	4.3200	1.2649	(1.84077, 6.79923)
λ	0.0140	0.01414	(0.11228, 0.16772)
β	2.4200	1.3693	(0.0000, 5.10384)
γ	5.1100	0.4641	(4.20034, 6.01966)

With the same sample and non-informative uniform priors $U(0.00, 3.20)$, $U(0.001, 0.01)$, $U(0.001, 0.10)$ and $U(-4.00, 3.00)$ for ρ_1, ρ_2, ρ_3 and ρ_4 respectively, we generate a set of 10000 Gibbs samples generated after burn in sample of size 1000. As table (2) shows the MC error is less than 5 % of the sample standard deviation which indicating convergence of the algorithm.

Table 2: Summary results for the posterior parameters in the case of the Lomax-W model

Parameter	Estimate	Standard Deviation	MC error	95 % Credible Interval
α	1.0180	0.01637	2.536E-4	(1.0010, 1.0620)
λ	0.6024	0.07521	6.588E-5	(0.4675, 0.7564)
β	1.0070	0.28680	0.002674	(2.8630, 3.9860)
γ	3.4040	0.00251	3.332E-5	(1.0010, 1.0100)

From Tables 1-2, one can note that in most cases the CIs based on maximum likelihood inference is larger than the credible intervals based on the posterior summaries.

5 CONCLUSIONS

The Weibull distribution may not give adequate fits in many data. For this reason, among many others, many authors suggested new families of distributions in the hope of adding flexibility in the modeling process. Many of these families are modification, extension or combination of existing one. In this paper we concerned with one of the families which introduced by [Gupta et al. (2015)] and [Gupta et al. (2016)]. Using this family we introduced a new generalization of Weibull distribution which called the Lomax-Weibull distribution. Some properties of the new distribution are investigated and maximum likelihood and Bayesian estimation of the four unknown parameters are discussed.

References

- Alexander, C., Cordeiro, G-M., Ortega, E-M-M. and Sarabia, J-M. (2012). Generalized Beta-Generated Distributions. Computational Statistics and Data Analysis 56, 1880-1897.
- Alzaatreh, A., Famoye, F. and Lee, C. A New Method for Generating Families of Continuous Distributions. Metron 71, 63-79.
- Alzaghal, A., Lee, C. and Famoye, F. Exponentiated T-X Family of Distributions with some Applications. International Journal of Probability and Statistics 2, 31-49.
- Cordeiro, G-M. and de Castro, M. A New Family of Generalized Distributions. Journal of Statistical Computation and Simulation 81, (7), 883-898.
- Cordeiro, G-M., Ortega, E-M-M. and da Cunha, D-C-C. The Exponentiated Generalized Class of Distributions. Journal of Data Science 11, 1-27.

- Gupta,V., Bhatt,M. and Gupta,M. The Lomax-Fréchet Distribution.Journal of Rajasthan Academy of Physical14(1),25-43.
- Gupta,J., Garg,M. and Gupta, J. The Lomax-Gumbel Distribution.Palestine Journal of Mathematics5(1), 35-42.
- Eugene,N., Lee,C. and Famoye, F. Beta-Normal Distribution and its Applications.Communications in Statistics-Theory and Methods31, 497-512.
- Jones,M-C. Families of Distributions Arising from Distributions of Order Statistics.Test13(1), 1-43.
- Zografos,K. and Balakrishnan,N. On Families of Beta-and Generalized Gamma-Generated Distributions and Associated Inference.Statistical Methodology6, 344-362.

ORIGINALITY REPORT

34%

SIMILARITY INDEX

17%

INTERNET SOURCES

32%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

R. M. Mandouh. "Logistic-modified Weibull distribution and parameter estimation", International Journal of Contemporary Mathematical Sciences, 2018

Publication

16%

2

www.m-hikari.com

Internet Source

2%

3

Submitted to TechKnowledge Turkey

Student Paper

2%

4

Giles, Emma L., Sharron Kuznesof, Beth Clark, Carmen Hubbard, and Lynn J. Frewer.

"Consumer acceptance of and willingness to pay for food nanotechnology: a systematic review", Journal of Nanoparticle Research, 2015.

Publication

1%

5

fs.gallup.unm.edu

Internet Source

1%

6

www.pjsor.com

Internet Source

1%

7	d-nb.info Internet Source	1 %
8	Tahir, M. H., Gauss M. Cordeiro, M. Mansoor, M. Zubair, and Morad Alizadeh. "The Weibull-Dagum Distribution: Properties and Applications", Communication in Statistics-Theory and Methods, 2016. Publication	1 %
9	Understanding Complex Systems, 2011. Publication	1 %
10	Aryal, Gokarna R. and Tsokos, Chris P.. "Transmuted Weibull Distribution: A Generalization of the Weibull Probability Distribution", European Journal of Pure & Applied Mathematics, 2011. Publication	1 %
11	Submitted to University of Washington Student Paper	1 %
12	oatao.univ-toulouse.fr Internet Source	1 %
13	Submitted to Covenant University Student Paper	1 %
14	www.btdt.ufpe.br Internet Source	1 %

15

Internet Source

<1 %

16

Elbatal, Ibrahim, Francesca Condino, and Filippo Domma. "Reflected Generalized Beta Inverse Weibull Distribution: definition and properties", Sankhya B, 2016.

Publication

<1 %

17

Zaindin, Mazen and Sarhan, Ammar M.. "NEW GENERALIZED WEIBULL DISTRIBUTION", Pakistan Journal of Statistics, 2011.

Publication

<1 %

18

Noga Alon. "Coin-Flipping Games Immune against Linear-Sized Coalitions", SIAM Journal on Computing, 1993

Publication

<1 %

19

www3.ufpe.br

Internet Source

<1 %

20

Submitted to CVC Nigeria Consortium

Student Paper

<1 %

21

A. Varchenko. "Resonance Relations, Holomorphic Trace Functions and Hypergeometric Solutions to qKZB and Macdonald Ruijsenaars Equations", International Mathematics Research Papers, 12/18/2007

Publication

<1 %

22	ww2.ppgbea.ufrpe.br Internet Source	<1 %
23	link.springer.com Internet Source	<1 %
24	home.tu-clausthal.de Internet Source	<1 %
25	Lee, Carl, Felix Famoye, and Ayman Y. Alzaatreh. "Methods for generating families of univariate continuous distributions in the recent decades", Wiley Interdisciplinary Reviews Computational Statistics, 2013. Publication	<1 %
26	jsdajournal.springeropen.com Internet Source	<1 %
27	www.scienpress.com Internet Source	<1 %
28	M.R. Mahmoud, R.M. Mandouh. "The (P-A-L) generalized exponential distribution: properties and estimation", International Mathematical Forum, 2017 Publication	<1 %

