

A SELF-STARTING FIVE-STEP EIGHT-ORDER BLOCK METHOD FOR STIFF ORDINARY DIFFERENTIAL EQUATIONS

Abstract. This paper examines the implementation of a self-starting five-step eight-order block method with two off-grid for stiff ordinary differential equations using interpolation and collocation procedures. The predictor schemes are the expanded using Taylor's series expansion. Multiple numerical integrators were produce and arrived at a discrete scheme. The discrete schemes are of uniform order eight and are assembled into a single block matrix equation. These equations are simultaneously applied to provide the approximate solution for stiff initial value problem for ordinary differential equations. The order of accuracy and stability of the block method is discussed and its accuracy is established numerically.

Keywords: block method, stiff, five-step, power series.

1. Introduction

A considerable literature exist for the conventional k-step linear multi-step methods for the solution of ordinary differential equations (ODE's) of the form

$$y' = f(x, y), \quad y(x_0) = y_0, \quad x \in [a, b] \quad (1)$$

The solution is in the range $a \leq x \leq b$, where a and b are finite and we assumed that f satisfies the Lipschitz condition, which guarantees the existence and uniqueness of the solution of the problem (1).

For the discrete solution of (1) by linear multi-step method has being studied by authors like [1] and continuous solution of (1) [2] and [3], [4]. One important advantage of the continuous over discrete approach is the ability to provide discrete schemes for simultaneous integration. These discrete schemes can be reformulated as general linear methods (GLM) [5]. The block methods are self-starting and can be applied to both stiff and non-stiff initial value problem in differential equations.

Definition 1. A stiff equation is a differential equations that are characterized as those whose exact solutions has a term of the form e^{-ct} , where C is a large positive constant.

More recently, authors like [6]), [7]), [8]), [9] and [10] to mention few, these authors proposed methods ranging from predictor- corrector to hybrid block method for initial value problem in ordinary differential equation.

In this work, we derived the continuous self-starting five-step order-eight block method with two off-grid using Taylor's series expansion. This would help in coming up with a more computationally reliable integrator that could solve stiff differential equations problems of the form (1).

2. Derivation technique of the self-starting block method

We consider the continuous hybrid formula to be the form

$$y(x) = \sum_{j=0}^{s+t-1} \alpha_j x^j \quad (2)$$

33 Interpolation and collocation procedures are use by choosing interpolation point t at a grid point and collocation
 34 points s at all points giving rise to $\xi = t + s$ system of equations whose coefficients are determined by using
 35 appropriate procedures. The first derivative of (2) is given by;

$$36 \quad y' = \sum_{j=0}^{s+t-1} j \alpha_j x^{j-1} \quad (3)$$

37 Then, substituting (3) in (1), we obtain

$$38 \quad f(x, y)' = \sum_{j=0}^{s+t-1} j \alpha_j x^{j-1} \quad (4)$$

39 Now, collocating (4) at point x_{n+s} , $s = 0(1)5$ and interpolating (2) at points $x_{n+\frac{3}{2}}$ and $x_{n+\frac{5}{2}}$ leads to the
 40 following system of equations;

$$41 \quad AX = U \quad (5)$$

$$42 \quad A = [a_0 \quad a_1 \quad a_2 \quad a_3 \quad a_4 \quad a_5 \quad a_6 \quad a_7 \quad a_8]^T,$$

$$43 \quad U = \begin{bmatrix} y_n & f_n & f_{n+1} & f_{n+\frac{3}{2}} & f_{n+2} & f_{n+\frac{5}{2}} & f_{n+3} & f_{n+4} & f_{n+5} \end{bmatrix}^T,$$

$$44 \quad X = \begin{bmatrix} 1 & x_n & x_n^2 & x_n^3 & x_n^4 & x_n^5 & x_n^6 & x_n^7 & x_n^8 \\ 0 & 1 & 2x_n & 3x_n^2 & 4x_n^3 & 5x_n^4 & 6x_n^5 & 7x_n^6 & 8x_n^7 \\ 0 & 1 & 2x_{n+1} & 3x_{n+1}^2 & 4x_{n+1}^3 & 5x_{n+1}^4 & 6x_{n+1}^5 & 7x_{n+1}^6 & 8x_{n+1}^7 \\ 0 & 1 & 2x_{n+\frac{3}{2}} & 3x_{n+\frac{3}{2}}^2 & 4x_{n+\frac{3}{2}}^3 & 5x_{n+\frac{3}{2}}^4 & 6x_{n+\frac{3}{2}}^5 & 7x_{n+\frac{3}{2}}^6 & 8x_{n+\frac{3}{2}}^7 \\ 0 & 1 & 2x_{n+2} & 3x_{n+2}^2 & 4x_{n+2}^3 & 5x_{n+2}^4 & 6x_{n+2}^5 & 7x_{n+2}^6 & 8x_{n+2}^7 \\ 0 & 1 & 2x_{n+\frac{5}{2}} & 3x_{n+\frac{5}{2}}^2 & 4x_{n+\frac{5}{2}}^3 & 5x_{n+\frac{5}{2}}^4 & 6x_{n+\frac{5}{2}}^5 & 7x_{n+\frac{5}{2}}^6 & 8x_{n+\frac{5}{2}}^7 \\ 0 & 1 & 2x_{n+3} & 3x_{n+3}^2 & 4x_{n+3}^3 & 5x_{n+3}^4 & 6x_{n+3}^5 & 7x_{n+3}^6 & 8x_{n+3}^7 \\ 0 & 1 & 2x_{n+4} & 3x_{n+4}^2 & 4x_{n+4}^3 & 5x_{n+4}^4 & 6x_{n+4}^5 & 7x_{n+4}^6 & 8x_{n+4}^7 \\ 0 & 1 & 2x_{n+5} & 3x_{n+5}^2 & 4x_{n+5}^3 & 5x_{n+5}^4 & 6x_{n+5}^5 & 7x_{n+5}^6 & 8x_{n+5}^7 \end{bmatrix}$$

45 Solving (5), for α'_j $s, j = 0(1)8$ and substituting back into (2) gives a continuous linear multistep method of the
 46 form

$$47 \quad y(x) = \alpha_0(x) y_n + h \sum_{j=0}^7 \beta_j(x) f_{n+j}, \quad (6)$$

48 Where $\alpha_o = 1$ and the coefficient of f_{n+j} gives

$$\begin{aligned}\beta_o &= \frac{-1}{75600} x^5 \left(-20874h^3 - 24990x_n h - 9576x_n^2 h - 1176x_n^3 + 4165xh^2 + 3192xx_n h \right) / h^7 \\ \beta_1 &= \frac{1}{30240} x^4 \left(\begin{aligned} &420945h^5 + 824040x_n h^3 + 549150x_n^2 h^2 + 151200x_n^3 h + 14700x_n^4 - 164808xh^3 \\ &219660xx_n h^2 - 90720xx_n^2 h - 11760xx_n^3 + 36610x^2 x_n h + 5880x^2 x_n^2 - 4320x^3 h \\ &- 1680x^3 x_n + 210x^4 \end{aligned} \right) / h^7 \\ \beta_2 &= \frac{-8}{945} x^5 \left(\begin{aligned} &-2100h^3 - 2940x_n h^2 - 168x_n^3 - 1260x_n^2 h + 490xh^2 + 84xx_n^2 + 420xx_n h - 24x^2 x_n \\ &- 60x^2 h + 3x^3 \end{aligned} \right) / h^7 \\ \beta_3 &= \frac{1}{1680} x^4 \left(\begin{aligned} &99295h^4 + 219380x_n h^3 + 160650x_n^2 h^2 + 47600x_n^3 h + 4900x_n^4 - 43876xh^{10} \\ &- 64260xh^9 x_n - 28560xh^8 x_n^2 - 3920xh^7 x_n^3 + 10710x^2 h^2 + 9520x^2 x_n h + 1960x^2 x_n^2 \\ &- 1360x^3 h - 560x^3 x + 70x^4 \end{aligned} \right) / h^7 \\ \beta_4 &= \frac{-8}{4725} x^4 \left(\begin{aligned} &59220x_n h^3 + 45150x_n^2 h^2 + 13860x_n^3 h + 1470x_n^4 + 25683h^4 - 11844xh^3 - 18060xx_n h^2 \\ &- 8316xx_n^2 h - 1176xx_n^3 + 3010x^2 h^2 + 2772x^2 x_n h + 588x^2 x_n^2 - 396x^3 h - 168x^3 x_n + 21x^4 \end{aligned} \right) / h^7\end{aligned}$$

49

$$\beta_5 = \frac{1}{15120} x^4 \left(\begin{aligned} &423150x_n^2 h^2 + 134400x_n^3 h + 14700x_n^4 + 535920x_n h^3 - 225015h^4 + 169260xx_n h^2 \\ &- 80640xx_n^2 h - 11760xx_n^3 - 107184xh^3 + 28210x^2 h^2 + 26880x^2 x_n h + 5880x^2 x_n^2 \\ &- 3840x^3 h - 1680x^3 x_n + 210x^4 \end{aligned} \right) / h^7$$

50

$$\beta_6 = \frac{-1}{30240} x^4 \left(\begin{aligned} &25200x_n^3 h + 2940x_n^4 + 74550x_n^2 h^2 + 89460x_n h^3 + 37191h^4 - 15120xx_n^2 - 2352xx_n^3 \\ &- 29820xx_n h^2 - 17892xh^3 + 5040x^2 x_n h + 4970x^2 h^2 - 720x^3 h - 336x^3 x_n + 42x^4 \end{aligned} \right) / h^7$$

$$\beta_7 = \frac{1}{8400} x^5 \left(-56x_n^3 - 336x_n^2 h - 630x_n h^2 - 364h^3 + 28x_n^2 x + 112xx_n h + 105xh^2 - 8x^2 x_n - 16x^2 h + x^3 \right) / h^7$$

51

(7)

52 Evaluating (7) at $x = 1(1)7$ gives a continuous discrete block formula of the form

53

$$A^{(0)}Y_m = Ey_n + hdf(y_n) + hbF(Y_m), \quad (8)$$

$$Y_m = \begin{bmatrix} y_{n+1} & y_{n+\frac{3}{2}} & y_{n+2} & y_{n+\frac{5}{2}} & y_{n+3} & y_{n+4} & y_{n+5} \end{bmatrix}^T$$

$$y_n = \begin{bmatrix} y_{n-4} & y_{n-3} & y_{n-\frac{5}{2}} & y_{n-2} & y_{n-\frac{3}{2}} & y_{n-1} & y_n \end{bmatrix}^T$$

$$F(Y_m) = \begin{bmatrix} f_{n+1} & f_{n+\frac{3}{2}} & f_{n+2} & f_{n+\frac{5}{2}} & f_{n+3} & f_{n+4} & f_{n+5} \end{bmatrix}$$

$$f(y_n) = \begin{bmatrix} f_{n-4} & f_{n-3} & f_{n-\frac{5}{2}} & f_{n-2} & f_{n-\frac{3}{2}} & f_{n-1} & f_n \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, E = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, d = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 37529 \\ 0 & 0 & 0 & 0 & 0 & 0 & 151200 \\ 0 & 0 & 0 & 0 & 0 & 0 & 7093 \\ 0 & 0 & 0 & 0 & 0 & 0 & 28672 \\ 0 & 0 & 0 & 0 & 0 & 0 & 26 \\ 0 & 0 & 0 & 0 & 0 & 0 & 105 \\ 0 & 0 & 0 & 0 & 0 & 0 & 191575 \\ 0 & 0 & 0 & 0 & 0 & 0 & 774144 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1387 \\ 0 & 0 & 0 & 0 & 0 & 0 & 5600 \\ 0 & 0 & 0 & 0 & 0 & 0 & 46 \\ 0 & 0 & 0 & 0 & 0 & 0 & 189 \\ 0 & 0 & 0 & 0 & 0 & 0 & 185 \\ 0 & 0 & 0 & 0 & 0 & 0 & 675 \end{bmatrix}$$

$$b = \begin{bmatrix} 74437 & -4096 & 2713 & -14864 & 15011 & -2171 & 27 \\ 30240 & 945 & 560 & 4725 & 15120 & 30240 & 5600 \\ 378927 & -623 & 333261 & -3411 & 69151 & -2007 & 135 \\ 143360 & 160 & 71680 & 1120 & 71680 & 28672 & 28672 \\ 4969 & -128 & 523 & -2944 & 103 & -67 & 1 \\ 1890 & 35 & 105 & 945 & 105 & 945 & 210 \\ 2040625 & -22375 & 75625 & -17245 & 370625 & -54125 & 135 \\ 774144 & 6048 & 14336 & 6048 & 387072 & 774144 & 28672 \\ 2943 & -128 & 2889 & -432 & 649 & -81 & 27 \\ 1120 & 35 & 560 & 175 & 560 & 1120 & 5600 \\ 2624 & -4096 & 232 & -4096 & 2624 & 46 & 0 \\ 945 & 945 & 35 & 945 & 945 & 189 & 0 \\ 10625 & 0 & -625 & 880 & -625 & 10625 & 185 \\ 6048 & & 336 & 189 & 336 & 6048 & 672 \end{bmatrix}$$

54

55

56 2.1 Analysis of Basic Properties of the Self-Starting Block Method

57 2.1.1 Order of the Self-Starting Block Method

58 Let the linear operator $L\{y(x); h\}$ associated with the block method (8) be defined as

$$59 \quad L\{y(x); h\} = A^{(0)}Y_m - Ey_n - h(df(y_n) + bF(Y_m)) \quad (9)$$

60 Expanding using Taylor's series and comparing the coefficients of h gives

$$61 \quad L\{y(x); h\} = c_0 y(x) + c_1 h y'(x) + c_2 h^2 y''(x) + \dots + c_p h^p y^{(p)}(x) + c_{p+1} h^{p+1} y^{(p+1)}(x) + \dots \quad (10)$$

62 **Definition 2.** The linear operator L and the associated continuous linear multistep method (9) are said to be of
63 order p if $c_0 = c_1 = c_2 = \dots = c_p = 0$ and $c_{p+1} \neq 0$ is called the error constant and the truncation error is
64 given by

$$65 \quad t_{n+k} = c_{p+1} h^{p+1} y^{(p+1)}(x_n) + O(h^{p+2}) \quad (11)$$

66 For our method

$$67 \quad L\{y(x); h\} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_{n+1} \\ y_{n+\frac{3}{2}} \\ y_{n+2} \\ y_{n+\frac{5}{2}} \\ y_{n+3} \\ y_{n+4} \\ y_{n+5} \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_{n-4} \\ y_{n-3} \\ y_{n-\frac{5}{2}} \\ y_{n-2} \\ y_{n-\frac{3}{2}} \\ y_{n-1} \\ y_n \end{bmatrix} \\
- h \begin{bmatrix} \frac{37529}{151200} & \frac{74437}{30240} & -\frac{4096}{945} & \frac{2713}{560} & -\frac{14864}{4725} & \frac{15011}{15120} & -\frac{2171}{30242} & \frac{27}{5600} \\ \frac{7093}{378927} & \frac{30240}{378927} & -\frac{623}{945} & \frac{333261}{560} & -\frac{3411}{4725} & \frac{69151}{15120} & -\frac{2007}{30242} & \frac{135}{5600} \\ \frac{28672}{26} & \frac{143360}{4969} & \frac{160}{-128} & \frac{71680}{523} & \frac{1120}{-2944} & \frac{71680}{103} & \frac{28672}{-67} & \frac{28672}{1} \\ \frac{105}{191575} & \frac{1890}{2040625} & \frac{35}{-22375} & \frac{105}{75625} & \frac{945}{-17245} & \frac{105}{370625} & \frac{945}{-54125} & \frac{210}{135} \\ \frac{774144}{1387} & \frac{774144}{2943} & \frac{6048}{-128} & \frac{14336}{2889} & \frac{6048}{-432} & \frac{387072}{649} & \frac{774144}{-81} & \frac{28672}{27} \\ \frac{5600}{46} & \frac{1120}{2624} & \frac{35}{-4096} & \frac{560}{232} & \frac{175}{-4096} & \frac{560}{2624} & \frac{1120}{46} & \frac{5600}{0} \\ \frac{189}{185} & \frac{945}{10625} & \frac{945}{0} & \frac{35}{-625} & \frac{945}{880} & \frac{945}{-625} & \frac{189}{10625} & \frac{185}{185} \\ \frac{672}{6048} & \frac{6048}{6048} & 0 & \frac{336}{336} & \frac{189}{189} & \frac{336}{336} & \frac{6048}{6048} & \frac{672}{672} \end{bmatrix} \begin{bmatrix} f_n \\ f_{n+1} \\ f_{n+\frac{3}{2}} \\ f_{n+2} \\ f_{n+\frac{5}{2}} \\ f_{n+3} \\ f_{n+4} \\ f_{n+5} \end{bmatrix}$$

68

69 Expanding in Taylor's series expansion gives and equating coefficients of the Taylor's series expansion to zero yield
70 a constant order eight with the following error constants

$$c_0 = c_1 = c_2 = \dots = c_8 = 0$$

$$c_9 = [-5.814(-04), 5.703(-04), 5.748(-04), -5.710(-04), -5.776(-04), -4.056(-04), -2.045(-03)]^T$$

2.2 Zero Stability

Definition 3. The block integrator (8) is said to be zero stable, if the roots $z_s, s = 1, 2, \dots, k$ of the first characteristic polynomial $\rho(z)$ defined by $\rho(z) \det(zA^{(0)} - E)$ satisfies $|z| \leq 1$ and every root satisfying $|z| \leq 1$ have multiplicity not exceeding the order of the differential equation. Moreover, as $h \rightarrow 0, \rho(z) = z^{r-\mu} (z-1)^\mu$ where μ is the order of the differential equation, r is the order of the matrix $A^{(0)}$ and E [11].

for our method

$$\rho(z) = z \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} = 0$$

$$\rho(z) = z^6(z-1) = 0 \Rightarrow z_1 = z_2 = z_3 = z_4 = z_5 = z_6 = 0, z_7 = 1.$$

Hence, the self-starting block integrator is zero-stable.

2.3 consistency

The block integrator (8) is consistent since it has order $\rho = 8 \geq 1$

2.4 convergence

The self-starting block integrator is convergent by consequence of Dahlquist theorem below.

Theorem 4. The necessary and sufficient conditions that a continuous LMM be convergent are that it be consistent and zero-stable. [12].

2.5 Region of Absolute Stability

Definition 5. Region of absolute stability is a region in the complex z plane, where $z = \lambda h$. It is defined as those values of z such that the numerical solutions of $y' = \lambda y$, satisfy $y_j \rightarrow 0$ as $j \rightarrow \infty$ for any initial condition.

The stability polynomial for our method is given by

$$\begin{aligned}
&= -h^7 \left(\left(\frac{5}{448} \right) w^7 + \left(\frac{5}{192} \right) w^6 \right) + h^6 \left(\left(\frac{67}{1896} \right) w^7 - \left(\frac{1247}{8064} \right) w^6 \right) - h^5 \left(\left(\frac{1177}{3840} \right) w^7 + \left(\frac{711}{1280} \right) w^6 \right) \\
&- h^4 \left(\left(\frac{451}{320} \right) w^6 - \left(\frac{853}{960} \right) w^7 \right) - h^3 \left(\left(\frac{355}{192} \right) w^7 + \left(\frac{165}{64} \right) w^6 \right) + h^2 \left(\left(\frac{85}{32} \right) w^7 - \left(\frac{105}{32} \right) w^6 \right) \\
&- h \left(\left(\frac{19}{8} \right) w^7 + \left(\frac{21}{8} \right) w^6 \right) + w^7 - w^6
\end{aligned}$$

This gives the stability region shown in Figure 2.5.1 shown below.




Figure 2.5.1

According to Fatunla, stiff algorithms have unbounded RAS, also Lambert showed that the stability region for L-stable schemes must encroach into the positive half of the complex plane.

3. Numerical Examples

We shall evaluate the performance of the self-starting block method on some challenging stiff problems and the results are displayed below.

The following notations are used in the tables below.

ERS - Error in Skwame *et al.* (2012)

ERM - Error in Mohammed and Yahaya (2012)

Problem 1. Consider the mildly stiff initial value problem

$$y' = -y, y(0) = 1, h = 0.1$$

With exact solution: $y(x) = e^{-x}$

This problem was solved by [8], by adopting fully implicit four points block backward formula.

Table 1. showing the result for stiff problem 1.

x	Exact Result	Computed Result	Error in our method	ERM
0.01	0.9048374180359595	0.9048374180555555	1.959599 (-011)	2.5292(-06)
0.02	0.8187307530779818	0.8187307555555556	2.477574 (-009)	2.0937(-06)
0.03	0.7408182206817179	0.7408182624999998	4.181828 (-008)	2.0079(-06)
0.04	0.6703200460356393	0.6703203555555557	3.095199 (-007)	1.6198(-06)
0.05	0.6065306597126334	0.6065321180555554	1.458543 (-006)	3.1608(-06)
0.06	0.5488116360940264	0.5488129556691562	1.319575 (-006)	2.7294(-06)
0.07	0.4965853037914095	0.4965864992843365	1.197493 (-006)	2.5457(-06)
0.08	0.4493289641172216	0.4493300698483614	1.105731 (-006)	2.1713(-06)
0.09	0.4065696597405992	0.4065708250308643	1.165290 (-006)	3.1008(-06)
0.10	0.3678794411714423	0.3678812102329587	1.769062 (-006)	2.7182(-06)

Problem 2. Consider the highly stiff initial value problem,

$$y' = -\lambda y, y(0) = 1, h = 0.01, \lambda = 10$$

With exact solution: $y(x) = e^{-\lambda x}$

[9] solved this problem. by adopting an L-Stable hybrid block Simpson's method of order six.

Table 2. showing the result for stiff problem 2.

x	Exact Result	Computed Result	Error in our method	ERS
0.01	0.9048374180359595	0.9048374180555555	1.959599 (-011)	6.28(-03)
0.02	0.8187307530779818	0.8187307555555556	2.477574 (-009)	1.88(-03)
0.03	0.7408182206817179	0.7408182624999998	4.181828 (-008)	3.26(-03)
0.04	0.6703200460356393	0.6703203555555557	3.095199 (-007)	1.06(-02)
0.05	0.6065306597126334	0.6065321180555554	1.458543 (-006)	3.85(-03)
0.06	0.5488116360940264	0.5488129556691562	1.319575 (-006)	1.45(-03)
0.07	0.4965853037914095	0.4965864992843365	1.197493 (-006)	5.02(-04)
0.08	0.4493289641172216	0.4493300698483614	1.105731 (-006)	2.76(-04)
0.09	0.4065696597405992	0.4065708250308643	1.165290 (-006)	1.01(-04)
0.10	0.3678794411714423	0.3678812102329587	1.769062 (-006)	3.74(-05)

3.1 Discussion of the Result

We consider two numerical examples in this paper. Mohammed, U., and Yahaya, Y. A., (2010) solved the first problem (which is a mildly stiff), where they proposed a fully implicit four points block backward difference formula. Our method gave a better approximation because we proposed a self-starting method, which does not, required a starting value. Skwame, Y., Sunday, J., and Ibijola, E. A., (2012) solved the second problem.

They adopted an L-Stable hybrid block Simpson's method of order six. Our method gave a better approximation, because the iteration per step in the new method is lower than the proposed method by Skwame, Y., Sunday, J., and Ibijola, E. A., (2012).

4. Conclusion

In this paper, we have presented a self-starting five-step eight-order block method for the solution of first order ordinary differential equations. The approximate solution adopted in this research produced a block method with L-stable stability region. This made it to perform well on stiff problems. The block method proposed was found to be zero-stable, consistent and convergent. The new block method was also found to perform better than the existing methods.

REFERENCES

136 [1] Lambert, J. D., Computational Methods in Ordinary Differential Equations. John Willey and Sons, New York,
137 1973

138 [2] Lie, I. and Norsett, P., (1989): Super convergences for the multi-step collocation.

139 [3] Onumanyi, *et al.* (1994): New linear multistep methods with continuous coefficients for the first order ordinary
140 initial value problems. *Nig. J. Math. Soc.* 13: 37-51.

141 [4] Onumanyi, *et al.* (1999): Continuous finite differential approximation for solving differential equations.
142 *Int. J. Comp. Math.* 72: 15-27
143

144 [5] Butcher, J. C. (1993): General linear method for the parallel solution of differential equations.
145 *App. Anal.* 2: 99-111.

146 [6] James, A. A., Adesanya, A. O. and Sunday, J. (2012): Uniform order continuous block hybrid method for the
147 solution of first order ordinary differential equations. "IOSR J. Math."

148 [7] Sunday, J., Adesanya, A. O., and Odekunle, M. R.: A self-starting four-step fifth-order block integrator for and
149 oscillatory differential equations. *J. Math. Comput. Sci.* 4 (2014), No. 1, 73-84

150 [8] Mohammed, U., and Yahaya, Y. A., (2010): Fully implicit four points block backward difference formula for
151 solving first-order IVP. "Leonado J. Sci." 16(1), 21-30.

152 [9] Skwame, Y., Sunday, J., and Ibijola, E. A., (2012): L-stable block hybrid Simpson's methods for Numerical
153 solution of IVP in stiff ODE's." *Int. J. Pure. Appl. Sci. Technol.* 11(2), 45-54.

154 [10] Donald, J. Z. *et al.* (2009): Construction of two steps Simpson's multistep method as parallel integrator for the
155 solution of ordinary differential equation. "Bagale J. Pure Appl. Sci." 7:1-12.

156 [11] Awoyemi, D. O. , Ademiluyi, R. A. and Amusegham, (2007): "off-grid points exploitation in the development
157 of more accurate collocation method for solution of ODE's" . *J. N. A. M. P.* 12, 379-386

158 [12] Dahlquist, G. G. Convergence and stability in the numerical integration of ordinary differential equations,
159 *Math. Scand.* 4 (1956), 33-50.

160