

Relations between various magnitudes which serve as signs of a right-angled triangle

Moshe Stupel

Shaanan & Gordon - Academic College of Education, Haifa, Israel

Avi Sigler

Shaanan- Academic College of Education, Haifa

Abstract

We present a proven scheme of 15 relations between the sides of a triangle, the radii of incircles and excircles of the triangle, the area and half the perimeter of the triangle, which serve as signs of a right-angled triangle, together with the 16th relation which is the **Pythagorean Theorem**. The relations are investigated dynamically using the computerized software.

Keywords


Relations between magnitudes in a right-angled triangle, dynamic investigation in a right-angled triangle.

Introduction

Relations between different magnitudes in a triangle, such as the radius of the circumcircle of the triangle, the radius of the incircle of the triangle, the radii of the excircles the triangle (which are tangent to one side from the outside and the continuations of the two other sides), as well as their relations with the side lengths the triangle and certain segments in the triangle, are a fascinating subject that can be investigated by hypothesizing and investigating the hypotheses dynamically using a computerized technological tool, or using formal mathematical proofs employing various mathematical tools and algebraic manipulations.

In connection with the excircles, the incircle, and the circumcircle, one should mention **the nine-point circle** [1-2] discovered by **Leonard Euler** in 1765. The nine points in the triangle through which the circle passes:

- The middles of the sides;
- The feet of the altitudes – the points at which altitudes meet the sides;
- The middles of the segments that connect the vertices of the triangle and the point of intersection of the altitudes.



The nine point circle for an arbitrary triangle $\triangle ABC$ is shown in Fig. 1.

Fig. 1.
The 9 point circle

Later, in 1822, Karl Wilhelm Feuerbach showed that this circle is also tangent to the excircles of the triangle [2-3]. This led to increased interest in finding the relations between the radii and various segments.

It is important to note that the center of the nine-point circle lies on “Euler’s line”, which passes through the point of intersection of the medians, the point of intersection of the altitudes and the point of intersection of the mid-perpendiculars. The length of the radius of the nine-point circle is equal to half the length of the radius of the circumcircle of the triangle $\triangle ABC$.

Several relations are known between the radii, one of which is $\frac{1}{r} = \frac{1}{R_a} + \frac{1}{R_b} + \frac{1}{R_c}$ [4].

Although the geometric properties and some of the connections have been known for many years, the issue is still relevant and appears in some publications[5-8]. Interesting connections exist in a right-angled triangle in the context of Pythagorean Theorem and its various representations.

The current activities are focused on finding 15 relations between various magnitudes that constitute signs for the existence of a right-angled triangle which supplement the well-known relation of the Pythagorean Theorem. Any relation is a necessary and sufficient condition for the triangle to be right-angled. Each relation is accompanied by a mathematical proof based on several known relations between the magnitudes in the triangle and its excircles, incircle and well-known geometrical properties.

Development of the relations that constitute signs for a right-angled triangle

Notation:

$p = \frac{a+b+c}{2}$ is half the perimeter of the triangle $\triangle ABC$

r is the radius of the incircle of the triangle $\triangle ABC$

R_a is the radius of the excircle of the triangle and is tangent to the side a .

R_b is the radius of the excircle of the triangle and is tangent to the side b .

R_c is the radius of the excircle of the triangle and is tangent to the side c .

R is the radius of the circumcircle of the triangle $\triangle ABC$

The triangle and the radii of the different circles are shown in Fig. 2.




Fig. 2

The triangle and its incircle, excircles and circumcircle

A GeoGebra applet was prepared which allows the vertices of the triangle to be dragged, and thus the lengths of its sides changed, resulting in an immediate demonstration of the effect on the lengths of the radii, which appear on screen at every stage.

It is possible to reach the applet using the following link:

Link 1: <https://www.geogebra.org/m/TueeM84U>

Known relations:

$$r = \frac{S_{\triangle}}{p}$$

$$R_a = \frac{S_{\triangle}}{p-a}$$

$$R_b = \frac{S_{\triangle}}{p-b}$$

$$R_c = \frac{S_{\triangle}}{p-c}$$

$$a + b - c = 2(p - c) , a + c - b = 2(p - b) , b + c - a = 2(p - a)$$

The right-angled triangle and its circles, together with the scheme of the relations between the magnitudes that constitute a precondition for the formation of such a triangle. In Fig. 3.




Fig. 3

A right-angled triangle, its circles and the scheme of the relations between its magnitudes

Mathematical development of the relations between the magnitudes in a right-angled triangle

$$(1) S = \frac{a \cdot b}{2} \Rightarrow a \cdot b = 2S$$

$$(2) a^2 + b^2 = c^2 \Rightarrow a^2 + b^2 - c^2 = 0 \Rightarrow \underbrace{a^2 + b^2 - c^2 + 2a \cdot b = 2a \cdot b}_{\Downarrow} = 4S$$

$$(3) (a + b)^2 - c^2 = 4S \Rightarrow \underbrace{(a + b + c) \cdot (a + b - c) = 4p \cdot (p - c)}_{\Downarrow} = 4S$$

$$(4) (a + b)^2 - c^2 = 4S \Rightarrow S = p \cdot (p - c)$$

$$(5) S = \sqrt{p(p-a)(p-b)(p-c)} = p(p-c) \Rightarrow (p-a) \cdot (p-b) = p(p-c) = S$$

$$(6) r = \frac{S}{p} = \frac{p(p-c)}{p} = p - c$$

$$(7) R_a = \frac{S}{p-a} = \frac{(p-a) \cdot (p-b)}{p-a} = p - b$$

$$(8) R_b = \frac{S}{p-b} = \frac{(p-a) \cdot (p-b)}{p-b} = p - a$$

$$(9) R_c = \frac{S}{p-c} = \frac{p \cdot (p-c)}{p-c} = p$$

$$(10) R_a + R_b + R_c = p - b + p - a + c = 2R \Rightarrow R = \frac{R_a + R_b}{2}$$

$$(11) R + r = \frac{a+b}{2}$$

looking at Fig. 4, one can write down the following relations:




Fig. 4

The relation between the radius of the incircle, the radius of the circumcircle and the legs of the triangle

$$BF = BD = a - r$$

$$AE = AF = b - r$$

$$AB = c = 2R = BF + AF = a - r + b - r \Rightarrow 2R + 2r = a + b$$

$\Downarrow$

$$R + r = \frac{a+b}{2}$$

$$(12) \quad r + R_a = p - c + p - b = a \quad (6, 7)$$

$$(13) \quad r + R_b = p - c + p - a = b \quad (6, 8)$$



$$(14) \quad R_c - r = p - (p - c) = c \quad (6, 9)$$

$$(15) \quad r + R_a + R_b = p - c + p - b + p - a = p = R_c \quad (6, 7, 8)$$

$$(16) \quad r \cdot R_c = (p - c) \cdot p = \frac{a \cdot b}{2} \quad (3, 6, 9)$$

$$(17) \quad R_a \cdot R_b = (p - b) \cdot (p - a) = \frac{a \cdot b}{2} \quad (7, 8)$$

Numerical example



A GeoGebra applet was prepared in order to illustrate the formulas, which contains a right-angled triangle with a fixed vertex C and vertices A and B that can be dragged, changing the lengths of the sides and accordingly – the radii of the circles shown on the screen. In each stage, the screen shows the values of the radii, the side lengths of the triangle, the area, and half the perimeter.

One can reach the applet using Link 2:

Link 2: <https://www.geogebra.org/m/p3k9KJzg>

Note: the proof of the relations between various magnitudes that constitute signs for the existence of a right-angled triangle was based on the fact that the triangle is given.

However, in order to prove that these relations are necessary and sufficient conditions one must also provide the proof in the other direction, where the relation is given and one has to prove that it necessarily gives a right-angled triangle. In some of the relations the proof is trivial and in others, a mathematical effort is required in order to obtain the proof. A check was made that this is true, and the reader should confirm this.

Summary

We presented a scheme of relations between the side lengths of a triangle, the radii of its excircles and incircle, the circumcircle and half the perimeter of the triangle. When one of the relations holds in itself, this means that the triangle is right-angled.

Bibliography

- [1]. Honsberger, R.(1995). "The Nine-Point Circle." , pp. 6-7 in Episodes in
Nineteenth and Twentieth Century Euclidean Geometry, by The Mathematical
Association of America, Washington, DC.
- [2]. Johnson, R.A.(2007). Advanced Euclidean Geometry, pp. 195-200. Dover
Publications, Mineola, New York.
- [3]. Pedoe , D. (1988). Geometry: A Comprehensive Course. pp. 56-63. In the Dover
Books on Mathematics Ed.
- [4]. Coxeter, H. S. M. and Greitzer, S. L.(1967). "The Nine-Point Circle.", pp. 20-22.
In the " Geometry Revisited ", by The Mathematical Association of America, New
York.
- [5]. Altshiller-Court, N. (1967). College Geometry An. Introduction to the Modern
Geometry of the Triangle and the Circle. Revised and Enlarged Dover
Publications, Inc. Mineola, New York. pp. 72-88, 115-119.
- [6]. D. W. Hansen, D. W. (2003). On inscribed and escribed circles of right triangles,
circumscribed triangles, and the four-square, three-square problem, Mathematics
Teacher, 96, pp 358-364
- [7]. Alperin, R.C. (2005). The modular tree of Pythagoras. American Mathematical
Monthly, 112(9), pp. 807–816.
- [8]. Mack. J. and Czernezkyj, V.(2010). The tree in Pythagoras' garden. Australian
Senior Mathematics Journal. Vol. 24(2), pp. 58-63.

College Geometry An Introduction to the Modern Geometry of the Triangle and the Circle Nathan Altshiller-Court Second Edition Revised and Enlarged Dover Publications, Inc. Mineola, New York