

Solving different order boundary value problems by Applying Homotopy Type Methods.

Muhammad Naeem¹, Zahid Khan², Sher Muhammad^{1,3} and Liaqat Khan^{3,4}

(muhammadnaee@yahoo.com; zahidkhalil90@gmail.com ; shermuhammad@cecos.edu.pk,
liaqatislamian@gmail.com)

¹Department of Basic Sciences, CECOS University of IT and Emerging Sciences, Peshawar, Pakistan;

²Department of Mathematics, Peshawar Model Degree College Hayatabad, Peshawar, 25000, KP, Pakistan;

³Department of Mathematics, Islamia College Peshawar 25000, KP, Pakistan;

⁴Department of Mathematics, Malala Yousafzai Danish School Jand, Attock, 43100, Pakistan;

AUTHORS' CONTRIBUTIONS:

This work was carried out in equal collaboration of all the authors. The final version of the manuscript has been read and approved by the authors.

*Corresponding author Email: shermuhammad@cecos.edu.pk

ABSTRACT

In this manuscript, the methods, Optimal Homotopy Asymptotic Method (OHAM), Homotopy Perturbation Method (HPM) and Homotopy Analysis Method (HAM) are applied to solve various order boundary value problems. These techniques give solutions in the form of a series. Their solutions and graphs are compared with their exact solutions and graphs. All these give profitable results, but the Optimal Homotopy Asymptotic Method is more precise, whose convergence is restrained optimizations and the convergence area can be accustomed according to the problem concerned. The results show that the suggested scheme is more active and relaxed to routine.

Keywords: OHAM, HPM, HAM, various order boundary value problems.

1. INTRODUCTION

Differential equations can be used to model various types of physical structures such as sociological, economical, biological and chemical, etc. Also in literature, physical problems are investigated by differential equations, which are mostly handled by the common methods, Splines (S) [10], Homotopy perturbation method (HPM) [21], Homotopy Analysis Method (HAM) [22, 23]. The non-perturbed techniques (DTM) [24], and ADM) [13, 16] concern nonlinear problems, but the region of convergence of their series solution is generally small. Freshly Herisanu and Marinca et al. [3-5] familiarized OHAM for approximate solution of nonlinear problems of shrill film flow of a fourth grade fluid down an upright cylinder. They used the

whispered method for understanding the conduct of nonlinear mechanical vibration of an electric contraption. By using this method they investigated the solution of nonlinear equations ascending in the study of state-run flow of a fourth grade fluid past a porous plate. This method supplies the need to switch the convergence. In general the OHAM solution settles with the exact solution. The graphs of the two solutions are conterminous. We have smeared OHAM to various types of boundary value problems and have investigated that the original exact solution approves with the numerical solution, the error noted is small. This method is operational and is relaxed to routine. The other relevant work [24-30] have been observed in these articles.

2.0: Analysis of the method OHAM.

Considering the boundary problem of the arrangement

$$L(F(y)) + g(y) + N(F(y)) = 0, B\left(F, \frac{dF}{dy}\right) = 0 \quad (2.0 a)$$

Where L is taken as linear operator y is independent variable $F(y)$ is an unknown function $g(y)$ is a notorious function, $N(F(y))$ is a nonlinear operator and B is a boundary operator. According to the idea of OHAM we construct a homotopy as given below

$$H(\theta(y, p), p) : R \times [0, 1] \rightarrow R$$

That satisfies the homotopy functional as given below

$$(1-p)[L(\theta(y, p)) + g(y)] = h(p)[L(\theta(y, p)) + g(y) + N(\theta(y, p))], B\left(\theta(y, p), \frac{\partial(\theta(y, p))}{\partial y}\right) = 0, \quad (2.0 b)$$

Where, $y \in R$, $p \in [0, 1]$ is an embedding parameter, $h(p)$ is a nonzero auxiliary function for $p \neq 0$, $h(0)=0$ and $\theta(y, p)$ is an Unknown function. Evidently, for $p=0$ and $p=1$. Manifestly, for $p=0$ and $p=1$ it restrains that $\theta(y, 1) = F(y)$ respectively. Thus as p varies from 0 to 1, the solution $\theta(y, p)$ approaches from $F_0(y)$ to $F_1(y)$ where $F_0(y)$ is obtained from Equation (2.0 b) for $p = 0$ and we have.

$$L(F_0(y)) + g(y) = 0, B\left(F_0, \frac{dF_0}{dy}\right) = 0, \quad (2.0 c)$$

Now choosing auxiliary function $h(p)$ in the following pattern

$$h(p) = pC_1 + p^2C_2 + \dots, \quad (2.0 d)$$

Where $C_1, C_2, \dots$ are constants to be determined later. $h(p)$ can be uttered in many customs as investigated by V. Marinca [3-5]. To acquire an approximate solution one can expand $\theta(y, p, C_i)$ in Taylor's series about p in the following Pattern.

$$\theta(y, p, C_i) = F_0(y) + \sum_{k=1}^{\infty} F_k(y, C_1, C_2, \dots, C_k) p^k, \quad (2.0 e)$$

Making use of equation (2.0 e) in equation (2.0 b) and comparing the coefficients of resembling powers of p we have the following linear equations zeroth order problem is given by equation (2.0 c) and the first and second order problems are given by equations (2.0 f) and (2.0 g)

$$L(F_1(y)) + g(y) = C_1 N_0(F_0(y)), B\left(F_1, \frac{dF_1}{dy}\right) = 0, \quad (2.0 f)$$

$$L(F_2(y)) - L(F_1(y)) = C_2 N_0(F_0(y)) + C_1 [L(F_1(y)) + N_1(F_0(y), F_1(y))], B\left(F_2, \frac{dF_2}{dy}\right) = 0, \quad (2.0 g)$$

The general governing equations for $F_k(y)$ are given by

$$L(F_k(y)) - L(F_{k-1}(y)) = C_k N_0(F_0(y)) + \sum_{i=1}^{k-1} C_i [L(F_{k-i}(y)) + N_{k-i}(F_0(y), F_0(y), \dots, F_{k-1}(y))], k = 2, 3, \dots, B\left(F_k, \frac{dF_k}{dy}\right) = 0, \quad (2.0 h)$$

Where $N_m(F_0(y), F_1(y), \dots, F_{k-1}(y))$ is the coefficient of p^m in the expansion of $N(\theta(y, p))$ about the embedding Parameter p

$$N(\theta(y, p, C_i)) = N_0(F_0(y)) + \sum_{m=1}^{\infty} N_m(F_0, F_1, \dots, F_m) p^m, \quad (2.0 i)$$

It has been investigated that the convergence of the series (2.0 e) depends on the auxiliary constants $C_1, C_2, \dots$. If it is convergent. At $p = 1$ then we have

$$\theta(y, C_i) = F_0(y) + \sum_{k=1}^{\infty} F_k(y, C_1, C_2, \dots, C_k), \quad (2.0 j)$$

The upshot of the m th order approximations are

$$\tilde{F}(y, C_1, C_2, \dots, C_m) = F_0(y) + \sum_{i=1}^m F_i(y, C_1, C_2, \dots, C_k), \quad (2.0 k)$$

Using equation (2.0 k) into equation (2.0 a) we get the residual

$$R(y, C_1, C_2, \dots, C_m) = L(\tilde{F}(y, C_1, C_2, \dots, C_m)) + g(y) + N(\tilde{F}(y, C_1, C_2, \dots, C_m)), \quad (2.0 l)$$

If $R=0$ then $\tilde{F}$ be the exact solution. Generally it does not happen especially in nonlinear problems .In order to get the Optimal Values of C_i 's $i=1,2,3...$ we first construct the Functional Values of C_i 's $i=1,2,3...$ we first construct the Functional.

$$J(C_1, C_2, \dots, C_m) = \int_a^b R^2(y, C_1, C_2, \dots, C_m) dy, \quad (2.0 m)$$

And then minimizing it using the basic calculus we get

$$\frac{\partial J}{\partial C_1} = 0, \frac{\partial J}{\partial C_2} = 0, \dots, \frac{\partial J}{\partial C_m} = 0. \quad (2.0 n)$$

Knowing the values of $C_1, C_2 \dots C_m$. The approximate solution of order m is determined. Where a, b lie in domain of the concerned problem using the least square method we get OHAM solution.

2.1 Fundamentals of HPM:

We show the basic idea of the homotopy perturbation method [21] and considering the nonlinear differential equation of the form:

$$C(z) - g(s) = 0, s \in \Omega \quad (2.1 a)$$

With boundary conditions

$$G\left(z, \frac{dz}{dn}\right) = 0, s \in \Gamma \quad (2.1 b)$$

Where C is a general differential operator, G is a boundary operator, z is a known analytical function,

Γ is the boundary of the domain Ω . The operator C can be divided into two parts, E and W

are linear and nonlinear . Equation (2.1a) takes the form:

$$E(z) + W(z) - g(s) = 0, \quad (2.1 c)$$

By the homotopy method proposed by Liao [23]. A homotopy can be constructed as

$v(s, j) : \Omega \times [0,1] \rightarrow R$ This satisfies:

$$T(v, j) = (1 - j)[E(v) - E(z_0)] + j[C(v) - g(s)] = 0, \quad (2.1 d)$$

$$\text{or } T(v, j) = E(v) - E(z_0) + jE(z_0) + j[W(v) - g(s)] = 0, \quad (2.1 e)$$

Where $s \in \Gamma$ and $j \in [0,1]$ is an embedding parameter, z_0 is an initial approximation of (22) which satisfies the boundary conditions using (2.1 d) we can easily guess that

$$T(v, 0) = E(v) - E(z_0) = 0, \quad (2.1 f)$$

$$T(v, 1) = C(v) - g(s) = 0, \quad (2.1 g)$$

And the altering technique of j from zero to one is just that of $T(v, j)$ from $E(v) - E(z_0)$ to $C(v) - g(s)$. In the Topology, this is called deformation, where $C(v) - g(s)$ and $E(v) - E(z_0)$ are called homotopic. The embedding parameter j is known initially. For $0 < j \leq 1$ equation (4) can be given as

$$v = v_0 + jv_1 + j^2v_2 + \dots, \quad (2.1 \text{ h})$$

The approximate solution of equation of (18) can be obtained as follows:

$$z = \lim_{j \rightarrow 1} v = v_0 + v_1 + v_2 + \dots, \quad (2.1 \text{ i})$$

2.2: Fundamentals of HAM.

We start by considering

$$N[u(x)] = 0, \quad (2.2 \text{ a})$$

Where N is a nonlinear operator $u(x)$ is an unknown function and x denotes independent variable. For simplicity we

Ignore all boundary conditions. We construct zeroth order functional as:

$$(1 - q)L[\phi(x, q) - u_0(x)] = qhH(x)N[\phi(x, q)], \quad (2.2 \text{ b})$$

Where $q \in [0, 1]$ is an embedding parameter, $h \neq 0$ is a nonzero auxiliary parameter $H(x)$ is an auxiliary function L

Is a linear operator $u_0(x)$ is an initial guess of $u(x)$, $\phi(x, q)$ is an unknown function, from equation (2.2b) $q = 0$ and $q = 1$ we get

$$\phi(x, 0) \text{ and } \phi(x, 1), \quad (2.2 \text{ c})$$

Thus as q increases from 0 to 1, the solution $\phi(x, q)$ varies from the $u_0(x)$ to $u(x)$. Expanding $\phi(x, q)$ in Taylor series

$$\text{We have} \quad \phi(x, q) = u_0(x) + \sum_{m=1}^{+\infty} u_m(x)q^m, \quad (2.2 \text{ d})$$

$$\text{Where} \quad u_m(x) = \frac{1}{m!} \frac{\partial^m \phi(x, q)}{\partial q^m}, \text{ at } q = 0 \quad (2.2 \text{ e})$$

If the linear operator, initial guess, the auxiliary parameter and auxiliary function are so properly chosen the series (2.3 e) Converges at $q = 1$ We have

$$u(x) = u_0(x) + \sum_{m=1}^{+\infty} u_m(x), \quad (2.2 \text{ f})$$

Which is one of the solution of the original non-linear equation approved by Liao [22] choosing $h = -1$ and $H(x) = 1$

Equation (2.2 b) becomes

$$(1-q)L[\phi(x, q) - u_0(x)] + qN[\phi(x, q)] = 0, \quad (2.2 \text{ g})$$

Which is used mostly in Homotopy Perturbation Method (HPM) whereby the solution is obtained directly, without using Taylor series the comparison between HPM and HAM can be found in [22]. As $H(x) = 1$ equation (2.2 b) becomes

$$(1-q)L[\phi(x, q) - u_0(x)] = qhN[\phi(x, q)], \quad (2.2 \text{ h})$$

This is often used in (HAM). In this case $H(x)$ will not be involved in setting of the base function. Differentiating equation (2.2b) m times with respect to q and the $q=0$ and finally dividing them by $m!$ we have the so called m^{th} -order deformation equation

$$L[u_m(x) - \chi_m u_{m-1}(x)] = hH(x)R_m(\bar{u}_{m-1}, x), \quad (2.2 \text{ i})$$

$$R_m(\bar{u}_{m-1}, x) = \frac{1}{(m-1)!} \frac{\partial^{m-1} N[\phi(x, q)]}{\partial q^{m-1}}, \quad \text{at } q = 0 \quad (2.2 \text{ j})$$

$$\bar{u}_m = \{u_0(x), u_1(x), \dots, u_m(x)\}, \quad (2.2 \text{ k})$$

And $\chi = \{0, m \leq 1, \chi = \{1, m > 1\}$ putting equation (2.2 d) in (2.2 j) we have

$$R_m(\bar{u}_{m-1}, x) = \frac{1}{(m-1)!} \frac{\partial^{m-1}}{\partial q^{m-1}} N \left[\sum_{n=0}^{+\infty} u_n(x) q^n \right] \cdot \text{at } q = 0 \quad (2.2 \text{ l})$$

It should be restrained that $u_m(x)$ for $m \geq 1$ is governed by the linear equation (2.2 i) with the linear boundary condition that is obtained from the original problem which can be solved by Matlab.

3. Numerical problems

Problem3.1.

Consider the following linear differential equation

$$F'(y) - F(y) - y \cos y + y \sin y - \sin y = 0, F(0) = 0, \quad (3.1 \text{ a})$$

The exact solution of the problem is

$$F(y) = y \sin y, \quad (3.1 \text{ b})$$

Applying the method mentioned above, the zeroth order problem is

$$F'_0(y) = 0, F_0(0) = 0, \quad (3.1 \text{ c})$$

Its solution is

$$F_0(y) = 0, \quad (3.1 \text{ d})$$

First order problem is

$$F_1'(y, C_1) = -y \cos y C_1 - \sin y C_1 + y \sin y C_1 - C_1 F_0(y) + (1 + C_1) F_0'(y), F_1(0) = 0, \quad (3.1 e)$$

Its solution is

$$F_1(y, C_1) = -y \cos y C_1 + \sin y C_1 - y \sin y C_1, \quad (3.1 f)$$

Second order problem is

$$F_2'(y, C_1, C_2) = \begin{cases} -y \cos y C_2 - \sin y C_2 + y \sin y C_2 - C_2 F_0(y) - C_1 F_1(y) + C_2 F_0'(y) + (1 + C_1) F_1', \\ F_2(0) = 0 \end{cases}, \quad (3.1 g)$$

Its solution is

$$F_2(y, C_1, C_2) = -y \cos y C_1 + \sin y C_1 - y \sin y C_1 - 2C_1^2 + 2 \cos y C_1^2 - 2y \cos y C_1^2 + 2 \sin y C_1^2 - y \cos y C_2 + \sin y C_2 - y \sin y C_2, \quad (3.1 h)$$

Third order problem is

$$F_3'(y, C_1, C_2, C_3) = -y \cos y C_3 - \sin y C_3 + y \sin y C_3 - C_3 F_0(y) - C_2 F_1(y) - C_1 F_2(y) + C_3 F_0'(y) + C_2 F_1'(y) + (1 + C_1) F_2'(y), F_3(0), \quad (3.1 i)$$

Its solution is

$$F_3(y, C_1, C_2, C_3) = -y \cos y C_1 + \sin y C_1 - y \sin y C_1 - 4C_1^2 + 4 \cos y C_1^2 - 4y \cos y C_1^2 + 4 \sin y C_1^2 - 6C_1^3 + 2y C_1^3 + 6 \cos y C_1^3 - 2y \cos y C_1^3 + 2y \sin y C_1^3 - y \cos y C_2 + \sin y C_2 - y \sin y C_2 - 4C_1 C_2 + 4 \cos y C_1 C_2 - 4y \cos y C_1 C_2 + 4 \sin y C_1 C_2 - y \cos y C_3 + \sin y C_3 - y \sin y C_3, \quad (3.1 j)$$

Now we use equations (3.1 d), (3.1 f), (3.1 h), (3.1 j), and the third order approximate solution by OHAM for $p=1$ is

$$\tilde{F}(y, C_1, C_2, C_3) = F_0(y) + F_1(y, C_1) + F_2(y, C_1, C_2) + F_3(y, C_1, C_2, C_3). \quad (3.1 k)$$

Using the technique mentioned in section 2 on the domain $a = 0, b = 1$. we use the residual

$$R = \tilde{F}' - \tilde{F} - y \cos y + y \sin y - \sin y \quad (3.1 l)$$

The following values of C_1, C_2, C_3 are obtained

$$C_1 = -1.1763684395430054, C_2 = 0.11688800369849017, C_3 = -0.0026970142170896472, \quad (3.1 m)$$

Considering the values of C_1, C_2, C_3 , the approximate OHAM solution becomes

$$\tilde{F}(y) = 1.01231y^2 - 0.0480322y^3 - 0.105409y^4 - 0.0223286y^5 + 0.00421533y^6 + 0.00112045y^7 - 0.0000878039y^8 - 0.0000237398y^9 + 1.11482 \times 10^{-6}y^{10} + 2.90161 \times 10^{-7}y^{11} - 9.50035 \times 10^{-9}y^{12} - 2.33658 \times 10^{-9}y^{13} + 5.79949 \times 10^{-11}y^{14} + 1.3396 \times 10^{-11}y^{15} + O(y^{16}), \quad (3.1 n)$$

The HPM solution is

$$\tilde{F}(y) = y^2 - 0.333333y^3 - 0.166667y^4 + 0.033333y^5 + y0.008333y^6 - 0.001190y^7 - 0.000198y^8 + O(y^9), \quad (3.1 \text{ o})$$

The HAM solution is


$$\tilde{F}(y) = 1.0026y^2 - 0.021482y^3 - 0.12265y^4 - 0.0223774y^5 + 0.00539169y^6 + 0.00109117y^7 + O(y^8). \quad (3.1 \text{ p})$$

The following table 3.1 displays values of exact solution (3.1 b) OHAM solution (3.1 n) and the error.



y	Exact sol	OHAM sol	HPM sol	HAM sol	Er OHAM	Er HPM	Er HAM
0.0	0.000000	0.000000	0.000000	0.0000000	0.0 E-0	0.0 E-0	0.0 E-0
0.1	0.00998334	0.0100643	0.00965034	0.00999199	8.0 E-6	3.3 E-4	8.0 E-6
0.2	0.0397379	0.0399325	0.0370779	0.0397289	1.9 E-4	2.6 E-3	4.9 E-6
0.3	0.0886561	0.0889059	0.0797368	0.0886099	2.4 E-4	8.9 E-3	4.6 E-5
0.4	0.155767	0.155987	0.134773	0.155695	2.1 E-4	2.0 E-2	7.2 E-5
0.5	0.239713	0.239861	0.199079	0.239691	1.4 E-4	4.0 E-2	2.1 E-5
0.6	0.33785	0.333884	0.269344	0.338939	9.8 E-5	6.9 E-2	1.5 E-4
0.7	0.450952	0.451076	0.342124	0.451411	1.2 E-4	1.0 E-1	4.5 E-4
0.8	0.573885	0.574113	0.413894	0.574712	2.2 E-4	1.5 E-1	8.2 E-4
0.9	0.704994	0.705338	0.481116	0.706085	3.4 E-4	2.2 E-1	1.0 E-3
1.0	0.841471	0.841762	0.540302	0.842428	2.9 E-4	3.0 E-1	9.5 E-4

Table 3.1

In the Table 3.1 the values of HPM, HAM and their errors are also given. From the table given above we conclude that the errors of the technique, OHAM are smaller than the errors of HPM and HAM.



(a) Graph of exact and OHAM solutions



(b) Graph of exact and HPM solutions

(c) Graph of exact and HAM solutions

Figure 3.1

From the figure 3.1 (a) we investigate that the graphs of OHAM and exact solutions coincide while in figure 3.1 (b) the graphs of HPM and exact solutions are not coincident completely and the graphs of figure 3.1 (c) are again coinciding but generally the graphs of OHAM are in best agreement with their exact solutions. From the table 3.1 given above **we conclude** that the exact and OHAM solutions are in best agreement in the domain $a = 0, b = 1$, solid curve elucidates exact solution, dotted curve **provides** OHAM solution. We conclude that the two curves coincide.

Problem 3.2:

For $y \in [0,1]$ we consider the following differential equation

$$F''(y) + 2F'(y) + F(y) = 0, F(0) = 1, F'(0) = 0, \quad (3.2 \text{ a})$$

The exact solution of the problem is

$$F(y) = e^{-y} + ye^{-y}, \quad (3.2 \text{ b})$$

Applying the technique, OHAM our zeroth order problem is

$$F_0''(y) + 2F_0'(y) + F_0(y) = 0, F_0(0) = 1, F_0'(0) = 0, \quad (3.2 \text{ c})$$

Its solution is

$$F_0(y) = \cos y, \quad (3.2 \text{ d})$$

First order problem is

$$F_1''(y, C_1) = -F_1(y) - 2F_1'(y) + (1 + C_1)[F_0(y) + 2F_0'(y) + F_0''(y)], F_1(0) = 0, F_1'(0) = 0, \quad (3.2 \text{ e})$$

Its solution is

$$F_1(y, C_1) = \frac{1}{2} \left(2y \cos y - \sin y + \cos 2y \sin y - \cos y \sin 2y + 2y \cos y C_1 - \sin y C_1 \right), \quad (3.2 \text{ f})$$

Second order problem is

$$F_2''(y, C_1, C_2) = \begin{pmatrix} C_2 F_0(y) - F_2(y) + 2C_2 F_0'(y) - 2F_2'(y) + C_2 F_0''(y) + (1 + C_1)[F_1(y) + 2F_1'(y) + F_1''(y)] \\ F_2(0) = 0, F_2'(0) = 0 \end{pmatrix}, \quad (3.2 \text{ g})$$

Its solution is

$$F_2(y, C_1, C_2) =$$

$$\frac{1}{4} \left(\begin{aligned} &\cos y + 4y \cos y + 2y^2 \cos y - \cos y \cos 2y - 2 \sin y + 2 \cos 2y \sin y + 2y \cos 2y \sin y - \\ &2 \cos y \sin 2y - 2y \cos y \sin 2y - \sin y \sin 2y + 2 \cos y C_1 + 8y \cos y C_1 + 4y^2 \cos y C_1 - \\ &2 \cos y \cos 2y C_1 - 4 \sin y C_1 + 4 \cos 2y \sin y C_1 + 4y \cos 2y \sin y C_1 - 4 \cos y \sin 2y C_1 - \\ &4y \cos y \sin 2y C_1 - 2 \sin y \sin 2y C_1 + \cos y C_1^2 + 4y \cos y C_1^2 + 2y^2 \cos y C_1^2 - \\ &\cos y \cos 2y C_1^2 - 2 \sin y C_1^2 + 2 \cos 2y \sin y C_1^2 + 2y \cos 2y \sin y C_1^2 - 2 \cos y \sin 2y C_1^2 \\ &- 2y \cos y \sin 2y C_1^2 - \sin y \sin 2y C_1^2 + 4y \cos y C_2 - 2 \sin y C_2 + 2 \cos 2y \sin y C_2 - \\ &2 \cos y \sin 2y C_2 \end{aligned} \right), \quad (3.2 h)$$

Third order problem is

$$F_0''(y, C_1, C_2, C_3) = \left(\begin{aligned} &C_3 F_0(y) + C_2 F_1(y) - F_3(y) + 2C_3 F_0'(y) + 2C_2 F_1'(y) - 2F_3'(y) + C_3 F_0''(y) + C_2 F_1''(y) \\ &(1 + C_1)[F_2(y) + 2F_2'(y) + F_2''(y)], F_3(0) = 0, F_3'(0) = 0 \end{aligned} \right), \quad (3.2 i)$$

Its solution is

$$F_3(y, C_1, C_2, C_3) = \frac{1}{2} \left(\begin{aligned} &C_2 F_0(y)[- \cos y - 2y \cos y + 2 \cos^2 y - \cos y \cos 2y + \sin y - 2y \sin y + 2 \cos^2 y \sin y - \\ &\cos 2y \sin y + 2 \sin^2 y - \sin y \sin 2y] + C_1 C_2 F_0(y)[- \cos y - 2y \cos y + 2 \cos^2 y - \\ &\cos y \cos 2y + \sin y - 2y \sin y + 2 \cos^2 y \sin y - \cos 2y \sin y + 2 \sin^2 y - \sin y \sin 2y] + \\ &C_3 F_0(y)[-2 \cos y + 2 \cos^2 y + 2 \sin^2 y] + F_1(y)[- \cos y - 2y \cos y + 2 \cos^2 y - \cos y \cos 2y \\ &+ \sin y - 2y \sin y + 2 \cos^2 y \sin y - \cos 2y \sin y + 2 \sin^2 y - \sin y \sin 2y] + C_1 F_1(y)[-2 \cos y \\ &- 4y \cos y + 4 \cos^2 y + 2 \cos y \cos 2y + 2 \sin y - 4y \sin y + 4 \cos^2 y \sin y - 2 \cos 2y \sin y + \\ &4 \sin^2 y - 2 \sin y \sin 2y] + C_1^2 F_1(y)[- \cos y - 2y \cos y + 2 \cos^2 y - \cos y \cos 2y + \sin y - \\ &2y \sin y + 2 \cos^2 y \sin y - \cos 2y \sin y + 2 \sin^2 y - \sin y \sin 2y] + C_2 F_1(y)[-2 \cos y + \\ &2 \cos^2 y + 2 \sin^2 y] \end{aligned} \right), \quad (3.2 j)$$

We use equations (3.2 d), (3.2 f), (3.2 h) and (3.2 j) we get third order approximate OHAM solution for $p = 1$

$$\tilde{F}(y, C_1, C_2, C_3) = F_0(y) + F_1(y, C_1) + F_2(y, C_1, C_2) + F_3(y, C_1, C_2, C_3), \quad (3.2 k)$$

Using the OHAM technique of section mentioned above on domain $a = 0, b = 1$ we use the residual R that is

$$R = \tilde{F}''(y) + 2\tilde{F}'(y) + \tilde{F}(y), \quad (3.2 l)$$

We have obtained the following values of C_1, C_2, C_3 .where

$$C_1 = -0.6157891011374226, C_2 = -1.648024704278988, C_3 = -0.48317914168990783, \quad (3.2 m)$$

From the above values of C_i 's we get the following approximate OHAM solution

$$\tilde{F}(y) = 1 - 0.424995 y^2 - 0.161002 y^3, \quad (3.2 n)$$

The HPM solution is

$$\tilde{F}(y) = 1 - 0.5y^2 + 0.333333y^3 + 0.04167y^4 + O(y^5), \quad (3.2 \text{ o})$$

The HAM solution is


$$\tilde{F}(y) = 1 - 0.464612y^2 + 0.209414y^3 - 0.00742224y^4 - 0.00671979y^5 + O(y^6). \quad (3.2 \text{ p})$$

The following table 3.2 displays values of the exact solution (3.2 b), OHAM solution (3.1 n) and Error of OHAM.



We compare the two solutions there exists similarity approximations between them also the values of HPM, HAM along with their errors are considered and the comparison is established between the errors of OHAM, HPM and HAM the errors of the technique, OHAM are smaller than the other two. All the mentioned values are described in the following table 3.2, their graphs are drawn in figure 3.2 below.

y	Exact sol	OHAM sol	HPM sol	HAM sol	Er OHAM	Er HPM	Er HAM
0.0	1.000000	1.000000	1.000000	1.000000	0.0 E-0	0.0 E-0	0.0 E-0
0.1	0.995321	0.99591	0.995338	0.995562	5.9 E-4	1.6 E-5	2.4 E-4
0.2	0.982477	0.984288	0.982733	0.983077	1.8 E-3	2.5 E-5	5.9 E-4
0.3	0.963064	0.966098	0.964338	0.963763	3.0 E-3	1.2 E-3	6.9 E-4
0.4	0.938448	0.942305	0.9424	0.938806	3.8 E-3	3.9 E-3	3.5 E-4
0.5	0.909796	0.923877	0.919271	0.90935	4.0 E-3	9.4 E-3	4.4 E-4
0.6	0.878099	0.881778	0.8974	0.876489	3.6 E-3	1.9 E-2	1.6 E-3
0.7	0.844195	0.846976	0.879337	0.841258	2.7 E-3	3.5 E-2	2.9 E-3
0.8	0.808792	0.810436	0.867733	0.804626	1.6 E-3	5.8 E-2	4.1 E-3
0.9	0.772482	0.773125	0.865338	0.76749	6.4 E-4	9.2 E-2	4.9 E-3
1.0	0.735759	0.736007	0.875	0.73066	2.4 E-4	1.3 E-1	5.0 E-3

Table 3.2



(a) Graph of exact and OHAM solutions



(b) Graph of exact and HPM solution

(c) Graph of exact and HAM solutions

Figure 3.2

From the Figure 3.2 (a) we conclude that the two graphs are coincident and also from the above Table: 3.2 We investigate that the OHAM solution and exact solution are in very good agreement on domain $a = 0, b = 1$ showing that the method is effective. Solid curve shows exact solution dotted curve shows OHAM solution, HPM solution and HAM solution. From figure 3.2(b) the HPM and exact solutions are not coincident again the HAM and exact solutions are coincident as shown in figure 3.2(c), However generally the OHAM errors are smaller and there exists accuracy in the values of OHAM and the method is reliable and effective.

Problem 3.3:

For $y \in [0,1]$ we consider the following linear differential equation

$$F'''(y) + F(y) - (7 - y^2)\cos y - (y^2 - 6y - 1)\sin y = 0, F(0) = 0, F'(0) = -1, F'(1) = 2\sin(1), \quad (3.3 a)$$

The exact solution of the problem is

$$F(y) = (y^2 - 1)\sin y, \quad (3.3 b)$$

Applying the technique of OHAM that is described in the above section. The Zeroth order problem is

$$F_0'''(y) = 0, F_0(0) = 0, F_0'(0) = -1, F_0'(1) = 2\sin(1), \quad (3.3 c)$$

Its solution is

$$F_0(y) = \frac{1}{2}(-2y + y^2(1 + 2\sin(1))), \quad (3.3 d)$$

First order problem is

$$F_1'''(y, C_1) = \begin{pmatrix} C_1[-7\cos y + y^2\cos y + \sin y + 6y\sin y - y^2\sin y + F_0(y)] + (1 + C_1)F_0'''(y) \\ F_1(0) = 0, F_1'(0) = 0, F_1'(1) = 0 \end{pmatrix} \quad (3.3 e)$$

Its solution is

$$F_1(y, C_1) = \frac{1}{240}C_1 \begin{pmatrix} -3120 - 240y + 135y^2 - 10y^4 + 2y^5 - 480y^2\cos(1) + 3120\cos y - 240y^2\cos y + \\ 950y^2\sin(1) + 4y^5\sin(1) + 240\sin y + 1440y\sin y - 240y^2\sin y \end{pmatrix}, \quad (3.3 f)$$

Second order problem is

$$F_2'''(y, C_1, C_2) = \begin{pmatrix} C_2[-7\cos y + y^2\cos y + \sin y + 6y\sin y - y^2\sin y + F_0(y) + F_0'''(y)] + \\ C_1F_1(y) + (1 + C_1)F_1'''(y), F_2(0) = 0, F_2'(0) = 0, F_2'(1) = 0 \end{pmatrix}, \quad (3.3 g)$$

Its solution is

$$F_2(y, C_1, C_2) = \frac{1}{40320}$$

$$\left(\begin{aligned} &C_1[-52416 - 40320y + 22680y^2 - 1680y^4 + 336y^5 - 80640y^2 \cos(1) + 524160 \cos y \\ &- 40320y^2 \cos y + 159600y^2 \sin(1) + 672y^5 \sin(1) + 40320 \sin y + 241920y \sin y - \\ &40320y^2 \sin y] + C_1^2[-1048320 + 1209600y - 468801y^2 - 87360y^3 - 3360y^4 + 714y^5 - \\ &8y^7 + y^8 + 446880y^2 \cos(1) - 1344y^5 \cos(1) + 1048320 \cos y + 483840y \cos y - \\ &80640y^2 \cos y + 475502y^2 \sin(1) + 3332y^5 \sin(1) + 2y^8 \sin(1) - 1693440 \sin y + \\ &483840y \sin y] + C_2[-524160 - 40320y + 22680y^2 - 1680y^4 + 336y^5 - 80640y^2 \cos(1) + \\ &524160 \cos y - 40320y^2 \cos y + 159600y^2 \sin(1) + 672y^5 \sin(1) + 40320 \sin y + \\ &241920y \sin y - 40320y^2 \sin y] \end{aligned} \right), \quad (3.3 \text{ h})$$

Third order problem is

$$F_3'''(y, C_1, C_2, C_3) = \left(\begin{aligned} &C_3[-7 \cos y + y^2 \cos y + \sin y + 6y \sin y - y^2 \sin y + F_0(y) + F_0'''(y)] + C_2[F_1(y) + F_1'''(y)] + \\ &C_1F_2(y) + (1 + C_1)F_2'''(y), F_3(0) = 0, F_3'(0) = 0, F_3'(1) = 0 \end{aligned} \right), \quad (3.3 \text{ i})$$

Its solution is obtained by using the method ,OHAM .The optimal values of the auxiliary constants C_1, C_2 and C_3 are obtained by using Galerkin's method or least square method, using these values we get the series solution.

$$F_3(y, C_1, C_2, C_3) = \frac{1}{159667200}$$

$$\begin{aligned}
& \left(C_1[-2075673600 - 159667200y + 89812800y^2 - 6652800y^4 + 1330560y^5 \right. \\
& - 319334400y^2 \cos(1) + 2075673600 \cos y - 159667200y^2 \cos y + \\
& 632016000y^2 \sin(1) + 2661120y^5 \sin(1) + 159667200 \sin y + 958003200y \sin y \\
& - 159667200y^2 \sin y] + C_1^2[-8302694400 + 9580032000y - 3712903920y^2 - \\
& 691891200y^3 - 26611200y^4 + 5654880y^5 - 63360y^7 + 7920y^8 + \\
& 3539289600y^2 \cos(1) - 10644480y^5 \cos(1) + 8302694400 \cos y + \\
& 3832012800y \cos y - 638668800y^2 \cos y + 3765975840y^2 \sin(1) + \\
& 26389440y^5 \sin(1) + 15840y^8 \sin(1) - 13412044800 \sin y + 3832012800y \sin y] \\
& + C_1^3[8302694400 + 14689382400y - 6081277257y^2 - 1037836800y^3 + \\
& 186278400y^4 - 28113426y^5 - 2882880y^6 - 95040y^7 + 12375y^8 - 44y^{10} + 4y^{11} \\
& + 7443992160y^2 \cos(1) + 24171840y^5 \cos(1) - 15840y^8 \cos(1) - 8302694400 \cos y \\
& + 5748019200y \cos y - 319334400y^2 \cos y - 1867972634y^2 \sin(1) + \\
& 44577852y^5 \sin(1) + 47190y^8 \sin(1) + 8y^{11} \sin(1) - 20437401600 \sin y \\
& + 319334400y^2 \sin y] + C_2[-2075673600 + 89812800y^2 - 6652800y^4 + 1330560y^5 \\
& - 319334400y^2 \cos(1) + 2075673600 \cos y - 159667200y^2 \cos y + \\
& 632016000y^2 \sin(1) + 2661120y^5 \sin(1) + 159667200 \sin y + 958003200y \sin y - \\
& 159667200y^2 \sin y] + C_1 C_2[-8302694400 + 9580032000y - 3712903920y^2 - \\
& 691891200y^3 - 26611200y^4 + 565880y^5 - 63360y^7 + 7920y^8 + 3539289600y^2 \cos(1) \\
& - 10644480y^5 \cos(1) + 8302694400 \cos y + 3832012800y \cos y - \\
& 638668800y^2 \cos y + 3765975840y^2 \sin(1) + 26389440y^5 \sin(1) + 15840y^8 \sin(1) - \\
& 13412044800 \sin y + 3822012800y \sin y] + C_3[-2075673600 - 159667200y + \\
& 89812800y^2 - 6652800y^4 + 1330560y^5 - 319334400y^2 \cos(1) + 2075673600 \cos y + \\
& 159667200y^2 \cos y + 632016000y^2 \sin(1) + 2661120y^5 \sin(1) + 159667200 \sin y + \\
& 958003200y \sin y - 159667200y^2 \sin y] \left. \right) , \tag{3.3 j}
\end{aligned}$$

Now we use equations (3.3 d), (3.3 f), (3.3 h) and (3.3 j) to get third order approximate solution by OHAM for $p = 1$ that is

$$\tilde{F}(y, C_1, C_2, C_3) = F_0(y) + F_1(y, C_1) + F_2(y, C_1, C_2) + F_3(y, C_1, C_2, C_3), \tag{3.3 k}$$

Using the proposed technique of section described above on the domain $a = 0, b = 1$ we use the residual

$$R = \tilde{F}'''(y) + \tilde{F}(y) - (7 - y^2) \cos y - (y^2 - 6y - 1) \sin y, \tag{3.3 l}$$

The following values of C_i 's are found

$$C_1 = 1.062670102836802, C_2 = -2.208103637790291, C_3 = 0.33369993779942651,$$

We use the above values of C_1, C_2, C_3 , the approximate OHAM solution is

$$\tilde{F}(y) = -y + 0.0000488077y^2 + 1.16623y^3 - 0.173884y^5 - 0.00092637y^6 + 0.00852856y^7 + 0.000200624y^8 - 0.000245363y^9 - 1.12908 \times 10^{-6}y^{10} + 4.06776 \times 10^{-6}y^{11} - 3.01577 \times 10^{-9}y^{12} - 4.10508 \times 10^{-8}y^{13} + 1.67752 \times 10^{-10}y^{14} + 2.89409 \times 10^{-10}y^{15} + O(y^{16}).$$

(3.3 m)

The HPM solution is

$$\tilde{F}(y) = -y + 1.66667y^2 + 0.008532y^7 - 0.000220y^9 + O(y^{11}).$$

(3.3 n)

The HAM solution is

$$\tilde{F}(y) = -0.563288 + 32.3288y - 0.776059y^2 - 2.25957y^3 - 0.00180541y^4 + 0.026541y^5 + O(y^6),$$


(3.3 o)

The following table displays the OHAM, HPM, HAM exact solutions and their errors.



y	Exact sol	OHAM sol	HPM sol	HAM sol	Er OHAM	Er HPM	Er HAM
0.0	0.000000	0.000000	0.000000	0.000000	0.0 E-0	0.0 E-0	0.0 E-0
0.1	-0.0988351	-0.0886448	-0.0988249	-0.0988314	1.9 E-4	1.0 E-5	3.6 E-6
0.2	-0.190723	-0.190034	-0.190682	-0.19071	6.8 E-4	4.0 E-5	1.2 E-5
0.3	-0.268923	-0.267535	-0.268833	-0.268899	1.3 E-3	9.0 E-5	2.4 E-5
0.4	-0.327111	-0.32492	-0.326954	-0.327074	2.1 E-3	1.5 E-4	3.7 E-5
0.5	-0.359569	-0.357602	-0.35933	-0.35952	3.0 E-3	2.3 E-4	4.9 E-5
0.6	-0.361371	-0.357602	-0.361042	-0.36131	3.7 E-3	3.2 E-4	6.0 E-5
0.7	-0.328551	-0.324145	-0.328131	-0.32848	4.4 E-3	4.2 E-4	7.1 E-5
0.8	-0.258248	-0.253368	-0.257746	-0.258169	4.8 E-3	5.0 E-4	7.9 E-5
0.9	-0.148832	-0.143667	-0.148271	-0.148747	5.1 E-3	5.6 E-4	8.4 E-5
1.0	0.000000	0.0052575	0.00058439	0.0000869	5.2 E-3	5.8 E-4	8.6 E-5

Table 3.3

We conclude that the errors of technique, OHAM are smaller than HPM and HAM solutions. From the above table 3.3 we conclude that OHAM and exact solutions are in best agreement and the values of the two columns are nearly equal.



(a) Graph of exact and OHAM solutions



(b) Graph of HPM and exact solutions

(c) Graph of HAM and exact solutions

Figure 3.3

In figure 3.3 (a), (b) and (c) we investigate that the graphs of OHAM, HPM and HAM are coincident with the graphs of their exact solutions, but generally we conclude that the technique, OHAM is more effective than the other two. From the above figure 3.3 we investigate that two graphs that is exact and OHAM solution graphs are coincident, this shows that the method OHAM is effective and reliable. Solid curve= exact solution, Dotted curve= OHAM solution, HPM solution and HAM solution.

References

- [1] A. M. Wazwaz, A reliable modification of Adomian decomposition method, App. Math. Comput. 02 (1999) 77-86.
- [2] A. M. Wazwaz, A new method for solving differential equations of the Lane –Emden type, Appl. Math. Comput 118(2001)287-310
- [3] V. Marinca, N. Herisanu, L. Nemes, Optimal Homotopy Asymptotic Method with Application to thin Film flow, Cent. Eur. J. phys. 6(3) (2008) 648-653.
- [4] V. Marinca , N. Herisanu, Application of Optimal Homotopy Asymptotic Method for solving Nonlinear Equations arising in heat transfer. Int comm. Heat Mass Transfer 35(2008) 710-715.
- [5] V. Marinca, N. Herisanu, C. Bota, B. Marinca, An Optimal Homotopy Asymptotic Method applied to the Steady flow of fourth –grade past a porous Plate, Appl. Math. Lett. 22(2) (2009) 245-251.
- [6] G. Adomian, Solving Frontier Problems of Physics: The decomposition method, vol. 60 of Fundamental Theories of physics, Kluwer academic publishers, USA, 1994.
- [7] Javed Ali, S .Islam, Siraj Ul Islam, Gulzaman, The solution of multipoint boundary value problems by the Optimal Homotopy Asymptotic Method, Computers and Mathematics with Applications 59(2010)2000-2006
- [8] M. Esmailpour, D.D. Ganji, Solution of the Jeffery-Hamel flow problem by optimal homotopy asymptotic Method, Computers and Mathematics with applications 59 (2010) 3405-3411
- [9] S.I.A. Tirmizi, On numerical solution of third order boundary-value problems, Commun. Appl. Numer. Math. 7(1991) 309-313.

- [10] H.N. Caglar, S.H. Caglar, E.H.T. wizell, The numerical solution of third order boundary-value problems with Fourth degree B-spline functions, *Int. J. Comput. Math.* 71(1999) 373-381.
- [11] M.K. Jain, Numerical Solution of Differential Equations, Wiley Eastern Ltd, New Delhi, 1984/
- [12] Q.K. Ghor, M. Ahmed, A. M. Siddiqui, *Int. J. Nonlinear Sci. Num. Simul*, 8(2) (2007)) 179-184.
- [13] Fakharian A, Hamidi Beheshti M. T, Najafi M, A New Algorithm for solving Reccati equation using Adomian Decomposition Method WSEAS international conference on Applied Mathematics, Istanbul Turkey, May 27-29 (2006) (pp 74-77)
- [14] B. Chanane, "Optimal Control of Nonlinear System: A Recursive Approach." *Computers Math Applic.*, 1998, vol 35, No., pp. 29-33.
- [15] J.B. burl, Linear Optimal control, Addison Wesley, 1999.
- [16] A.M. Wazwaz, S.M. Al-Syed, "A New modification of Adomian's Decomposition Method for Linear and Nonlinear Operators, " *Appl Math. Comput.*, 2001, vol. 122, pp. 393-405.
- [17] A.H. Nayfeh problems in perturbation, J. Wiley, New York, 1985.
- [18] W.T. Reid, Riccati Differential equations (Mathematics in series and engineering, vol. 86) New York; Academic press. ; 1972
- [19] J.E. Dunn, k. R. Rajagopal, Fluids of differential type: critical review and thermodynamic Analysis, *Int. J, Eng. Sci.* 33 (1995) 689-729.
- [20] B.D. Anderson, J.B. Moore, Optimal Control-Linear quadratic methods Prentice-Hall, New Jersey 1999.
- [21] J. H. He, Limit cycle and bifurcation of nonlinear pronlems, *Chaos, Soli-tons and Fractals*, 26(2005), 827-833.
- [22] S. J. Liao, Comparison between the homotopy analysis method and homotopy perturbation method, *Appl. Math. Comput.* !69, 1186-1194 (2005).
- [23] S. J. Liao, An approximate solution technique not depending on small parameters; A special example, *Internet, J. Non-Linear Mech*, 30(3), (1995) 371-380.
- [24] M. Al-Smadi, A. Freihat, O. Abu Arqub, N. Shawagfeh, A novel multistep generalized differential transform method for solving fractional-order Lü chaotic and hyper chaotic systems, *Journal of Computational Analysis & Applications* 19 (2015) 713-724.
- [25] N. Shawagfeh, O. Abu Arqub, S. Momani, Analytical solution of nonlinear second-order periodic boundary value problem using reproducing kernel method, *Journal of Computational Analysis and Applications* 16 (2014) 750-762.

[26] M. Al-Smadi, O. Abu Arqub, N. Shawagfeh, S. Momani, Numerical investigations for systems of second-order periodic boundary value problems using reproducing kernel method, *Applied Mathematics and Computation* 291 (2016) 137-148.

[27] Z. Abo-Hammour, O. Abu Arqub, S. Momani, N. Shawagfeh, Optimization Solution of Troesch's and Bratu's Problems of Ordinary Type Using Novel Continuous Genetic Algorithm, *Discrete Dynamics in Nature and Society*, vol. 2014, Article ID 401696, 15 pages, 2014. doi.10.1155/2014/401696.

[28] O. Abu Arqub, Fitted reproducing kernel Hilbert space method for the solutions of some certain classes of time-fractional partial differential equations subject to initial and Neumann boundary conditions, *Computers & Mathematics with Applications*, 73 (2017) 1243–1261.

[29] O. Abu Arqub, Approximate solutions of DASs with non-classical boundary conditions using novel reproducing kernel algorithm, *Fundamental Informaticae* 146 (2016) 231-254.

[30] O. Abu Arqub, An iterative method for solving fourth-order boundary value problems of mixed type integro-differential equations, *Journal of Computational Analysis and Applications* 18 (2015) 857-874.

Acknowledgement:

The authors are thankful to the anonymous reviewers and the Editor-in-Chief for the productive comments that led to positive enhancement in the paper.

Competing financial interests:

The author declares that they have no competing interests.