

Antiproliferative activity by ethanolic extract of *Alpinia galanga* Willd in inoculated breast carcinoma cells of C3H mice

ABSTRACT

Background

We investigated antiproliferative activity of ethanolic extract of *Alpinia galanga* Willd (galangal) in the inoculated breast carcinoma of C3H mice, that lead to decreased of tumor volume. In some previous studies ethanolic extract of galangal contains active substances I-Acetoxychavicol acetate (ACA), may be used as anti-cancer activity by inhibiting cell proliferation through inhibition of NFκB (Nuclear Factor Kappa B).

Methods

This experimental study was designed by post test only controlled design group, using 32 C3H mice. The C3H mice were inoculated with tumor cells and then divided into 4 groups: control group (K) and three treatment groups (P1,P2 and P3) given graded doses of ethanolic extract of galangal (225mg/kgBW/day, 450mg/kgBW/day and 675mg/kgBW/day) for 2 weeks. After all mice were terminated, we count the tumor volume of each mice. Correlation between tumor volume and graded doses of ethanolic extract of galangal was analyse by Pearson correlation.

Result

There were no significant differences in tumor volume between control and treatment group at week 1 and 2 but in the end of this study (pra termination week), there is a marked differences of tumor volume between groups. Statistical analysis by Pearson correlation showed a significant result.

Conclusion

Ethanolic extract of *Alpinia galanga* Willd at graded doses can inhibit breast carcinoma proliferative activity that lead to decrease of tumor volume.

Keywords: *antiproliferative activity, tumor volume, breast carcinoma cells, ethanolic extract of galangal*

Background

Breast carcinoma is the second common cancer in Indonesian women, according to data from the Agency for Cancer Registration, Indonesian Association of Pathologist (IAPI) in 2006. but it is the first common cancer in Padang, West Sumatera Several treatment modality such as surgery, radiotherapy, hormonal therapy are well known by now but the mortality rate is still high. Problems often arise in our society is that the cost is not affordable for the treatment of breast cancer¹.

It has been recognized that a large groups of therapeutic agents can stop proliferative activity by many mechanism. Chemopreventive material from plants have prospects as a cancer inhibitor and

has fewer side effects. Antitumor and anticancer compounds in plants of which have been widely recognized by the public is *Alpinia galanga* Willd (galangal). Besides used as a food flavoring, galangal is often used as a traditional medicine to treat gastric disorders, eliminate bloating, anti-fungal, eliminate itching, increase appetite, and sore throat ².

Most of the major mediators of tumor growth is a chronic inflammatory, and NFκB (*Nuclear Factor Kappa B*) is one of the major inflammatory transcription factor that plays a role in the regulation of several tumor cell growth. NFκB is a therapeutic target of several medicinal plants and also found in tumor tissue. Galangal is one of the herbs that have anti-inflammatory effects and have the inhibitory pathway in NFκB, thereby reducing the activity of cell proliferation. Inhibition of cell proliferative activity could lead to decreased of tumor volume ^{3,4,5}.

The major phytoconstituents which have been isolated from the galangal is *I-Acetoxychavicol Acetat* (ACA) and hydroxychavicol acetate (HCA). The extracts of galangal showed acetylcholinesterase-inhibitory, platelet-activating factor (PAF)-inhibitory, antimicrobial, antibacterial, antifungal, antioxidant and apoptosis activities. It has been reported that extract of galangal also showed anti-inflammatory, immunostimulating, antinociceptive, hepatoprotective and anticancer activities. Rusmarilin (2003) proved ethyl acetate extracts of galangal inhibit the proliferation of cancer cells in culture using either a groove cancer cells and primary human cancer cells. Ethyl acetate extract of galangal 100 µg/ml inhibits 100% of lung cancer cell proliferation and at a concentration of 50 µg/ml inhibits 95% of cell proliferative activity. It is also reported that concentration of 200 µg/ml could inhibits 76,36% proliferative activity of serous ovarian papillary adenocarcinoma ^{6,7,8}.

Based on the above we are interest to studied the effect of the ethanolic extract of galangal (*Alpinia galanga*) to the activity of cell proliferation that lead to decrease of tumor volume of breast carcinoma C3H mice with different dose level (225 mg/kg/day, 475 mg/kg day and 675 mg/kg/day).

Methods

This study was an experimental study using post-test only research design controled design group. The sample used is a strain of C3H mice as much as 32 tails, aged 3-4 months, body weight 15-25 grams were obtained from the Anatomical Pathology Laboratory of the Faculty of Medicine, University of Indonesia. Protocols of current study were approved by the Ethical Committee of Faculty of Medicine, Andalas University.

76

77 **Preparation of the galangal extract**

78 The fresh red galangal, 9 months old, that has been harvested, cut into small pieces, dried in
79 an oven 50°C and pulverized to make powder. The dried powder (40 g) was mixed with ethanol
80 (70%), by slowly warming, ethanol has a boiling point of 60 ° C, continuously extracted about 10
81 times. then evaporated with a *rotary vacuum evaporator* into a *crude* extract. Before oral
82 administration to mice, a crude extract of galangal mixed with aquabidest to make a 0,2 ml solution

83

84 **Tumor inoculation, administration of ethanolic extract of galangal and measurement of tumor** 85 **volume**

86

87 Thirty two strains C3H female mice were divided into 4 groups: control group (K), treatment
88 1 (P1), Treatment 2 (P2). Treatment 3 (P3), each group consisted of 8 mice C3H. Tumor inoculation
89 is done by injecting a slurry tumors derived from donor mice subcutaneously in the right axillar
90 region at a dose of 0.2 ml using insulin syringe, containing approximately 10^6 tumor cells alive. After
91 the tumor growth, we measure the tumor volume and terminated one mouse of each groups to make
92 haematoxyllin eosin slide of tumors to prove the microscopic features of tumor cells. .

93 Having proved correct for tumor growth, further mice were treated as follows: Group K were
94 given a standard diet for 2 weeks and did not receive treatment (ethanolic extract of galangal). P1
95 given a standard diet, and ethanolic extract of galangal received 225 mg/kg/day for 2 weeks. P2 given
96 a standard diet, ethanolic extract of galangal 450 mg/kg/day for 2 weeks. P3 given a standard diet,
97 ethanolic extract of galangal 675 mg/kg/day for 2 weeks.

98 Body weight of mice were weighed at the start of the study, subsequent to the first week, the
99 second week of the beginning and the end of the second week before termination. Measurement of
100 tumor volume started 1 week after inoculation, the second week of the beginning and the end of the
101 second week before termination. After the treatment period ended, all the mice were terminated, cut
102 the tumor tissue and measure the tumor volume using the ruler (in mm^3)..

103


104 **Results**

105 **Morphological evaluation**

106 At the beginning of the first week after inoculation, the tumor mass has been raised at the site
107 of inoculation. One mouse of each groups were randomly taken and sacrifice to examined tumor cells

108 through histopathological examination with HE staining (*hematoxylin eosin*). From the results of the
109 entire slides obtained positive adenocarcinoma tumor cells (Figure 1).

110



111

112 Figure 1. Microscopic features of breast carcinoma of C3H mice (Haematoxyllin eosin).

113 A. Breast carcinoma, one week after inoculation (Haematoxyllin eosin, 100 x).

114 B. Breast carcinoma one week after inoculation (Haematoxyllin eosin, 400 x).

115

116 Measurement of tumor volume

117 Before the termination, we measure the body weight of all mice. The weight of the smallest
118 mice prior to final termination is 18,4 g and largest is 25,8 g. After the termination, we cut the tumor
119 and performed the measurement of tumor volume. The smallest mice tumor volume contained in P3
120 group is 12.6 mm³ and the largest found in the control group which is 216.4 mm³.

121



122

123

Figure 2. Tumor volume of each groups

Table 1. Tumor volume of each groups

Groups	Tumor volume					
	M1		M2		After termination	
K 1	26,6	Mean =	49,2	Mean =	182,5	Mean =
K 2	16,6	23,2	12,9	33,46	192,5	176,04
K 3	19,7		39,5		161,8	
K 4	23,8		25,1		187,4	
K 5	25,8		25,1		216,4	
K 6	19,8		23,2		149,7	
K 7	30,1		59,2		142,0	
K 8	22,8		-		-	
P 1.1	17,4	Mean =	15,0	Mean =	17,1	Mean =
P 1.2	20,1	21,84	61,8	36,7	93,6	51,53
P 1.3	16,2		35,7		47,8	
P 1.4	17,8		34,1		52,8	
P 1.5	32,5		48,9		61,3	
P 1.6	15,2		27,7		45,7	
P 1.7	33,7		33,7		42,4	
P 1.8	20,2		-		-	
P 2.1	25,8	Mean =	60,5	Mean =	65,4	Mean =
P 2.2	20,9	25,56	19,9	38,07	33,1	53,57
P 2.3	24,2		32,1		53,6	
P 2.4	27,6		33,8		51,5	
P 2.5	29,1		53,9		43,6	
P 2.6	30,9		31,4		63,9	
P 2.7	20,4		34,9		63,9	
P 2.8	29,4		-		-	
P 3.1	17,6	Mean =	18,3	Mean =	54,1	Mean =
P 3.2	22,3	22,4	19,7	29,93	25,8	44,91
P 3.3	18,8		25,3		12,6	
P 3.4	18,9		40,4		71,6	
P 3.5	29,4		44,0		42,1	
P 3.6	20,8		30,1		59,2	
P 3.7	29,0		31,7		49,0	
P 3.8	18,4		-		-	

From the table above shows that the mean tumor volume of the control and treatment groups on week 1 and week 2 is not too different but the mean tumor volume after termination shows marked differences. Statistical analysis by Pearson ($p = 0,000$) shows that there is a significant relationship between the dose of galangal with a reduction in tumor volume.

Discussion

Several studies have shown that a number of herbal medicine have anticancer potential in vitro and in vivo. Those herbal medicine might be good candidate for the development of anti-cancer drugs. Inhibition of cancer cell proliferation, growth factor signaling, induction of apoptosis, etc., contributes to chemopreventive properties⁸.

We used a strain C3H mice in this study because this mice types have a tendency to get a malignancy through genetic susceptibility and transmission of virus MMTV (*Mouse Mammary Tumor Virus*). Using the technique of inoculation in the form of transplant tumor cells will lead to the development of tumor cells. Growth of tumors in mice in the first week after inoculation until the termination varies. In the control group tumor volume tends to increase in size, while in the treatment group tumor volume tends to shrink due to the effects of galangal extract. Turusov and Mohr (1994) states that the growth of mammary tumors in mice C3H strain could reach 2.4 mm per week.⁹

Table 1 shows the mean of tumor volume for the control and treatment groups. The mean of the highest found in the control group and the lowest in the treatment group 3 (P3). It can be concluded that the ethanolic extract of galangal at a dose of 225mg/kg/day had inhibit proliferative activity, and when the dose was increased up to 675mg/kg/hr, the tumor volume is decreased more smallest.

Galangal (*Alpinia galanga*) with ACA and HCA as active substance, has the potential to reduce the incidence or mortality of cancer by reducing the activity of proliferation. This can be explained that most of the major mediators of tumor growth is a chronic inflammatory and NFκB is one of the major inflammatory transcription factor that plays a role in the regulation of several tumor cell growth. NFκB also found in tumor tissue and is the molecular target of several medicinal plants that have anti-cancer effects. Galangal containing ACA has a role inhibiting NFκB through the inhibition of cell cycle regulatory proteins cyclin D1 which results in barriers to the transition phase of the cell cycle from G1 to S, this activity decreases cell proliferation and tumor cells do not grow.^{3,4,5}. Ethanolic extract of galangal has been reported to posses antioxidant activity through strong superoxide anion scavenging activity⁷.

Several previous studies have use galangal as an alternative therapy for cancer because galangal has cytotoxic activity. Hartono (2009) which uses ethyl acetate extract of galangal as well

as research of Herla Rusmarilin (2003) has been proved that the ethyl acetate extract of galangal was found to inhibit the proliferation of cancer cells in culture good using groove cancer cells and primary human cancer cells.^{6,7,10} Similarly, Lee and Houghton (2005) and Chudiwal et al (2010) also have proved the efficacy of anti-cancer ACA against MCF7 (*Michigan Cancer Foundation 7*) on breast cancer cells and COR L23 in lung cancer.^{7,11}

Conclusion

Administration of ethanolic extract of *Alpinia galanga* Willd in graded doses could inhibit cell proliferative activity of breast carcinoma C3H mice, that lead to decrease of tumor volume.

Ethical approval

The authors have obtained ethical approval from Ethical Commision, Faculty of Medicine, Andalas University, Padang, Indonesia.

References

1. Association of Cancer Registry IAPI. Histopathological Data. Indonesian Health Ministry, 2006.
2. Sinaga Erna, *Alpinia Galanga* (L).Willd, Herbal and Traditional Medicine Center of Research and Development /P3TO Unas, 2000.
3. Gautam Sethi, Bokyoung Sung, Bharat B. Aggarwal, 2008, Nuclear Factor κ B activation: From bench to bedside, minireview, *Experimental Biology and Medicine*, 233:21-31
4. Shishir Shisodia, Bharat B. Aggarwal, 2004, Nuclear factor κ B: a friend or a foe in cancer?, *Biochemical Pharmacology*, Elsevier, 68:1071-1080
5. Bharat B. Aggarwal, Shishir Shisodia, 2006, Molecular targets of dietary agents for prevention and therapy of cancer, Elsevier, *Biochemical Pharmacology* 71: 1397-1421
6. Rusmarilin H. Anticancer activity of local *alpinia galangal* L (SW) rhizome extracts on cancer cell line of human and mice transplanted with primary tumor cell. [Disertation], Institut Pertanian Bogor.2003
7. Chudiwal AK, Jain DP, Somani RS. *Alpinia galangal* wild-an Overview on Phyto pharmacological properties. *Indian Journal of Natural Products and Resources* .2010 Juni; Vol.I(2).pp.143-149
8. Samarghandian S, Hadjzadeh MAR, Afshari JT, Hosseini M. Antiproliferative activity and induction of apoptotic by ethanolic extract of *Alpinia galanga* rhizhome in human breast carcinoma cell line. *BMC Complementary and Alternative Medicine*, 14 : 192, 2014.

- 202 9. Turusov VS and U.Mohr, 1994, Pathology of Tumours in Laboratory Animals, Volume II
203 Tumours of the mouse, second edition, International Agency for Research on Cancer, Lyon
204 France
- 205 10. Hartono NWB. The effects of extract galangal to cell proliferative activity and apoptotic index
206 of breast adenocarcinoma of C3H mice.[Thesis]. Diponegoro University, 2009
- 207 11. Lee CC , Houghton P. Cytotoxicity of plants from Malaysia and Thailand used traditionally
208 to treat cancer . Elsevier. J Ethnopharmacol.2005 September 14;100(3): 237-43
209