

Original research Articles**Effects of various concentrations of Quail egg solution on glycemia and antioxidant parameters of alloxan-induced diabetic rats****ABSTRACT**

This study evaluated the effects of different concentrations of quail egg solution on glycemia and some antioxidant markers in alloxan-induced diabetic rats. Thirty adult male albino rats were assigned to 5 groups of 6 rats per group. Groups 2, 3, 4 and 5 were made diabetic by single intraperitoneal injection of 160 mg/kg alloxan monohydrate. Upon establishment of diabetes (fasting blood glucose levels above 126 mg/dl), the rats in groups 2, 3 and 4 were respectively administered with 30, 15 and 7.5 mg/ml orally daily for 21 days. Groups 1 and 5 were administered with distilled water. Fasting blood glucose (FBG) levels of the rats were assessed 1 h, 6 h, 24 h and 21 days post treatment. On the 21st day, blood samples were collected for malondialdehyde (MDA), superoxide dismutase (SOD), and reduced glutathione (RSH) assays. Results indicate that administration of quail egg solution at the concentration of 30 mg/ml to the diabetic rats significantly ($p < 0.05$) reduced the FBG from 343.80 to 87.20 mg/dl on day 21 post treatment. There was significant ($p < 0.05$) reduction in the mean MDA values of the 30 mg/ml-treated groups compared to the negative control group. The SOD activity and glutathione levels of 30 mg/ml-treated diabetic rats were significantly ($p < 0.05$) higher compared to that of the diabetic untreated group. It was concluded that administration of quail egg solution especially at the concentration of 30 mg/ml to the diabetic rats resulted in hypoglycemia and improvement in the levels and activities of *in vivo* antioxidant parameters.

Key words: Quail egg, glycemia, antioxidant, diabetic rats

INTRODUCTION

Diabetes mellitus is a chronic metabolic disorder that is fast becoming a global problem with huge social, health and economic consequences. It is estimated that in 2010, there were about 285 million people (approximately 6.4 % of the adult population) suffering from diabetes. This number is estimated to increase to 430 million in near future in the absence of better control or cure [1]. Data from world health organization (WHO), suggest that Nigeria has the greatest number of people living with diabetes in Africa [2]. Traditionally, types 1 and 2 diabetes mellitus are recognized by WHO, although gestational diabetes mellitus and other types of diabetes have been described. Basically, the clinical signs of diabetes mellitus include polyuria, polydipsia, polyphagia, weakness, asthenia blurred vision and impaired wound healing. Diagnosis involves assay of fasting blood glucose levels, measurement of glycosylated haemoglobin, assay of urinary sugar and plasma insulin values while treatment tailored towards reduction in the fasting blood sugar levels are achieved by dietary control, exercise and use of oral hypoglycemic [3]

Oxidative stress associated with diabetes mellitus has been widely reported. Diabetic patients present with increased levels of malondialdehyde consequent upon exuberant cellular lipid peroxidation. Increased lipid peroxidation presents a close relationship with the high glycemic levels and oxidative stress in diabetes mellitus [4]. Superoxide dismutase and glutathione are antioxidants that protect against lipid peroxidation. Assay of MDA, SOD and glutathione are important in assessment of severity and amelioration of diabetes mellitus [5, 6]

The synthetic drugs in use for treatment of diabetes mellitus are not only expensive but have complicated mode of intake and have several side effects. Therefore the search for alternative therapy has been advocated for. Numerous nutritional and therapeutic values of quail egg have been reported. The quail eggs are rich sources of antioxidants, minerals and vitamins [7]. Experts in natural medicine believe that quail has positive effects on people with hypertension, liver problem, hyperlipidemia and anaemia [8]

This study evaluated the effects of quail egg solution on glycemia and *in vivo* antioxidant parameters of alloxan-induced diabetic rats

66

67

68

69

70

71 MATERIALS AND METHODS

72 Animals

73 Adult male Wistar albino rats of 10 to 16 weeks and average weight of 160 ± 15 g were obtained
74 from the Animal House of the Faculty of Biological Sciences, University of Nigeria, Nsukka,
75 Enugu state, Nigeria. The animals were acclimatized for the duration of 7 days under standard
76 environmental conditions with a 12 h light/dark cycle maintained on a regular feed (vital feed)
77 and water *ad libitum*.

78 Quail Egg

79 Quail eggs used were obtained from the Faculty of Veterinary Medicine, University of Nigeria,
80 Nsukka, Enugu state, Nigeria Farm. The freshly laid eggs weighed between 10-15 g.

81

82 Experimental Design

83 Thirty adult male albino wistar rats were assigned into 5 groups of 6 rats per group. Following
84 establishment of diabetes mellitus on the 2nd day post induction, the rats were treated with
85 different concentrations of quail egg solution as follows:

GROUP	TREATMENT
One	Non diabetic rats administered 10ml/kg distilled water (positive control)
Two	Diabetic rats administered 30 mg/ml quail egg solution (highest concentration)
Three	Diabetic rats administered 15 mg/ml quail egg solution (medium concentration)
Four	Diabetic rats administered 7.5 mg/ml quail egg solution (lowest concentration)

Five	Diabetic rats administered 10 ml/kg distilled water (Negative control)
------	--

86

87 Upon establishment of diabetes, the quail egg solution was administered daily through the oral
88 route for 21 days. The FBG levels were assessed 1 h, 6 h 24 h and 21 days post treatment while
89 blood for assay of oxidative stress markers was collected on the 21st day post treatment

90 **Preparation of Quail Egg**

91 An empty beaker was weighed (A g). The shells of the quail eggs were broken with spatula and
92 the contents emptied into the beaker. The weight of the beaker and the contents were recorded as
93 B g. The weight of the contents of the egg alone was obtained by subtracting the weight of the
94 beaker and its contents from the weight of the beaker alone. Thus the weight of the egg yolk and
95 albumen, C was expressed mathematically thus:

$$96 \quad C (g) = B (g) - A (g)$$

97 C (g) was solubilized in a calculated quantity of distilled water to make a desired concentration
98 of quail egg solution and thereafter, serial dilutions of the stock solution were made for the
99 different groups

100

101 **Induction of Experimental Diabetes mellitus**

102 Diabetes was induced in rats using the method described by [9]. The rats were fasted for 16 h
103 prior to induction of diabetes. Diabetes was induced by single intraperitoneal injection of
104 alloxan monohydrate at the dose of 160 mg/kg. Diabetes was established on day two post
105 induction on confirmation of FBG levels above 7 mmol/l or 126 mg/dl

106

107

108

109

110 **Blood Collection**

111 Blood samples were collected from the animals into EDTA bottle using orbital techniques for
112 haematological determination. Blood samples were collected from the retrobulbar plexus of the
113 median canthus of the eye of the rats [10].

114

115 **Estimation of Superoxide dismutase**

116 Superoxide dismutase activity was assayed by the method of [11]. 0.5 ml of plasma was
117 diluted to 1.0 ml with ice cold water, followed by 2.5 ml ethanol and 1.5 ml chloroform (chilled
118 reagent). The mixture was shaken for 60 seconds at 4⁰C and then centrifuged. The enzyme
119 activity in the supernatant was determined as follows. The assay mixture contained 1.2 ml of
120 sodium pyrophosphate buffer, 0.1 ml of PMS and 0.3 ml of NBT and approximately diluted
121 enzyme preparation in a total volume of 3 ml. The reaction was started by the addition of 0.2 ml
122 NADH. After incubation at 30⁰C for 90 seconds, the reaction was stopped by the addition of 1
123 ml glacial acetic acid. The reaction mixture was stirred vigorously and shaken with 4 ml n-
124 butanol. The mixture was allowed to stand for 10 minutes, centrifuged and butanol layer was
125 separated. The colour intensity of the chromogen in the butanol layer was measured in a
126 spectrophotometer at 520 nm. A system devoid of enzyme served as control. One unit of enzyme
127 activity is defined as the enzyme concentration, which gives 50% inhibition of NBT reduction in
128 one minute under assay conditions. SOD activity was expressed as U/ml of plasma.

129

130 **Estimation of Lipid Peroxidation (Malondialdehyde)**

131 Lipid peroxidation was estimated by measuring spectrophotometrically, the level of the
132 lipid peroxidation product, malondialdehyde (MDA) as described by [12]. A volume, 0.1 ml of
133 the serum was mixed with 0.9 ml of H₂O in a test tube. A volume, 0.5 ml of 25% TCA
134 (trichloroacetic acid) and 0.5 ml of 1% TBA (thiobarbituric acid) in 0.3% NaOH were also added
135 to the mixture. The mixture was boiled for 40 minutes in water-bath and then cooled in cold

water. Then 0.1 ml of 20% sodium dodecyl sulfate (SDS) was added to the cooled solution and mixed properly. The absorbance was taken at wavelength 532nm and 600 nm against a blank.

$$\% \text{ TBARS} = \frac{A_{532} - A_{600}}{0.5271 \times 0.1} \times 100 \quad (\text{mg/dl})$$

$$0.5271 \times 0.1$$

Estimation of Reduced Glutathione

The reduced glutathione level was determined by the method of [13]. This method was based on the development of yellow colour when 5,5'- dithio-bis-2-nitrobenzoic (DTNB) is added to compound containing sulphydryl groups. The colour developed was read at 412 nm in spectrophotometer.

Statistical Analyses

Data obtained were analyzed using One-way Analysis of Variance (ANOVA). Variant means were separated using Duncans Multiple range Pos hoc Test. Results were presented as Mean $\pm$ Standard Error of the Mean (Mean $\pm$ SEM)

RESULTS

Table 1: Effect of quail egg solution on the fasting blood glucose levels of alloxan-induced diabetic rats

GROUP	FASTING BLOOD GLUCOSE LEVELS (mg/dl)					
	PRE INDUCTION	POST INDUCTION (PI)	1 h PI	6 h PI	24 h PI	21 Days PI
1	84.80 ± 4.39 ^a	84.60 ±4.31 ^a	85.20± 4.32 ^a	84.40 ±2.01 ^a	86.80 ± 2.35 ^a	83.80 ±2.09 ^a
2	92.80 ± 1.98 ^{ab}	343.80 ±6.66 ^b	235.00 ±16.88 ^b	158.60 ±20.15 ^b	109.60 ± 4.15 ^a	87.20 ±3.22 ^a
3	91.40 ± 2.24 ^{ab}	371.00 ±5.73 ^b	335.20± 4.71 ^c	319.20 ±37.67 ^c	267.40 ±44.37 ^b	233.60 ± 14.29 ^b
4	96.80 ± 1.11 ^b	344.60 ±4.35 ^b	326.20± 5.04 ^c	311.20 ±5.81 ^c	281.00 ±16.80 ^b	274.40 ± 10.79 ^c
5	92.00 ± 1.47 ^{ab}	342.00 ±5.00 ^b	343.40± 7.39 ^c	353.00 ±10.28 ^c	347.20 ±14.46 ^c	329.20 ±12.46 ^d

Different superscript along the same column indicate significant difference at $p<0.05$

There was a significant ($p<0.05$) decrease in the FBG levels of the groups treated with quail egg solution 24 h post treatment when compared to that of the diabetic untreated group. The group treated with 30 mg/ml of quail egg (group 2) showed the most profound decrease in FBG levels when compared with the other treated groups. The FBG levels of 30 mg/ml-treated rats was comparable to that of the normal control groups 21 days post treatment.




Figure 1: Effect of subchronic administration of quail egg solution on malondialdehyde values of alloxan-induced diabetic rats

The plasma malondialdehyde values of rats in group 2 were significantly ($p < 0.05$) reduced compared to that of the negative control group but was statistically comparable ($p > 0.05$) to those of the normal control group.




Figure 2: Plasma superoxide dismutase activity of alloxan-induced diabetic rats treated with quail egg solution

There was significant ($p < 0.05$) increase in the mean activity of SOD for the group treated with 30 mg/ml of quail egg solution when compared with the diabetic untreated group. There was no significant ($p > 0.05$) difference in the mean activity of SOD of the 15 and 7.5 mg/kg treated rats when compared to the diabetic untreated group




Figure 3: Reduced glutathione values of alloxanized rats treated with varying concentrations of quail egg solution

There was a significant ($p < 0.05$) increase in the reduced glutathione concentration in the groups treated with quail egg solution when compared to the diabetic untreated group. There was no significant ($p > 0.05$) difference in the reduced glutathione levels between the 15 and 7.5 mg/dl quail egg solution-treated groups.

DISCUSSION

The elevated glucose levels observed in groups 2-5 rats compared to the group 1 rats (**Table 1**) was attributed to the effect of alloxan monohydrate administered to the rats in groups 2-5. Alloxan monohydrate, a glucose analogue has been shown to produce hyperglycemia through either partial or complete destruction of beta cells of the islets of langerhans which secretes insulin [14]. Insulin deficiency leads to increased blood glucose levels [15]. The elevated glucose level was significantly reduced in all the treated groups especially in the 30 mg/ml-treated diabetic rats which decreased from 343.80 ± 6.00 mg/dl to 87.20 ± 3.22 mg/dl 21 days post-treatment. The reduction was credited to the effect of quail egg solution. Quail egg contains important nutrients such as amino acids (leucine, valine and alanine), vitamins (vitamin E) and minerals such as zinc [16]. Leucine has been reported to lower blood glucose levels in type 2 diabetic patients [17]. Leucine is one of the most potent insulin secretagogues among the branched chain amino acids that facilitate glucose induced insulin release from pancreatic beta cells [17]. Alanine equally plays a key role in maintaining glucose levels in the body by helping to convert glucose to energy [18]. Marvin et al, [19] reported that supplementation of vitamin E might alter insulin receptors in muscle or adipose tissue by increasing membrane motility.

The results obtained for MDA showed that the diabetic rats treated with the highest concentration of quail egg solution had significantly lower levels of MDA compared to other groups (**Figure 1**). This indicates that at very high concentration, quail egg may mitigate lipid peroxidation. This observation may be attributed to antioxidant content of quail egg such as Vitamin E. Antioxidants are known to protect against lipid peroxidation [20].

Similarly, the diabetic rats treated with quail egg solutions demonstrated significant increases in the SOD activities compared to the negative control group (**Figure 2**). This may be related to the Zinc content of quail egg [21]. Zinc is a co-factor for the enzyme SOD this facilitates the activity of SOD [22].

The depleted reduced glutathione in the diabetic untreated rats compared to the treated group (**Figure 3**) is a consequence of diabetes. Reduced glutathione level is usually depleted in diabetic conditions due to hyperglycemia which channels glucose to the polyol pathway depleting NADPH required for glutathione regeneration [23]. Mitigation of hyperglycemia may therefore improve the levels of reduced glutathione. It is also possible that the zinc content of quail egg may inhibit glutathione depletion [24]. In conclusion, administration of quail egg solution especially at the concentration of 30 mg/ml to diabetic rats resulted in hypoglycaemia and improvement in the *in vivo* antioxidant marker parameters

263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

CONSENT

Not applicable

ETHICAL APPROVAL

All authors hereby declare that “principles of laboratory animal care” (NIH publication No 85-23, revised 1985) were followed, as well as specific national laws. All experimemts have been examined and approved by the appropriate ethics committee.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. King, H, Aubert, R.E, and Herman WH (1998) Global burden of diabetes, 1995-2025: prevalence, numerical estimates and projections. *Diabetes care*, 21 (9): 1414-1431
2. Chinenye S and Young E (2011) state of Diabetes care in Nigeria. *Health Journal* 11(4): 1-11
3. WHO (1999). "Definition Diagnosis and Classification of Diabetes Mellitus and its complications.
4. Yang Z, laubach VE, French BA and Kron IL (2009) acute hyperglycemia enhances oxidative stress and exacerbate myocardial infarction by activating nicotinamide adenine dinucleotide phosphate oxidase during reperfusion. *Journal of thoracic and cardiovascular surgery* 137(3):723-729
5. Kim, CH (2013) Expression of extracellular superoxide dismutase protein in diabetes, *Archives of plastic surgery* 40 (5): 517-521
6. Calabresse V, Cornelius C and Leso V (2012) oxidative stress , glutathione status, sirtuin and cellular stress response in type 2 diabetes, *Biochemica et Biophysica Acta*, 1822 (5); 729-736
7. Lalwani P (2011) quai egg nutrition http://www.buzzle.com/articles/quail_egg-nutrition.html. Accessed 20th September, 2012
8. Genchev, A (2012) quality and composition of Japanese quail eggs. *Trakia Journal of Sciences* 10(2): 91-101
9. Venogopal, P.M., Prince, P.S.M., Pari, .L. (1998).Hypoglycemic activities of *Syzigium cumini* seeds;effect on lipid peroxidation in alloxan diabetic rats", *Journal of Ethnopharmacology* **61**:1-7.
10. Parasuraman, S., Raveendran, R., and Kesavan, .R. (2010). Blood sample collection in small laboratory animals. *J Pharmacol Pharmacother* **1**(2): 87-93
11. Kakkar, SB; Viswanaathan, P (1989) "A Modified Spectrophotometric Assay of Superoxide dismutase" *Indian J. Biochem. Biophysics*, 21: 130-132.
12. Wallin, B; Rosengren, B; Shertzer, HG; Camejo, G (1993) "Lipoprotein Oxidation and Measurement of TBARS Formation in a Single Microliter Plate; its use for Evaluation of Antioxidants". *Anal. Biochem.* 208:10-15
13. Beutler E, Duron O, Kelly BM (1963) Improved method for determination of blood glutathione. *J Lab. Clin. Med.* **61**: 882-888

14. Lenzen, S (2008) "The Mechanisms of Alloxan and Streptozotocin-Induced Diabetes" *Diabetologia*; 51(2): 216-226. doi: 10.1007/s00125-0070886-7. PMID 18087688
15. Mclethie NG (2002) Alloxan diabetes mellitus, albeit a minor one; *Journal of Royal College of Physicians of Edinburgh* 32 (2): 134-142
16. Tanarson T, Wanna T, Wattasit S (2013) Nutrient benefits of quail (*cortunix cortunix japonica*) eggs. *International Journal of Research and Publication* 3(5): 1-8
17. van Loon LJ (2012) leucine as pharmaconutrient in health and disease. *Current Opinion in Clinical nutrition and Metabolic care* 15(1): 71-77
18. Vitalhealthzone, Alanine (2007a) <http://www.vitahealthzone.com/nutrition/aminoacids/alanine.html> Accessed 20th september, 2014
19. Marvin L, Bierenbaum MD and Frank J (1985) the effect of supplemental vitamin E on serum parameters in diabetics, post coronary and normal subjects. *Nutritional Research Journal* 31: 1171
20. Rikans LE and Hornbrook KR (1997) Lipid peroxidation, antioxidant protection and aging. *Biochimica et Biophysica (BBA)-Molecular Basis of Disease* 1362(3):116-127
21. Shaheen AA and Abd El-Fattah AA (1995) effect of dietary zinc on lipid peroxidation, glutathione, protein thiols levels and superoxide dismutase activity in rat tissues. *International Journal of Biochemistry and Cell Biology*; 27: 89-95
22. Rae TD, Torres AS, Pufahl RA and O'Halloran TV (2001) Mechanism of Cu Zn-superoxide dismutase activation by the human metallochaperon hccs. *Journal of Biological Chemistry* 275:5166-5176
23. Moussa SA (2008) oxidative stress in diabetes mellitus; *Romanian Journal of Biophysics* 18: 225-236
24. Gibbs PNB, Gore MG and Jordan PM (1985) investigation of the effect of metal ions on the reactivity of thiol groups in human 5-aminolevulinatedehydratase; *Biochemistry Journal*; 225:573-580