

Case studies

Ulcerated Giant Lipoma over Nape of Neck – A Rare Case

Abstract - Lipoma is benign tumor composed of fat cells of adult type. Lipoma called as universal tumor as it can occur anywhere in body. Lipoma with size of more than 10 cm in one dimension or weighing a minimum of 1000 gm. called as giant lipoma [1]. Surgical excision is the treatment of choice for lipoma. We report a case of giant lipoma (4.5 kg.) with decubitus ulcer over nape of neck which is rare entity.

Keywords – Benign tumor, Giant lipoma, Lipoma, Tumor, Universal tumor.

Introduction – lipoma is a common benign tumor. The annual incidence of lipoma is one per 1,000 persons. Lipomas are generally slow-growing tumor with a firm rubbery consistency. While about 80% of lipomas are less than 5 cm in diameter, some can reach more than 20 cm and weigh several kilograms. Usually lipomas are asymptomatic but can cause pain when they compress nerves. Solitary lesions are seen about 80% of the time, commonly in women, while multiple lesions are most common in young men [2]. Lipoma is seen in all age group though most commonly seen in fifth and sixth decade [3]. It constitutes five percent of all benign tumors of body and can be found anywhere in the body [4]. Lipoma in head and neck region is not commonly encountered (13%). The first case of lipoma in the neck region was reported over 100 years ago [5]. Amongst the head and neck lipoma, the most commonest location is anterior neck [6].

Case report – We report a case of ulcerated giant lipoma size 15x15 cm, 4.5 kg. weigh, over nape of neck in 60 year old patient. A 60 year old male patient came in outdoor with chief complaint of swelling over nape of neck since 12 years. Initially swelling was small in size, with time swelling was increasing in size gradually. Patient had difficulty during sleeping in supine position and difficulty in neck movement. Since three years he got developed ulcer over summit of swelling. This ulcerated lesion made suspicious diagnosis of malignancy. Cosmetically he felt guilty due to the swelling. Swelling was globular, firm in consistency, freely mobile over base, ulcer over summit with slough. There were visible dilated veins over swelling. Due to weight swelling was hanging down and giving appearance like pedunculated swelling.

Fine needle aspiration cytology report of swelling was showing fat cells. CT scan report revealed that mass was well defined and present in sub-cutaneous plane. There was no communication with spinal cord. Other heretofore logical investigations were within normal limit. Patient was operated in plan OT under general anaesthesia and lipoma was excised with ulcer after giving elliptical incision. Wound closed after placing a drain. Drain was removed on third post-operative day. Patient discharge without any complication on eight post-operative days. Histopathology of biopsy report revealed as lipoma without any evidence of malignancy.

Discussion - A lipoma is a benign tumor composed of adipose tissue (body fat). It is the most common benign form of soft tissue tumor [7]. According to WHO classification of soft tumors these can be classified into nine groups, including lipoma, lipomatosis, lipoblastoma, angioliipoma, myoliipoma of soft tissues, chondroid lipoma, spindle cell lipoma, and finally hibernoma and pleomorphic lipoma [8]. Lipomas are tumors with unexplained pathogenesis and etiology. Hereditary factors have been reported in conditions such as familial multiple lipomatosis [9]. Giant lipomas are defined by Sanchez et al as lesions with size of at least 10 cm in one dimension or weighing a minimum of 1,000 gm. [1]. After long time lipoma may undergo myxomatous degeneration, saponification, calcification, infection, ulceration due to repeated trauma and malignant transformation. Rarely malignant transformation of lipoma into liposarcoma has been described [10, 11]. Most of the lipomas are small but some

lipomas are huge in size that they resemble like malignant swelling. Ulcerated huge lipoma swelling with bleeding arouse suspect of malignancy [12]. FNAC, USG, CT scan and MRI are useful diagnostic tool for final diagnosis.

Conclusion – lipoma is a common benign tumor. Huge lipoma with ulcer over summit over nape of neck is not common entity. Such type of swelling resemble like malignancy. So, this type of swelling should be operated after proper diagnosis. Surgical excision is the treatment of choice for lipoma.



Fig -1: Showing giant ulcerated swelling.



Fig -2: Showing visible dilated vein.



Fig – 3: Excised giant lipoma skin side view.



Fig - 4: Excised giant lipoma internal view.

51 **References –**

- 52 1. Sanchez MR, Golomb FM, Moy JA, Potozkin JR. Giant lipoma: Case report and review of the literature. J
53 Am Acad Dermatol. 1993; 28:266.
- 54 2. Luba MC, Bangs SA, Mohler AM, Stulberg DL. Common benign skin tumours. Am Fam Physician 2003;
55 67: 729-738.
- 56 3. Salam G. Lipoma excision. Am Fam Physician 2002; 65:901-905.
- 57 4. Enzinger FM, Weiss SW. Benign lipomatous tumors. In: Enzinger FM, Weiss SW, eds. Soft Tissue
58 Tumors. 2nd edn. St Louis: Mosby; 1988.p.301-345.
- 59 5. Horne, W.J., Lipoma or Cystoma of the Neck. Proc R Soc Med, 1908. 1(Laryngol Sect): p. 38-39.
- 60 6. Barnes L. tumors & tumor like lesions of the head & neck: In: Barnes L, ed. surgical pathology of the head
61 and neck. New York, NY:Dekker; 1985: 747-758.
- 62 7. Bancroft LW, Kransdorf MJ, Peterson JJ, O'Connor MI (October 2006). "Benign fatty tumors:
63 classification, clinical course, imaging appearance and treatment". Skeletal Radiol. 35 (10): 719–733.
64 doi:10.1007/s002560060189y. PMID 16927086.
- 65 8. Murphey MD, Carroll JF, Flemming DJ, Pope TL, Gannon FH, Kransdorf MJ. From the archives of the
66 AFIP: Benign musculoskeletal lipomatous lesions. Radiographics 2004; 24:1433-1466.
- 67 9. Leffell DJ, Braveman AM. Familiar multiple lipomatosis. Report of a case and a review of the literature. J
68 Am Acad Dermatol 1986 Aug; 15:257-259.
- 69 10. Mentzel T. Cutaneous Lipomatous Neoplasms. Semi Diagn Pathol, 2001, 18: 250-257.
- 70 11. Mentzel T. Biological continuum of benign, atypical, and malignant mesenchymal neoplasms – does it
71 exist? J Pathol, 2000, 190: 523-525.
- 72 12. Gluscek S. Giant lipoma of thigh. Wiad Lek. 1987 Jun 15; 40(12):845-848.