



Original Research Article

ASSESSMENT OF THE RETICULAR FIBRES ON CADMIUM INDUCED LIVER

DAMAGE FOLLOWING ORAL ADMINISTRATION OF MORINGA OLEIFERA OIL

AND ANACARDIUM OCCIDENTALE OIL IN WISTAR RATS.

Abstract

Background: Cadmium is one of the most toxic pollutants in the environment. This study aim at assessment of the reticular fibres on cadmium induced liver damage following oral administration of *moringa oleifera oil and anacardium occidentale oil* in wistar rats.

Methods: Thirty-five wistar rats (80-180g) were randomly selected and divided into seven groups after being well acclimatized for two (2) weeks. Each group contained five rats. Group A which was the control group received phosphate buffer, Group B received Cadmium only (3.5mg/kg body weight). Group C received Vitamin C and Vitamin E, Group D received Vitamin C and Vitamin E and Cadmium. Group E received Cadmium and Moringa oleifera oil, Group F received Cadmium and Cashew nut oil and Group G received Cadmium (3.5mg/kg body weight) and Moringa oleifera oil and Cashew nut oil

Results: Microscopic examination revealed normal histological architecture of the liver, the portal vein and the reticular fibres were seen clearly and there was orderly arrangement of the reticular fibres in group A. In group B, there was enlargement of the portal vein and distortion of the arrangement of the reticular fibres. In group C, resolution was observed and the reticular fibres were seen though not orderly arranged. In group D, the portal vein and reticular fibres were seen though not well arranged and the sinusoidal beds were a little bit distorted. In groups E, F and G, there was restoration of the reticular fibres and the portal vein was seen unlike the

group B which was induced with Cadmium only. This therefore suggests that *Moringa oleifera* oil and Cashew nut oil have ameliorative effects that led to the regeneration of the damaged and distorted reticular fibres.

Conclusion: The histological architecture of the livers of the treated rats showed the central vein and restoration of the hepatocytes these show ameliorative effect of *Moringa oleifera* seed oil and Cashew nut oil on Cadmium induced liver damage.

Key words: *Moringa oleifera*, *Anacardium occidentale*, Oxidative stress, Liver, Cadmium.

Introduction

Cadmium is a metal with an oxidation state of +2. It is chemically similar to zinc and occurs naturally with zinc and lead in sulfide ores [23]. It is a soft white solid whose density is 8.64 g/cm³, melts at 320.9 °C and boils at 765 °C at 100 kPa. It is soluble in dilute nitric and concentrated in sulphuric acids [11][18][21]

Cadmium metal is used mainly as an anticorrosive, electroplated onto steel. Cadmium sulfide and selenide are commonly used as pigments in plastics. Cadmium compounds are used in electric batteries, electronic components and nuclear reactors [7][18]. Fertilizers produced from phosphate ores constitute a major source of diffuse cadmium pollution. Cadmium (Cd) is one of the most toxic metal ions of the environment which is bound in the air, food and water [13]. Cadmium has been classified as a number 1 category human carcinogen by the International Agency for Research on Cancer of USA because of its carcinogenic properties [12]. Cadmium is a potent human carcinogen and has been associated with cancers of the lung, prostate, pancreas, and kidney.

46 Cadmium can cause osteoporosis, anemia, non-hypertrophic emphysema, irreversible renal
 47 tubular injury, eosinophilia, anosmia and chronic rhinitis. Cadmium intoxication is responsible
 48 for alterations in various metabolic processes [22] and the inhibition of nucleic acid and protein
 49 synthesis (Beyersmann and Hechtenberg [3]. Cadmium has been reported extensively as being
 50 carcinogenic, mutagenic and teratogenic under experimental conditions. Cadmium's action of
 51 being teratogenic has sometimes been attributed to placental or yolk sac damage in rodents [5], it
 52 has also been reported that cadmium is found in early organogenesis-stage embryonic
 53 tissues[16], indicating that embryonic cells may be the direct targets of cadmium action.
 54 *Moringa oleifera* is the most widely cultivated species of a monogeneric family, the
 55 Moringaceae, that is native to the sub-Himalayan tracts of India, Pakistan, Bangladesh and
 56 Afghanistan [8]. It is a small, fast,growing, evergreen, or deciduous tree that usually grows up to
 57 10 or 12 m in height. It is a perennial softwood tree with timber of low quality, but which for
 58 centuries has been advocated for traditional, medicinal and industrial uses [8]. Moringa seed oil
 59 (yield 30-40% by weight), also known as Ben oil, is a sweet non-sticking, non-drying oil that
 60 resists rancidity. It has been used in salads, for fine machine lubrication, and in the manufacture
 61 of perfume and hair care products [20]. Some of its activities include; Antihypertensive and
 62 cholesterol lowering [4]; Hepatoprotective and anticancer, [10]; As biosorbent (for the removal
 63 of Cadmium); Antibacterial and antifungal activities and Antitumor activities amongst others
 64 [17][19]. *Anacardium occidentale* tree is a native of Brazil and the Lower Amazons. Cashew nut
 65 is a high value edible nut. It yields two "Oils" one of these found, between the seed coat (or
 66 pericarp) and the nuts, is called the Cashew Nut Shell Liquid (CNSL). It is not a triglyceride and
 67 contains a high proportion of phenolic compound. It is used in industry as a raw material for
 68 brake lining compounds, as a water proofing agent, a preservative and in the manufacturing of

69 paints and plastics. Cashew apples are sometimes made locally into fruit drinks, wines and
 70 pickles. In some countries they are also Osmo-Sol dried to produce a date- like caramel. An
 71 ability of cashew apples to supply and fortify the nutritional requirement for vitamin C,
 72 particularly in Africa was reported by Akinwale [2]. The importance of the Cashew Nut Kernel
 73 Oil and Cashew Nut Shell Liquid (CNSL) cannot be overemphasized. The fat of nut is
 74 completely natural and unprocessed which is best for the body. It is especially rich in Linoleic
 75 acid (Omega-3) and is least damaging to heart and arteries. In fact, it constitutes about 47% of
 76 the total weight of the nut. Nuts often produce oil half their weight. Cashew has what is called
 77 the 'good fat. Cashew has the right combination of fat and the ratio of saturated to
 78 monounsaturated and polyunsaturated is 1:2:1 which is ideal for human consumption. The
 79 relative abundance of monounsaturated fatty acids in cashew nut is conducive to the promotion
 80 of good health and that the relative abundance of fat in cashew nut in no way poses a nutritional
 81 risk [1]. It has been shown to have antifungal, medicinal and antidiarrhoeal properties amongst
 82 others [14]. It has been reported that continuous exposure and intoxication of liver to different
 83 types of exogenous compounds on a daily basis may lead to hepatic dysfunction [15]

84 Vitamin C (Ascorbic acid) is a water-soluble antioxidant occurring in the organism as an
 85 ascorbic anion. It also acts as a scavenger of free radicals and plays an important role in
 86 regeneration of α -tocopherol [25]. Ascorbic acid is a powerful antioxidant because it can donate
 87 a hydrogen atom and form a relatively stable ascorbyl free radical. As a scavenger of reactive
 88 oxygen and nitrogen oxide species, ascorbic acid has been shown to be effective against the
 89 superoxide radical ion, hydrogen peroxide, the hydroxyl radical and singlet oxygen [23]. Water-
 90 soluble vitamin C (ascorbic acid) also provides intracellular and extracellular aqueous-phase
 91 antioxidant capacity primarily by scavenging oxygen free radicals.

Vitamin E (α -tocopherol) is a fat-soluble vitamin known to be one of the most potent endogenous antioxidants. α -tocopherol is a term that encompasses a group of potent, lipid soluble, chain-breaking antioxidants that prevents the propagation of free radical reactions. Vitamin E is an important antioxidant, which is suggested to play a physicochemical role in the stabilisation of bio-membrane by virtue of lipid-lipid interaction between the vitamin and the unsaturated fatty acids [6]. Reports have shown that antioxidants like vitamin C and Vitamin E have shown protection against cadmium induced toxicity in different animal models [26][27]. Supplementation of these natural antioxidants reduced ROS levels, lipid peroxidation, haematological values and enzymatic and non-enzymatic components of antioxidant defense system.

Materials and Method

Extract preparation

2330g of Moringa seed was purchased from Mararaba market in Nasarawa State. The oil was extracted using the following procedures. The husks were removed from the seeds and they were were heated inside an oven at a temperature of 40°C and were pounded using a mortar and pestle in order to separate the chaff from the seeds. The seed extract was then grinded into powder using a grinding mill. The powdered extract of Moringa Oleifera was dissolved in 466ml of water at ambient temperature for two days and was later filtered. The aqueous extract was then poured into molten mesh and was placed on an oil extractor machine. The seed oil was removed at high temperature and pressure. 2000g of Cashew nut was purchased from Kuchikau in Nasarawa state. The oil was extracted using the following procedure. The nuts were heated inside an oven at a temperature of 40°C and were grinded into powder using a grinding mill. The

114 extract was poured into a molten mesh and placed in an oil extractor machine. The oil was
115 removed at high temperature and pressure and both oil **extract** were kept at room temperature.

116 **Drug Preparation**

117 The Cadmium Sulphate Solution ($3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$) was prepared by dissolving 9.9198mg of
118 Cadmium Sulphate salt (CdS) in 5ml of 0.9% w/v Phosphate buffer. The Ascorbic acid (Vitamin
119 C) was prepared by dissolving 5mg of Vitamin C in 10ml of 0.9% w/v Phosphate buffer. The
120 Vitamin E (Alpha Tocopherol) was prepared by dissolving 6mg of Vitamin E in 20ml of olive
121 oil.

122 **Experimental animals:**

123 Thirty five (35) Wistar rats **weighting** between 80g-180g were used for this research work. The
124 rats were randomly selected into seven groups as follow A, B, C, and D, E, F and G of five **(5)**
125 rats each. They were kept in the animal house of Bingham University, Nigeria and given feed
126 and water *ad libitum*. The treatment for the various groups was administered accordingly.

127 **Chemical and Extract Administration**

128 0.32ml (40mg/kg body weight) of Moringa Oleifera and Cashew nut seed oil, 0.16ml (20mg/kg
129 body weight) **each of Moringa oleifera seed oil and Cashew nut oil**, 0.8ml **(100mg/kg body**
130 **weight) of Vitamin C**, **0.3ml of Vitamin E (300mg/kg body weight)** and 5ml of 0.9% w/v of
131 Phosphate buffer were administered orally to the experimental rats according to their individual
132 groups for a period of four (4) weeks using a 2ml syringe with an oral cannular at the tip. The
133 administration was done by holding the rats with a glove using the left hand in such a way that
134 the neck region was being held so that the rats would be stable while the extracts were being
135 administered orally. The treatment was done every morning after which the animals were fed.

Animal Treatment

The animals were treated as shown below in the table

Table 1: Number of Animals in each group and Dosage of treatment given:

Groups	Number of Animals	Dosage
A	5	5ml of 0.9% w/v Phosphate buffer per kg bw
B	5	3.5mg/kg bw Cadmium only
C	5	100mg/kg Vitamin C (0.8ml) and 300mg/kg Vitamin E (0.3ml)
D	5	100mg/kg Vitamin C (0.8ml) and 300mg/kg (0.3ml) Vitamin E + 3.5mg/kg bw Cd
E	5	3.5mg/kg bw Cadmium+ 40mg/kg Moringa oleifera seed oil (0.32ml)
F	5	3.5mg/kg bw Cadmium+ 40mg/kg Cashew nut oil (0.32ml)
G	5	3.5mg/kg bw Cadmium+ Moringa oleifera seed oil + Cashew nut oil 20mg/kg each (0.16ml each).

Control Groups

The animals were grouped into three control groups which are; the normal control (A), the negative control (B) and the positive control (C). The Normal Control Group (A) received 5ml of 0.9% w/v Phosphate buffer orally for a period of four (4) weeks. The Negative Control Group (B) was induced intraperitoneally with 3.5mg/kg $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$.

145 The Positive Control Group (C) received 100mg/kg (0.8ml) Vitamin C and 300mg/kg
146 (0.3ml) Vitamin E orally for a period of four (4) weeks.

147 Prophylactic Treatment Group

148 The animals in Group D were the prophylactic group that received 100mg/kg (0.8ml)
149 Vitamin C and 300mg/kg (0.3ml) Vitamin E for a period of four (4) weeks followed by
150  intraperitoneal injection of 3.5mg/kg body weight $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$. The rats were then
151 sacrificed after 24 hours.

152 Therapeutic Treatment Groups

153 The animals in Groups E, F and G were the therapeutic control group. The animals in
154 each group were injected intraperitoneally with 3.5mg/kg $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$. After 48 hours
155 of the injection, the animals in Group E and F received 40mg/kg body weight of Moringa
156 Oleifera Seed Oil and Cashew Nut Oil for a period of four (4) weeks. The animals in
157 Group G received 20mg/kg each (0.16ml each) of both Moringa Oleifera Seed oil and
158 Cashew nut oil.

159 Animal Sacrifice

160 The animals were sacrificed twenty four (24) hours through cervical dislocation after which food
161 and water had already being withdrawn from them. The animals were laid in a supine position on
162 the dissecting board, the two hind limbs were held and the tail was pulled until a sound was held
163 which indicated that the animal was dead. The thoracic and abdominal cavities were exposed
164 adequately by using a surgical blade to make a midline incision through the skin of the
165 abdominal wall from the xiphisternum to the pubic symphysis. After the abdominal cavity had

166 been adequately exposed, the liver was removed and fixed in 10% formosaline for histological
167 analysis.

168 **Histological Preparation of Tissues**

169 The liver was fixed in 10% formosaline to preserve the cell and tissue in a life-like state
170 as much as possible and to prevent autolysis. The liver was passed through ascending
171 grades of alcohol (70%, 80%, 90% and absolute 100%) to gradually remove its water
172 contents. The sample was placed in a clearing agent which is xylene to remove the
173 alcohol. This improves the refractive index of the tissue. This is the immersion of the
174 tissue in molten paraffin wax so that the holes left by the alcohol would be filled up. This
175 gives the tissue support. The tissue is placed into an embedding mould which is filled
176 with more paraffin wax and allowed to solidify. This is done in order to make the tissue
177 compact for sectioning. The block is trimmed to remove the excess wax. The block of
178 tissue was placed in a microtome and trimmed to expose the surface. The microtome was
179 set to 3-5 micron and the tissue was sectioned. The sections were picked with forceps and
180 placed in a water bath to float out and spread well. It was picked with a slide and the slide
181 was placed on a hot plate in order for the tissue to slick to the slide. The slides were
182 arranged on a staining rack and dewaxed in two changes of xylene for 10 minutes. It was
183 then hydrated in descending grades of alcohol and rinsed. The stain was applied and the
184 slides left to dry. Photomicrographs were taken with a JVC colour video digital camera
185 (JVC, China) mounted on an Olympus light microscope (Olympus UK Ltd, Essex,UK) to
186 demonstrate the reticular fibres.

187



188

RESULT AND DISCUSSION




189

Group A

190

Photomicrograph of the liver of experimental control group received phosphate buffer



191

192

Fig. 1: Histological demonstration of the liver using silver staining techniques

193

[X400] showing the reticular fibres (RF) and portal vein (PV).

194

195

196

197


198

199

200 **Group B**

201 **Photomicrograph of the liver of experimental administered Cadmium only**

202 **(3.5mg/kg body weight).**



203

204 **Fig.2: Histological demonstration of the liver using silver staining techniques**

205 **[X400] showing the enlarged portal vein (EPV) and reticular fibre**

206

207

208


209

210

211 **Group C**

212 **Photomicrograph of the liver of experimental animal administered Vitamin C and**

213 **Vitamin E,**



214
215 **Fig.3: Histological demonstration of the liver using silver staining techniques**

216 **[X400]** showing the reticular fibres (RF) and **central vein (CV)**

217 **Group D**

218 **Photomicrograph of the liver of experimental animal administered Vitamin C and**

219 **Vitamin E and Cadmium.**




Fig.4: Histological demonstration of the liver using silver staining techniques [X400] showing the portal vein (PV) and degeneration of reticular fibres (DRF)

Group E

Photomicrograph of the liver of experimental animal administered Cadmium and Moringa oleifera oil




Fig.5: Histological demonstration of the liver using silver staining techniques [X400] showing the central vein (CV) and reticular fibres (RF).

Group F

Photomicrograph of the liver of experimental animal administered Cadmium and Cashew nut oil




Fig.6: Histological demonstration of the liver using silver staining techniques [X400] showing the portal vein

Group G

Photomicrograph of the liver of experimental animal administered Cadmium (3.5mg/kg body weight) and Moringa oleifera oil and Cashew nut oil




Fig.7: Histological demonstration of the liver using silver staining techniques [X400] showing the blood vessel (BV) and reticular fibres (RF).

Silver staining aids the visualization of the intracellular and extracellular components such as DNA and proteins such as type III collagen and reticulin fibres by the deposition of metallic silver particles on the target of interest [28].

Present study revealed normal histological architecture of the liver, the portal vein and the reticular fibres were seen clearly and there was orderly arrangement of the reticular fibres in group A. In group B, there was enlargement of the portal vein and distortion of the arrangement of the reticular fibres. This thus indicates hepatic dysfunction due to the generation of reactive oxygen species. In group C, resolution was observed and the reticular fibres were seen though not orderly arranged. This suggests that the antioxidant activities of the Vitamins can help to enhance the liver functions. In group D, the portal vein and reticular fibres were seen though not



255 well arranged and the sinusoidal beds were a little bit distorted. This suggests that the antioxidant
 256 activities of the Vitamins have helped to curb the effects of free oxygen radicals generated by
 257 Cadmium although not highly potent. This is supported by the previous research by Flora *et. al.*,
 258 [6] that antioxidant has been found to heal the free radical mediated cell damage. In groups E, F
 259 and G, there was restoration of the reticular fibres and the portal vein was seen unlike the group
 260 B which was induced with Cadmium only. This therefore suggests that Moringa oleifera oil and
 261 Cashew nut oil have ameliorative effects that led to the regeneration of the damaged and
 262 distorted reticular fibres. We therefore deduced from our finding that vitamin C, vitamin E,
 263 Moringa oleifera oil and Cashew nut oil attenuate the effects of cadmium on liver of wistar rats.

264 **Conclusion**

265 The histological architecture of the livers of the treated rats showed the central vein and
 266 restoration of the hepatocytes these show ameliorative effect of Moringa oleifera seed oil and
 267 Cashew nut oil on Cadmium induced liver damage.

268 **CONSENT**

269 It is not applicable.

270 **ETHICAL APPROVAL**

271 All the authors hereby declare that all the experiments have been examined and approved by the
 272 appropriate ethics committee and have therefore been performed in line with the ethical
 273 procedure laid down by Nigeria Medical Ethical Association for Accreditation of Laboratory
 274 Animal Care.

275

276

277 REFERENCES

278

279 [1] Achal (2002). Cashew: Nutrition and medicinal value. *Colarado state University*, pp:
280 159-165.

281 [2] Akinwale, T. O. (2000). Cashew apple juice: its use in fortifying the nutritional
282 quality of some tropical fruits. *Eur Food Res Technology*, **211**:205-207.

283 [3] Beyersmann, D and Hechtenberg S. (1997). Cadmium, gene regulation, and cellular
284 signalling in mammalian cells. *Toxicol Appl Pharmacol*, **144**:247–261.

285 [4] Faizi, S., Siddiqui, B.S, Saleem, R, Siddiqui, S, Aftab, K, Gilani, A.H (1995). Fully
286 acetylated carbamate and hypotensive thiocarbamate glycosides from *Moringa*
287 *oleifera*. *Phytochemistry*, **38**:957–963.

288 [5] Feuston, M.H and Scott, W.J Jr. (1985). Cadmium- induced forelimb ectrodactyly: a
289 proposed mechanism of teratogenesis. *Teratology*, **32**: 407-419.

290 [6] Flora S.J.S. (2002). Lead in the Environment: Prevention and Treatment. *J Environ*
291 *Biol*; **23**:29-44.

292 [7] Friberg, L., Nordberg, G.F, Vouk, V.B. eds. (1986). Handbook of the toxicology of
293 metals. Vol. II. Amsterdam, *Elsevie*; pp. 130–184.

294 [8] Fulgie, L.J. (1999). The Miracle Tree: *Moringa oleifera*: Natural Nutrition for the
295 Tropics. Church World Service; pp 172

296 [9] Gibbon, D and Pain, A. (1981). Crops of the drier regions of the tropics. *Longman*
297 *Singapore publications Ltd*, pp 66-67.

- 298 [10] Gilani, A.H., Janbaz, K.H and Shah, B.H. (1997). Quercetin exhibits
299 hepatoprotective activity in rats. *Biochem. Soc. Trans.* 25: 85.
- 300 [11] IARC (1976) *Cadmium, nickel, some epoxides, miscellaneous industrial chemicals*
301 *and general considerations on volatile anaesthetics*. Lyon, International Agency for
302 Research on Cancer, pp. 39–74.
- 303 [12] IARC, International Agency for Research on Cancer, Beryllium, cadmium, mercury,
304 and exposures in the glass manufacturing industry (1993). In: International Agency
305 for Research on Cancer Monographs on the Evaluation of Carcinogenic Risks to
306 Humans. *IARC Scientific Publications Lyon*, **58**:119–237.
- 307 [13] Lin, S., Sandow, M, Decandido, S, Waslien, C and Naughton, B. (1992). The effect
308 of vitamin D3 and 1,25- dihydroxy vitamin D3 on the toxic symptoms of cadmium
309 exposed rats. *Nutr. Res.* **12**:1393.
- 310 [14] Mustapha, Y; Hasfat, S. (2007). Antibacterial activities of *Anacardium occidentale*
311 leaf extract against some selected bacterial isolates. *IJPAS.*, 1(1):40-43.
- 312 [15] Nithya, N., Chandrakumar, K, Ganevan, V and Senthilkumar, S (2012). Efficacy of
313 *Momordica charantia* in attenuating abnormalities in cyclophosphamide
314 intoxicated rats. *J. Pharmacol. Toxicol.* **7**(1): 38-45.
- 315 [16] Ochi, T., Takahashi, K and Ohsawa, M. (1987). Indirect evidence for the induction
316 of a prooxidant state by cadmium chloride in cultured mammalian cells and a
317 possible mechanism for the induction. *Mutat. Res.* **180**:257–266.

- [17] Rao, V.A., Devi, P.U and Kamath, R. (2001). In vivo radioprotective effect of Moringa oleifera leaf extract on gastric ulcer models in rats. *Phytother Res*, **9**: 463–465
- [18] Ros, J.P.M and Slooff W, eds. (1987). Integrated criteria document. *Cadmium*. Bilthoven, National Institute of Public Health and Environmental Protection (Report No. 758476004).
- [19] Sharma P, Kumari P, Srivastava MM, Srivastava S (2006). Removal of cadmium from aqueous system by shelled Moringa oleifera Lam. seed powder. *Bioresour Technol*; **97**: 299–305.
- [20] Tsaknis, J.V and Spiliotis (1999). Quality changes of *Moringa oleifera*, variety Mbololo of Kenya, seed oil during frying. *Grasas y Aceites*, **50**(1): 37-48.
- [21] Ware, GW., ed. (1989). Cadmium. US Environmental Protection Agency Office of Drinking Water health advisories. *Reviews of Environmental Contamination and Toxicology*, **107**:25–37.
- [22] Webb., M. and In: Brown, S.S. (Ed.) (1977). Clinical Chemistry and Chemical Toxicology of Metals. Elsevier, Amsterdam; pp:51–64.
- [23]Weber, P., Bendich, A and Schalch (1996). Ascorbic acid and human health – a review of recent data relevant to human requirements. *Int J Vit Nutr Res*, **66**:19-30.
- [24] WHO (in preparation) (2010). Safety evaluation of certain food additives and contaminants in food. Geneva, World Health Organization. *WHO Food Additives Series*, No. 64.

- 340 [25] Young, I.S., Woodside, I.S. (2001). Antioxidants in health and disease. *J Clin*
341 *Pathol*, **54**: 176- 186.
- 342 [26] Ognjanovic, B.I., Pavlovic, S.Z, Maletic, S.D, Zikic R.V, Stajn A.S, Radojicic R.M,
343 Saicic Z.S, and Petrovic V.M. (2003). Protective influence of vitamin E on
344 antioxidant defense system in the blood of rats treated with cadmium. *Physiol Res*,
345 **52**:563-70.
- 346 [27] Beytut, E., Yuce, A, Kamiloglu, N.N and Aksakal M. (2003). Role of dietary
347 vitamin E in cadmium-induced oxidative damage in rabbit's blood, liver and
348 kidneys. *Int J Vitam Nutr Res*, **73** : 351-5.
- 349 [28] Schwint, O.A, Labraga, M, Cervino C.O, Haffar, M, Sequeiros, P.H and Marcos,
350 H.J. (2004).A modification of the staining technique of reticular fibres for image
351 analysis of the cardiac collagen network". *Cardiovasc. Patho.* **13**(4): 213- 20.
352