

The Nutraceuticals: A Voluminous Torrent in Pharmaceuticals- Coupling Health & Drugs

ABSTRACT

The present review is focused on the nutraceuticals which are present in our surroundings having an excellent impact over the health of humans but are not known for their pharmaceutical use. Nutraceuticals include vitamins, minerals and other dietary supplements, which may be herbs, enzymes, animal extracts etc. Various researchers have proved that nutraceuticals are having an important role in reducing the risk of various diseases such as diabetes, cardiovascular diseases, Parkinson's disease etc. with very few or negligible side effects.

Keywords: Nutrients, Drugs, Dietary supplements, Functional foods.

1. INTRODUCTION

"Let food be your medicine and medicine be your food" the great line said by great scientist Hippocrates is all related with the chemicals having nutritious and therapeutic value i.e. Nutraceuticals. Which states the value of nutraceuticals in **therapeutics**. [1]

Nutraceuticals are a multifarious product category which has a number of synonyms that are used internationally. The term "Nutraceutical" was given by Stephen De Felice who was the founder and chairman of the Foundation for Innovation in Medicine. This term has been a part of the industry lexicon for almost a decade. [2] The term "Nutraceutical" is derived from the combination of two words, "nutrient" which means a nourishing food or food component and second "pharmaceutical" which means a medical drug. Nutraceuticals may contain substances that are "natural" intended to treat or prevent number of diseases but may not be generally recognized as safe. Hence these are the food products intended for health and medical benefit. [3, 4] It have been proved by research that nutraceuticals are useful in providing protection from a number of diseases like diabetes, cancer, cardiac disease, hypertension etc e.g. carotenoids and anti-oxidants found in carrots help in avoiding chronic diseases, by preventing free radical damage. [5] **In the present scenario** Nutraceuticals have become more popular in modern society because of negligible chances of adverse effects of pharmaceuticals, ease of self-medication and aging in population.

2. TYPES OF NUTRACEUTICALS

2.1 Dietary Supplements

Dietary supplements (ds) have basic objective to provide **nutrition** which are otherwise not consumed in sufficient quantities like herbs, minerals, vitamins, or products obtained from plant sources, animal sources such as yeasts, fungus, algae, seafood and many more e.g. **Enzymes**, energy bars, amino acids, and liquid food supplements. **U.S. authorities** say that dietary supplements may be regarded as foods, while elsewhere they may be classified as drugs or other products. [6, 7]

2.2 Functional Foods

Japan introduced the concept of functional food in 1980s, to promote health or reduce the risk of diseases. The functional food include those food items which are advised to be consumed as part of the normal diet containing biologically active constituents offering the potential to enhance health or reduce

risk of various diseases. Various examples are food that comprises fatty acids, vitamins, specific minerals or dietary fibers, food with added biologically active substances such as phyto-chemicals or other antioxidants and probiotics that have live beneficial cultures. [8] Some examples of functional food products are; milk, cheese and eggs (enriched with omega-3 fatty acids); probiotics (yogurt enhanced with live active cultures); fruit juices and drinks (having antioxidant); cereals and grains such as wheat, oat, barley (having enriched amounts of dietary fibre); modified fatty acid vegetable oils; and soy, canola and hemp (vegetable proteins) , legumes and fruit products.[9, 10].

As per established requirement of functional food in Japan: food should be consumed-

- in its natural form, rather than a prepared dosage form like capsule, tablet, or powder;
- daily , in sufficient quantity ; and
- in a way which should be regulating a biological process to prevent or cure disease.[11]

2.3 Drugs, Dietary supplements and food additives

A drug is a substance used in diagnosis, cure, mitigation, treatment, or prevention of diseases, while dietary suppliments are any substances that are either deliberately added to food to enhance its shelf-life, nutrition, texture, or other aspect of quality or which unintentionally contaminates food (indirect additive) are known as food additives. (Figure 1) [12]




Figure 1: Diagrammatic Representation of Nutraceuticals Nutrients

Nutrients are the nutritive constituents present in food that a person consumes for his/her survival and growth. Macronutrients provide the bulk energy required for functioning of metabolic system, whereas micronutrients are helpful in providing the necessary cofactors for metabolism. Plenty of these nutrients are available in the environment. These nutrients are helpful in building and repairing of tissues, to regulate body processes. [13] Inorganic chemical compounds like water, minerals, and oxygen should also be considered as nutrients. [14]

2.4 Herbals

In ancient time; a large number of herbs were used to prevent and treat a number of diseases. A plant containing a number of non-nutritive phyto-chemicals provides health benefits if included in diet . [15, 16, 17] Nutraceuticals as herbals are big boon to human being in aspect of improving their health and to prevent them from chronic diseases e.g. willow bark (*Salix nigra*), helps as anti-inflammatory, analgesic, anti-arthritic, astringent as well as antipyretic. [18]

2.5 Phyto-chemicals

Phyto-chemicals are the plant components having bio-activities which are used to get health benefits. But their use always requires to be defended with some scientific rational in food, as potential nutraceutical. Phyto-chemicals are having following health benefits:

- (1) These are used as substrates for biochemical reactions.
- (2) These are used as cofactors of enzymatic reactions.
- (3) These work as inhibitors of enzymatic reactions.
- (4) These are used as absorbents/sequester which bind to unwanted constituents present in intestine and eliminate them.
- (5) These act as ligands which agonize or antagonize cell surface or intracellular receptors.
- (6) These are used as scavengers of various reactive or toxic chemicals.
- (7) These are used as compounds to increase the absorption and or improve stability of many essential nutrients.
- (8) These work as selective growth factors for gut friendly bacteria.
- (9) Fermented phyto-chemicals are beneficial for non-pathogenic bacteria found in GI tract.
- (10) These are selective inhibitors of deleterious intestinal bacteria. Phyto-chemicals which consists terpenoids, phenolics, alkaloids and fiber, are extensively examined and used for their ability to get health advantage. [19]

2.6 Probiotic/ Prebiotics

Probiotic bacteria are "living microbes taken in tolerable quantity for health advantage by recipient". These may interact with commensal bacteria to have a direct impact on the host. [20] Metchnikoff was the first person who successfully revamped the toxic flora of the large intestine into a host-sympathetic colony of *Bacillus bulgaricus*, found by Hord. [21, 22] Hence probiotics are gut friendly bacteria which aids in digestion and absorption of some nutrients. They act by eliminating the disease causing pathogens, like yeasts, other bacteria and viruses which mutually develop advantageous symbiosis within gastrointestinal tract. (Table 1) [23] The Japanese were the first to recognize the value of non-digestible oligosaccharides, and added these in feed of piglets to relieve and prevent from the diarrhoea. It was observed that fructo-oligosaccharides and galacto-oligosaccharides cause an increase in intestinal bifido-bacteria which stimulated their growth in the human gut. Hence a prebiotic is "a selectively fermented ingredient that causes changes in the activity of the gastrointestinal microbiota that are beneficial for human health". [24] A number of marketed nutraceuticals are available in market as mentioned in Table 2.

Table 1: Different species of microbes used as Probiotic

Genus	Species
Lactobacillus	acidophilus
	delbrueckii
	brevis

	fermentum
	gasseri
	johnsonii
	paracasei
	plantarum
	reuteri
	rhamnosus
	salivarius
Bifidobacterium	adolescentis
	animalisb
	breve
	bifidum
	infantis
	longum
Streptococcus	thermophilus
	salivarius
Saccharomyces	cerevisiae
Escherichia	coli
Enterococcus	faecium
Bacillus	coagulansc
	clausii

PRODUCTS	CATEGORY	CONTENTS	MANUFACTURER
Coral calcium	Calcium supplement	Calcium and trace minerals	Nature's answer, Hauppauge, NY, USA
Weight smart™	Nutritional supplement	Vitamins and trace elements	Bayer corporation, Morristown, NJ, USA
Omega woman	Immune supplement	Antioxidants, vitamins and phytochemicals (eg. Lycopene, and resveratrol)	Wassen, Surrey, U.K
Appetite Intercept™	Appetite suppressant	Caffeine, tyrosine and Phenylalanine	Natrol, Chatsworth, CA, USA
Chaser™	Hangover supplement	Activated calcium carbonate, and vegetable carbon	Living essentials, Walled lake MI, USA
Rox® Glucon-D Glucose-D	Energy drink	Taurine, caffeine and glucuronolactone Glucose	Rox America, Spartanburg, SA, USA Dabur
Mushroom optimizer™	Immune supplement	Mushrooms polysaccharides and Folic acid	Jarrow formulas, Los Angeles, CA, USA
Biovinca™	Neurotonic	Vinpocetine	Cyvox nutrition, Irvine, USA
Proplus®	Nutritional supplement	Soy proteins	Campbell soup company, Camden, NJ, USA
Snapple-aday™	Meal replacement beverage	Vitamins and minerals	Snapple beverage group, White Plains, NY, USA
WelLife®	Amino acid supplement	Granulated-L-glutamine	Daesang America Inc., Hackensack, NJ, USA
PNer plus™	Neuropathic pain supplement	Vitamin and other natural supplement	NeuroHelp, San Antonio, Texas, USA
Olivenol™	Dietary supplement	Natural antioxidant, Hydroxytyrosol	Cre Agri, Hayward, CA, USA
Threptin®	Diskettes Protein supplements	Proteins and vitamin B	Raptakos, Brett & Co. Ltd., Mumbai, India
GRD®	Nutritional supplement	Proteins, vitamins, minerals and carbohydrates	Zydus Cadila Ltd. Ahmedabad, India
Proteinex®	Protein supplement	Predigested proteins, vitamins, minerals and carbohydrates	Pfizer Ltd., Mumbai, India
Calcirol D-3®	Calcium supplement	Calcium and vitamins	Cadilla healthcare limited, Ahmedabad, India
Appetite Intercept	Appetite suppressant	Caffeine, tyrosine and phenylalanine	Natrol, Chatsworth, CA, USA
Betafactor™	Immune supplement	Beta glycan	Ameridan International Inc. USA
Brainspeed Memory®	Brain Health supplement	vitamin and minerals	Natrol, Chatsworth, CA, USA
Red bull®	Energy drink	Taurine, Caffeine, Glucuronolactone, b-group vitamins	Austrian red bull GmbH
5 hour energy®	Energy drink	Vitamins, tyrosin, Taurine, malic acid, caffeine, Glucuronolactone	Living essential, USA
Revital®	Daily health supplement	Ginseng, vitamin and minerals	Ranbaxy, India
Becadexamine®	Nutritional supplement	Multivitamins	GSK, India
Glowelle®	Beauty drink	Antioxidants, vitamins and fruit extracts	Nestle, India
Threptin® Diskettes	Protein supplement	Protein and vitamin B	Raptakos Brett & co. Ltd, India
HiOwna	Nutritional supplement	Protein, multivitamins, minerals and antioxidant	Himalaya herbal hdi
PediSure®	Nutritional supplement	Protein, multivitamins, minerals and antioxidant	Abbott India Ltd, India
Orgazyme®	Sex stimulating	L-arginine & L-ornithine	
R-Gene® 10	Growth stimulant	L- arginine HCl	Pfizer Inc, NY
Alamin SE®	Protein supplement	L-arginine & other Protein	Albert David Ltd., India
Albumen Care	Protein supplement	L-arginine & other Protein with multivitamins and minerals	B.V. Bio-Corp Pvt. Ltd., India
Argipreg	Protein Supplement	L-arginine, proanthocyanidins	Manikind Pharma Pvt. Ltd, India

Table 2: List of marketed nutraceuticals

3. PHARMACOLOGICAL USE OF NUTRACEUTICALS

Better life quality is achieved with food items filled with nutrient values and remarkably potent in preventing diseases which may be cancer, diabetes, heart diseases, hypertension etc as . (Table 3 & Figure 2) Such products may be dietary supplements, food additives, phyto-chemicals, prebiotics, probiotics, genetically modified food, processed food and plant products.




Figure 2: Pharmacological Uses of Nutraceuticals

Table 3: Detailed Review on Various Nutraceuticals

Types of Nutraceuticals	Sources	Active Constituents	Applications
Dietary Fibre	Whole grain foods wheat and corn bran, nuts	Insoluble Fibre	Reduce chances of colon or breast cancer (anticancer) ^[74, 75] , maintain health of digestive tract ⁷⁶
	Oats, barley	Beta-Glucan	Reduce risk of cardiovascular disease, lower down Low Density Lipids and total cholesterol ^[77,78]
	Beans e.g. Legumes, oats, barley and some fibrous fruits	Soluble Fibre	Anticancer (Colon Cancer), Digestive ^[79,80]
Fatty Acids	Salmon and other fish oils	Long chain omega-3 Fatty Acids-DHA/EPA	Reduce risk of CVD Improve mental, visual functions ⁸¹⁻⁸³
	Cheese, meat products	Conjugated Linoleic Acid (CLA)	Improving of body composition Decreases chances of certain cancers ⁸⁴⁻⁸⁶
Phenolics	Fruits	Anthocyanidins	Antioxidant ; reduce risk of cancer ⁸⁷⁻⁸⁹
	Green Tea	Catechins	Antitumor ⁹⁰
	Citrus	Flavonoids	Antioxidative activity, Prevention of coronary heart disease, hepato-protective, Effective in inflammation and cancer. ⁹¹
	Cocoa, Chocolate, Cranberries & cranberry products	Tannins	Anti-microbial, Reduce risk of cardiovascular disease ⁹²
	Corn, soy, wheat, wood oils	Plant Sterols, Stanol ester	Lower blood cholesterol levels by inhibiting cholesterol absorption ⁹³
Carotenoids	Tomatoes	Lycopene	Antioxidant, protect against prostate cancer ⁹⁴
	Corn, various fruits, egg yolk, spinach	Lutin	Antioxidant, Muscle regeneration, anti cancer activity, protect eyes against age related muscular degenerations, cataract ^{95,96}

	Carrots, various fruits (Guava, papaya, Water melon etc) and vegetables (tomatoes etc)	Beta carotene	Antioxidant, protection of cornea against UV light.
	Soya beans	Saponins	Effective against colon cancer, reduces cholesterol level ⁹⁷
Probiotics/ Prebiotics	Curd	Lactobacillus	Antibacterial, acute diarrhea ⁹⁸
	Whole grains, onions, combination of Pro & Prebiotics	Fructo–oligosaccharides	Improve GI health, restore gut flora ⁹⁹
Phytochemical	Grains	Tocotrienols and tocopherols	The growth of diverse tumors cell lines was suppressed via initiation of apoptosis and concomitant arrest of cells in the G1 phase of the cell cycle ¹⁰⁰
	Cereal grain, dairy & egg products and plants oil	Phytosterols	Exhibit antioxidant, anti-inflammatory, anti-neoplastic, anti-pyretic & immune-modulating activity, decrease cholesterol ¹⁰¹⁻¹⁰³
	Various plants, wholegrain	Phenolic constituents	Antioxidants, Anti-hyperglycemic, and anti hypertensive ¹⁰⁴
	Grapes, berries, cocoa, green tea, acacia spp.	Catechin & gallic acids	Antioxidants, Antiradical property, cyto-protective. ^{105,106}
	Soybeans	Isoflavonoids	Treating cancers & attenuates bone loss ^{107, 108}

129

130 3.1 In cardiovascular disease

131 It is not easy to set up a clear impact of nutrition/physical exercise on major cardio vascular diseases
132 because history of cardiovascular diseases is too long. [25] The effect of calcium over hypertension and
133 pre-eclampsia (a condition in pregnancy characterized by high blood pressure, sometimes with fluid
134 retention and proteinuria) is unpredictable as well as ambiguous. [26] Treatment with vitamin C and
135 selenium need further study to observe its effect on mortality. [27] It has been observed that some

nutraceuticals may be useful to forbid and superintend the risk of thrombosis in women with thrombophilic gene mutations. [28]

Vitamins, minerals, omega-3 poly-unsaturated fatty acids (n-3 PUFAs), dietary fibers and antioxidants, as nutraceuticals and physical exercise are advised to prevent and treat cardio vascular diseases. Researchers have proved that polyphenols found in grapes and in wine are helpful in reducing arterial disease by altering cellular metabolism and signaling. [29]

Onion, black grapes, cherries, cruciferous vegetables, grapefruits, red wine, apples and berries are good sources of flavonoids [30] and also available as flavones and flavonols which are beneficial for the treatment of cardiovascular diseases.[31, 32, 33] Antioxidant activity of ascorbic acid, alpha-tocopherol, and beta-carotene as have been studied and reviewed.[34]

3.2 In management of diabetes

Various *in-vitro* and *in-vivo* studies (animal) have proved that plant polyphenols including phenolic acids, stilbenes, lignans and flavonoids are effective nutraceuticals in diabetes and its prevention. Although human clinical trials are required to check the efficacy of poly-phenol compounds in treatment of diabetes. [35]

Vitamin C (ascorbic acid) is a chain-breaking antioxidant which avoids the propagation of chain reactions that may lead to a reduction in protein glycation. It has been reported that ascorbic acid helps in reducing diabetes-induced sorbitol in animals. [15]. Ascorbic acid (800 mg/day) partially replenishes ascorbic acid levels in patients with type 2 diabetes but endothelial dysfunction or insulin resistance is not effected at all. [16].

N. Bunyapraphatsara *et al* observed a combined effect of Aloevera juice with glibenclamide in diabetic patients and concluded that glibenclamide alone did not show any effect while Aloevera juice showed significant reduction in fasting blood glucose level and triglycerides within two weeks and four weeks respectively. Although it did not show any effect on level of cholesterol but the effectiveness of Aloe vera was found in the treatment of diabetes. [36] Acacia is a non-starch polysaccharide which is not digested in the intestine, but generate short chain fatty acids in large bowel; hence, it produces extensive biological effects. Philips AO *et al* conducted study over the extract of Acacia Arabica and confirmed the anti-diabetic effect of acacia resulted by increasing the insulin release. [37] Hou *et al* demonstrated significant hypoglycemic effects of Acacia Arabica powder in healthy rabbits, whereas no significant decrease in blood sugar was observed in the alloxan-induced diabetic rabbits was observed. [38] Wadood *et al* concluded that Acacia arabica initiates the release of insulin from pancreatic beta cells of normal rabbits. [39] Antidiabetic activity of glycerrhiza in non insulin dependent diabetic model was observed by Takii. [40]

High intake of isoflavone (20–100 mg/day) is helpful in lowering of rate of mortality in diabetes of type 2, osteoporosis, cardiac disease and certain cancers.[41] Docosahexaenoic acid is vital for neurovisual development which helps in modulating insulin resistance and nurture the advocacy for essential fatty acids in pregnancy in women with gestational diabetes mellitus. [42]

Omega-3 fatty acids helps in reducing blood glucose tolerance in patients predisposed to diabetes. Insulin is needed to synthesize long chain n-3 fatty acids; thus heart may be victim to their depletion in case of diabetes. Hence it is concluded that ethyl esters of n-3 fatty acids may be advantageous in diabetic patients. [43] Lipoic acid is a well known antioxidant, and in Germany being used to cure diabetic neuropathy. Lipoic acid; as a long-term dietary supplement is aimed at the prophylactic protection of diabetics from complications. [44] α -Lipoic acid enhances insulin sensitivity by approximately 18–20% in patients suffering from type 2 diabetes. [45] Clinical trial studies on α -lipoic acid reported advantageous in the treatment of diabetic neuropathy.[46]

Dietary fibers obtained from psyllium have been used to reduce weight and lipid levels in hyperlipidemia as dietary supplement. [47] Intake of Chromium supplements may be helpful to enhance sensitivity to insulin and boost glucose tolerance in type II diabetic patient. [48] Magnesium-rich diet intake may reduce risk of diabetes by improving insulin sensitivity.[49] Diabetes management is supported by one of the nutraceuticals Biotin which increase insulin production and stimulates liver glucokinase activity, thus improves the uptake of glucose in muscle cells. [50, 51, 52] Pharmacological effect of epinephrine can be inhibited by Azadirachta Indica which results in enhanced utilization of peripheral glucose [53, 54] and reduce hypo-glycaemic activity without change in the serum cortisol level. [55, 56]

Kernels of *Eugenia jambolana* are useful in diabetes management; their aqueous/alcoholic extract shows hypoglycemic effect. [57] Green Tulsi (*Ocimum sanctum*) leaves extract also reduces blood sugar significantly by cortisol inhibiting potency as proved in both normal and alloxan induced diabetic rats. [56, 58]

3.3 In Parkinson's disease

Latif S *et al* concluded that diet enriched with vitamin E may decrease the chances of Parkinson's disease [59] while Brower V reported that creatine is helpful in management of Parkinson's disease by decreasing the clinical symptoms. [60]

Antioxidant vitamin supplements such as tocopherol, ascorbic acid and beta-carotene are the abundantly occurring nutraceuticals. As per various earlier literature vitamin E supplements are becoming popular in treatment of Parkinson's disease, whereas epidemiological studies reported that vitamins C and E rich diets are associated with decreased risk of Parkinson's disease. [61, 62]

3.4 In Alzheimer's disease

Literatures have been reported that fulvic acid, an active principle of Shilajit is highly effective against brain disorders exclusively and in combination with vitamin B complex. [63] Treatment of patient suffering from Alzheimer's disease with donepezil and vitamin E found effective although future study was suggested to check and compare additive as well as individual effect. [64] Wettstein A. *et al* reported that mild to moderate Alzheimer's dementia could be treated with metrifonate, donepezil, rivastigmine which are second-generation cholinesterase inhibitors. [65]

Hager *et al* found stable cognitive function especially in those patients who were administered with 600 mg Alpha-lipoic acid along with acetyl-cholinesterase inhibitors, in comparison to the patient only on the therapy of standard acetyl-cholinesterase inhibitors since last 337 days. [66] Huperzine-alpha is a plant alkaloid derived from Club moss plant, Huperzine serrata, which is a member of the Lycopodium species. Huperzine-alpha is in phase III clinical trial in the USA and is available as a dietary supplement.[67] The meta-analysis of Huperzine A reported here highlights that this treatment has certain significant improvement for patients with Alzheimer's disease and Vascular Dementia, and longer durations may result in better efficacy for patients with Alzheimer's disease. [68]

Literature survey supports that Lipoic acid also helps to improve potential of mitochondrial membrane, memory loss due to ageing and brain ailments as well as in patient suffering from Parkinson's and Alzheimer's disease. [69]

3.5 In Erectile Dysfunctioning

L-arginine in combination with pycnogenol (a product obtained from the pine bark of *Pinus pinaster*) is found safe and effective in mild to moderate erectile dysfunction in Japanese patients. [70] When Patients suffering from moderate to severe ED and dyslipidemia were kept on Niacin rich diet, significant improvement was observed in patients. [71]

Kaempferia Parviflora Wall. Ex. Baker (KP), which is a Thai plant with name, Kra-Chai-Dum and rhizomes of which are used as a traditional medicine to alleviate male impotency, improve male libido, as energizer, control blood pressure and also reduce stomachache. Study reveals that KP is a potential nutraceutical compound effective in male erectile dysfunction caused due to ageing. [72]

3.6 Osteoarthritis

Chondroitin sulfate and Glucosamine both are commonly used to lighten the symptoms of osteoarthritis. Chondroitin sulfate and Glucosamine both act as nutraceuticals and possible mechanism of their anti- inflammatory activity may be due to synthesis of NO and PGE2.[73]

Conclusion

In the present scenario Nutraceuticals have become more popular in modern society and became important due to increasing applications of Nutraceuticals which serves as a part of growing pharmaceutical industry. Modern society is now being aware about the food product that is beneficial for them in aspects of health and nutrition owing to very few or no side effects. Nutraceuticals are being used to treat fatal diseases like cancer, diabetes, cardiovascular diseases; as well as Parkinson's disease, osteoarthritis etc.

REFERENCES

1. Bagchi D. Nutraceuticals and functional foods regulations in United States and around the world. Toxicol. 2006; 221: 1-3.
2. Granato H. Regulatory Concerns Cloud Functional Food, Nutraceutical Markets. c 2009 <http://www.naturalproductsinsider.com/articles/2000/11/regulatory-concerns-cloud-functional-food-nutraceutical.aspx>
3. Ross S. Functional foods: the Food and Drug Administration perspective. Am J Clin Nut. 2000; 71: 1735-38.
4. Palthur, MP, Palthur SSS, Chitta SK. Nutraceuticals: Concept and regulatory scenario. Int J Pharm Pharm Sci. 2010; 2: 14-20.
5. Cindy, Oliveri S. Nutraceuticals, Phytochemicals, and Antioxidants-What Are They All About. OSU Extension Fact Sheet. 2003; 5051-98
6. Dietary Supplements What You Need to Know, USFDA, May 2006 <http://www.fda.gov/downloads/Food/DietarySupplements/UCM240978.pdf> (last cited on Jan 20, 2016).
7. Dietary supplement health and education act 1994, public law 103-417, 103rd congress http://ods.od.nih.gov/pdf/About/DSHEA_Wording.aspx (last cited on Jan 20, 2016).
8. <http://www.eufic.org/article/en/expid/basics-functional-foods/> (last cited on Jan 20, 2016).
9. Hathcock J. Dietary supplements: How they are used and regulated. J Nut. 2001; 131: 1114-7.
10. Dureja H, Kaushik D, Kumar V. Developments in nutraceuticals. Ind J Pharmacol. 2003; 35: 363-72.
11. Hardy G. Nutraceuticals and functional foods: introduction and meaning. Nutrition. 2000; 16(7-8): 688-9.

- 262 12. Camire ME, Kantor MA. Dietary Supplements: Nutritional and Legal Considerations, Food technol.
263 1999; 53(7): 87-96.
- 264 13. Whitney, Elano, Sharo R. Understanding nutrition. 10th ed. Thomson-Wadsworth; 2005.
- 265 14. Frances S, Ellie W. Nutrition: concepts and controversies.1^{3th} ed. Cengage learning 2007.
- 266 15. Hamid AA, Luan YS. Functional properties of dietary fiber prepared from defatted rice bran. Food
267 Chem. 2000; 68: 15 - 9.
- 268 16. Mermel V. Old paths new directions: use of functional foods in the treatment of obesity. Trends Food
269 Sci Tech. 2004; 15: 532-40.
- 270 17. Dulloo AG, Duret C, Rohrer D. Efficacy of a green tea extract rich in catechin polyphenols and
271 caffeine in increasing 24-h energy expenditure and fat oxidation in humans. Am J Clin Nutr. 1999; 70:
272 1040-5.
- 273 18. Ehrlich SD. (Willow bark), private practice specializing in complementary and alternative medicine
274 Phoenix AZ Review. VeriMed Healthcare Network, 2008.
- 275 19. Cora JD, German JB. Phytochemicals: nutraceuticals and human health. J Sci Food Agric, 2000;
276 80:1744-56.
- 277 20. FAO/WHO (2001) Joint FAO/WHO Expert Consultation on evaluation of health and nutritional
278 properties of probiotics in food including powder milk with live lactic acid bacteria Cordoba, Argentina,
279 October 2001.
- 280 21. Hord NG. Eukaryotic microbiotic crosstalk: potential mechanisms for health benefits of prebiotics and
281 probiotics. Annu Rev Nutr. 2008; 28: 215–31.
- 282 22. Michail S, Sylvester F, Fuchs G, Issenma R. Clinical efficacy of probiotics: Review of the evidence
283 with focus on children, clinical practice guideline. J Pediatr Gastroenterol Nutr. 2006; 43(4):550-7.
- 284 23. Holzapfel WH, Haberer P, Geisen R, Bjorkroth J, Schillinger U. Taxonomy and important features of
285 probiotic microorganisms in food and nutrition. Am J Clin Nutr. 2001; 73: 365S-73S.
- 286 24. Gibson GR. Dietary prebiotic: current status and new definition. IFIS Func Foods Bull. 2011; 7: 1–19.
- 287 25. Ignarro LJ, Balestrieri ML, Napoli C. Nutrition, physical activity, and cardiovascular disease: an
288 update. Cardio Res. 2007; 73: 326-40.
- 289 26. Trumbo PR, Ellwood KC. Supplemental calcium and risk reduction of hypertension. Nutr Rev. 2007;
290 65: 78-87.
- 291 27. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Mortality in randomized trials of
292 antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis.
293 JAMA. 2007; 297: 842-57.
- 294 28. Ravi SMT. Nutrigenetics and nutraceuticals: the next wave riding on personalized medicine. Trans
295 Res. 2007; 149:55-61.
- 296 29. German JB, Walzem RL. The health benefits of wine. Ann Rev Nutr. 2000; 20: 561-93.
- 297 30. Hollman PCH, Hertog MGL, Katan MB. Analysis and health effects of flavonoids. Food Chem. 1996;
298 57:43- 6.

- 299 31. Yao LH, Jiang YM, Shi J, Tomás-Barberán FA, Datta N, Singanusong R, Chen SS. Flavonoids in
300 food and their health benefits. *Plant Foods Hum Nutr.* 2004; 59(3): 113-22.
301
- 302 32. Cook NC, Samman S. Flavonoids - Chemistry, metabolism, cardio-protective effects, and dietary
303 sources. *J Nutri Biochem.* 1996;7:66-76.
- 304 33. Hollman PCH, Feskens EJ, Katan MB. Tea flavonols in cardiovascular disease and cancer
305 epidemiology. *Proc Soc Exper Biol Med.* 1999; 220:198-202.
- 306 34. Niki E, Noguchi N, Tsuchihashi H, Gotoh N. Interaction among vitamin C, vitamin E, and beta-
307 carotene. *Am J Cli Nutr.* 1995; 62(6Suppl):1322S-6S.
308
- 309 35. Zahra B, Parvin M, Fereidoun A. Dietary polyphenols as potential nutraceuticals in management of
310 diabetes: a review. *J Diabetes Metab Disord.* 2013; 12(1):43.
- 311 36. Bunyapraphatsara N, Yongchaiyudha S, Rungpitarangsi V, Chokechaijaroenporn O. Antidiabetic
312 activity of Aloe vera L juice II Clinical trial in diabetes mellitus patients in combination with
313 glibenclamide. *Phytomed.* 1996;3(3): 245-8.
- 314 37. Philips AO, Philips GO. Biofunctional behavior and health benefits of a specific Gum Arabic. *Food*
315 *hydrocoll.* 2011; 25(2): 165-9.
- 316 38. Hou CC, Lin SJ, Cheng JT, Hsu FL. Antidiabetic dimeric guianolides and a lignan glycoside from
317 *Lactuca indica*. *J Nat Prod.* 2003; 66(5):625-9.
- 318 39. Wadood A, Wadood N, Shah SA. Effects of *Acacia arabica* and *Caralluma edulis* on blood glucose
319 levels of normal and alloxan diabetic rabbits. *J Pak Med Assoc.* 1989; 39(8): 208-12.
- 320 40. Takii H, Kometani T, Nishimura T, Nakae T, Okada S, Fushikii T. Antidiabetic effect of glycyrrhizin in
321 genetic diabetes *Licorice* in diabetes. *Biol Pharma Bull.* 2000; 24: 484-7.
- 322 41. Brouns F. Soya isoflavones: a new and promising ingredient for the health foods sector. *Food Res*
323 *Int.* 2002; 35: 187-93.
- 324 42. Thomas B, Ghebremeskel K, Lowy C, Crawford M, Shore R N. Nutrient intake of women with and
325 without gestational diabetes with a specific focus on fatty acids. *Nutrition.* 2006; 22: 230-6.
- 326 43. Sirtori CR, Galli C. Fatty acids and the Omega 3. *Biomed Pharmacother.* 2002; 56: 397-406.
- 327 44. Coleman MD, Eason RC, Bailey CJ. The therapeutic use of lipoic acid in diabetes: a current
328 perspective. *Envir Toxi Pharmacol.* 2001; 10: 167-72.
- 329 45. Singh U, Jialal L. Alpha-lipoic acid supplementation and diabetes. *Nutr Rev.* 2008;66: 646–57.
- 330 46. Ziegler D, Reljanovic M, Mehnert H, Gries FA. Alpha-lipoic acid in the treatment of diabetic
331 polyneuropathy in Germany: Current evidence from clinical trials. *Exp Clin Endo Dia.* 1999;107: 421–
332 30.
- 333 47. Baljit S. Psyllium as therapeutic and drug delivery agent. *Int J Pharm.* 2007; 334:1-14.
- 334 48. Lau FC, Bagchi M, Sen CK, Bagchi D. Nutrigenomic basis of beneficial effects of chromium (III) on
335 obesity and diabetes. *Mol Cell Biochem.* 2008; 3:171–10.
- 336 49. McCarty MF. Nutraceutical resources for diabetes prevention—an update. *Med Hypo.* 2005; 64:151–
337 8.

- 338 50. Dakshinamurti K, Cheah-Tan C. Biotin-mediated synthesis of hepatic glucokinase in the rat. Arch
339 Biochem Biophys. 1968; 127: 17-21.
- 340 51. Furukawa Y. Enhancement of glucose-induced insulin secretion and modification of glucose
341 metabolism by biotin. Nippon Rinsho. 1999; 57: 2261-9.
- 342 52. Wang ZQ, Zhang XH, Cefalu WT. Chromium picolinate and biotin enhance glycogen synthesis and
343 glycogen synthase gene expression in human skeletal muscle culture. Diab Res Clin Pract. 2000; 50:
344 395.
- 345 53. Chattopadhyay RR, Chattopadhyay RN, Nandy AK, Poddar G, Maitra SK. The effect of fresh leaves
346 of *Azadirachta indica* on glucose uptake and glycogen content in the isolated rat hemi diaphragm.
347 BullCal Sch Trop Med. 1987; 35: 8–12.
- 348 54. Chattopadhyay RR. Possible mechanism of antihyperglycemic effect of *Azadirachta indica* leaf
349 extract part IV. Gen Pharmacol. 1996; 27:431–4.
- 350 55. Chattopadhyay RR. A comparative evaluation of some blood sugar lowering agents of plant origin. J
351 Ethnopharmacol. 1999;67:367–72.
- 352 56. Gholap S, Kar A. Hypoglycaemic effects of some plant extracts are possibly mediated through
353 inhibition in corticosteroid concentration. Pharmazie. 2004;59:876–8.
- 354 57. Sharma SB, Nasir A, Prabhu K,M Murthy PS, Dev G. Hypoglycaemic and hypolipidemic effect of
355 ethanolic extract of seeds of *Eugenia jambolana* in alloxan-induced diabetic rabbits. J
356 Ethnopharmacol. 2003;85:201–6.
- 357 58. Rai V, Iyer U, Mani UV. Effect of *Tulsi* (*Ocimum sanctum*) leaf powder supplementation on blood
358 sugar levels serum lipids and tissue lipids in diabetic rats. Plant Foods Hum Nutri. 1997; 50:9–16.
- 359 59. Latif S, Anwar F, Ashraf M, Gilani AH. *Moringa oleifera*: a food plant with multiple medicinal uses.
360 Phytother Res. 2007; 21:17-25.
- 361 60. Brower VA. Nutraceutical a day may keep the doctor away. EMBO reports. 2005; 8:708- 11.
- 362 61. Maxwell SRJ. Antioxidant vitamin supplements: update of their potential benefits and possible
363 risks. Drug Saf. 1999; 21:253–66.
- 364 62. Anderson C, Checkoway H, Franklin GM et.al, Dietary factors in Parkinson's disease: the role of food
365 groups and specific foods. Mov Disord. 1999; 14:21–7.
- 366 63. Carlos C, Gonzalo F, Patricio F, Fernando C, Ricardo M. Can Nutraceuticals Prevent Alzheimer's
367 Disease? Potential Therapeutic Role of a Formulation Containing Shilajit and Complex B Vitamins.
368 Arch Med. 2012; 43(8): 699–704.
- 369 64. Klatte ET, Scharre DW, Nagaraja HN, Davis RA, Beversdorf DQ. Combination therapy of donepezil
370 and vitamin E in Alzheimer disease Alzheimer. Dis Asso Disord. 2003; 17: 113-6.
- 371 65. Wettstein A, Stadtärztlicher DZ. Cholinesterase inhibitors and Gingko extracts--are they comparable
372 in the treatment of dementia? Comparison of published placebo-controlled efficacy studies of at least
373 six months' duration. Phytomedicine. 2000; 6(6): 393-401.
- 374 66. Hager K, Marahrens A, Kenklies M, Riederer P, Munch G. Alpha-lipoic acid as a new treatment option
375 for Alzheimer type dementia. Arch Gerontol Geriatr. 2001;32:275-82.

- 376 67. Grunwald J, Raveh L, Doctor BP. Huperzine A as a pretreatment candidate drug against nerve agent
377 toxicity. *Life Sci.* 1994; 54: 991-7.
- 378 68. Shu-huai X, Chun-Xiao Z, Rui Z, Li An. Huperzine A in the Treatment of Alzheimer's Disease and
379 Vascular Dementia: A Meta-Analysis. *Evid Based Complement Alternat Med.* 2014 Article ID 363985
380 [10 http://dxdoiorg/101155/2014/363985](http://dxdoiorg/101155/2014/363985)
- 381 69. Kramer K, Elkins MHR. Alpha lipoic acid: Nature's supreme Antioxidant. In: Elkins MHR. USA: Wood
382 land Publishing; 1998.
- 383 70. Aoki H, Nagao J, Ueda T, Strong JM, Schonlau F, Yu-Jing S, Lu Y, Horie S. Clinical assessment of a
384 supplement of Pycnogenol and L-arginine in Japanese patients with mild to moderate erectile
385 dysfunction. *Phytother Res.* 2012; 26(2): 204-7.
- 386 71. Ng CF, Lee CP, Ho AL, Lee VW. Effect of niacin on erectile function in men suffering erectile
387 dysfunction and dyslipidemia. *J Sex Med.* 2011; 8(10): 2883-93.
- 388 72. Yenjai CK, Prasanphen S, Daodee V, Wongpanich, Kittakoop P. Bioactive flavonoids from
389 *Kaempferia parviflora*. *Fitoterapia.* 2004; 75(1):89-92.
- 390 73. Kalioraa AC, Dedoussisa GVZ, Schmidt H. Dietary antioxidants in preventing atherogenesis.
391 *Atherosclerosis.* 2006; 187(1):1-17.
- 392 74. Adom KK, Liu RH. Antioxidant activity of grains. *J Agric Food Chem.* 2002; 50 (21): 6182-7.
- 393 75. Ferguson LR, Harris P. Protection against cancer by wheat bran: role of dietary fibre and
394 Phytochemicals. *Euro J Can Pre.* 1999; 8(1): 17-25.
- 395 76. Engberg RM, Hedemann MS, Steinfeldt S, Jensen BB. Influence of Whole Wheat and Xylanase on
396 Broiler Performance and Microbial Composition and Activity in the Digestive Tract. *Poul Sci.* 2004;
397 83(6): 925-38.
- 398 77. Keogh GF, Cooper GJS, Mulvey TB, McArdle BH, Coles GD, Monro JA, Poppitt SD. Randomized
399 controlled crossover study of the effect of a highly β -glucan-enriched barley on cardiovascular
400 disease risk factors in mildly hypercholesterolemic men. *Am J Clin Nutr.* 2003; 78(4): 711-8.
- 401 78. Theuwissen E, Mensink RP. Water-soluble dietary fibers and cardiovascular disease. *Physiol*
402 *Behav.* 2008;94(2):285-92.
- 403 79. McIntyre A, Gibson PR, Young GP. Butyrate production from dietary fibre and protection against large
404 bowel cancer in a rat model. *Gut.* 1993; 34: 386-91.
- 405 80. Malkki Y, Virtanen E. Gastrointestinal effect of oat bran and oat gum: a review. *LWT- Food Sci*
406 *Technol.* 2001;34(6):337-47.
- 407 81. Popp-Snijders C, Schouten JA, Heine RJ, Vander MJ, Vander VE. A Dietary supplementation of
408 omega-3 polyunsaturated fatty acids improves insulin sensitivity in non-insulin-dependent diabetes.
409 *Diabetes Res.* 1987; 4(3):141-7.
- 410 82. Connor WE, Importance of n-3 fatty acids in health and disease. *Am J clin Nutr.* 2000; 71(1): 171S-
411 5S.
- 412 83. Young G, Conquer J. Omega-3 fatty acids and neuropsychiatric disorders. *Reprod Nutrition Dev.*
413 2005; 45(1): 1-2 8.

- 414 84. Lee KN. Kritchevsky D, Parizaa MW. Conjugated linoleic acid and atherosclerosis in rabbits.
415 Atherosclerosis. 1994;108(1):19-25.
- 416 85. Clement I, Banni S, Angioni E, Carta G, McGinley J, Thompson HJ, Barbano D, Bauman D.
417 Conjugated Linoleic Acid–Enriched Butter Fat Alters Mammary Gland Morphogenesis and Reduces
418 Cancer Risk in Rats. J Nutr. 1999;
419 129(12): 2135-42.
- 420 86. Shultz TD, Chewab BP Seamana WR, Luedeck LO. Effect of conjugated linoleic acid on body
421 composition in mice. Cancer Lett. 1992; 63(2):125–33.
- 422 87. Wang H. Cao G. Prior RL. Oxygen Radical Absorbing Capacity of Anthocyanins. J Agric Food Chem.
423 1997;45(2):304–9.
- 424 88. Kanga SY, Seeramb NP, Narib MG, Bourquin LD. Tart cherry anthocyanins inhibit tumor
425 development in ApcMin mice and reduce proliferation of human colon cancer cells. Cancer Lett.
426 2003;194(1):13–9.
- 427 89. Li-Shu W, Stoner GD. Anthocyanins and their role in cancer prevention. Cancer Lett.
428 2008;269(2):281–90.
- 429 90. Liang G, Tang A, Lin X, Li L, Zhang S, Huang Z, Tang H, Li QQ. Green tea catechins augment the
430 antitumor activity of doxorubicin in an in vivo mouse model for chemoresistant liver cancer. Int J
431 Oncol. 2010; 37(1):111-23.
- 432 91. Kumar S. Pandey AK. Chemistry and Biological Activities of Flavonoids: An Overview. Scientific
433 World J. 2013; 1-16.
- 434 92. Lim H, Darah I, Jain K. Antimicrobial activities of tannins extracted from rhizophora apiculata barks. J
435 Trop Forest Sci. 2006; 18(1):59—65.
- 436 93. Katan MB, Grundy SM, Jones P. Efficacy and Safety of Plant Stanols and Sterols in the Management
437 of Blood Cholesterol Levels. Mayo Clin Proc. 2003; 78(8):965–78.
- 438 94. Takeoka GR, Dao L, Flessa S, Gillespie DM, Jewell WT, Huebner B, Bertow D, Ebeler
439 SE. Processing Effects on Lycopene Content and Antioxidant Activity of Tomatoes. J Agric Food
440 Chem. 2001; 49(8):3713–17.
- 441 95. Naguib YMA. Antioxidant Activities of Astaxanthin and Related Carotenoids. J Agric Food
442 Chem. 2000; 48(4):1150–54.
- 443 96. Olmedilla B, Granado F, Blanco I, Vaquero M, Cajigal C. Lutein in patients with cataracts and age-
444 related macular degeneration: a long-term supplementation study. J Sci Food Agri. 2001;81(9):904–
445 9.
- 446 97. Kim HY, Yu R, Kim JS, Kim YK, Sung MK. Antiproliferative crude soy saponin extract modulates the
447 expression of Ikb α protein kinase C and cyclooxygenase-2 in human colon cancer cells. Cancer Lett.
448 2004; 210(1):1–6.
- 449 98. Isolauri E, Rautanen T, Juntunen M, Sillanaukee P, Koivula T. A Human Lactobacillus Strain
450 (Lactobacillus Casei sp strain GG) Promotes Recovery From Acute Diarrhea in
451 Children. Pediatric.1991; 88(1):90 -7.
- 452 99. Fuller R, Gibson GR. Probiotics and prebiotics: microflora management for improved gut health.
453 Clinic Microbiol Infec. 1998; 9:477–80.

- 454 100. Ju J, Picinich SC, Yang Z. Cancer-preventive activities of tocopherols and tocotrienols.
455 Carcinogenesis. 2010; 31(4):533–42.
- 456 101. Wang T, Hicks KB, Moreau R. Antioxidant activity of phytosterol oryzanol and other phytosterol
457 conjugates. J Ame Oil Chem Soci. 2002; 79(12):1201-6.
- 458 102. Rozner S, Garti N. The activity and absorption relationship of cholesterol and phytosterols.
459 Colloids and Surfaces A: Physicochem Engg Aspec. 2006; 282-283:435-56.
- 460 103. Garcia MD, Saenz MT, Gomez MA. Topical anti-inflammatory activity of phytosterol isolated from
461 Ezyngium foetidum on chronic and acute inflammatory model. Phytother Res.1999; 13(1):78-80.
- 462 104. Ranilla LG, Kwon Y, Apostolidis E, Shetty K. Phenolic compounds antioxidant activity and in vitro
463 inhibitory potential against key enzymes relevant for hyperglycemia and hypertension of commonly
464 used medicinal plants herbs and spices in Latin America. Bioresour Tech. 2010; 101(12):4676–89.
- 465 105. Morel I, Lescoat G, Cogrel P, Sergent O, Pasdeloup N. Antioxidant and iron-chelating activities of
466 the flavonoids catechin quercetin and diosmetin on iron-loaded rat hepatocyte cultures. Biochem
467 Pharmacol. 1993; 45(1):13-9.
- 468 106. Kroes BH, Vanden BAJ, Quarlesvan UHC, Labadie RP. Anti-inflammatory activity of gallic acid.
469 Planta Medica. 1992; 58(6):499-504.
- 470 107. Birt DF, Hendrich S, Wang W. Dietary agents in cancer prevention: flavonoids and isoflavonoids.
471 Pharmacol Thera. 2001; 90(2–3):157–77.
- 472 108. Alekel DL, St G, Peterson CT, Hanson KB, Stewart, Toda T. Isoflavone-rich soy protein isolate
473 attenuates bone loss in the lumbar spine of perimenopausal women. Am J Clin Nut. 2000; 72(3):844-
474 52.