

ADVERSE HAEMATOLOGICAL CHANGES IN PLASMODIUM INFECTED MICE FOLLOWING TREATMENT WITH AZADIRACHTA INDICA (NEEM) LEAF EXTRACT

Abstract

Background: *Azadirachta indica* (neem) is an effective natural remedy for malaria treatment in some tropical regions of Africa. This study investigated the impact of *Azadirachta indica* leaf extract on some haematological indices of *Plasmodium berghei-berghei* infected mice. **Materials and Methods:** Twenty-five (25) mice weighing between 19kg and 31kg were randomly selected into five (5) groups of 5 animals per group. Groups 1, 2 and 3 were inoculated on day 0, intraperitoneally about 1% (4.5×10^4) *Plasmodium berghei berghei* parasitized red blood cells. Groups 1 and 2 were treated with 50 mg/kg and 75mg/kg of the extract respectively by oral gavage, while group 3 animals were given normal saline. Groups 4 and 5 were not infected but group 4 was treated with 75mg/kg of the extract and 5 served as normal control mice. Treatment lasted for 14 days. Whole blood was obtained for haematological analysis using Sysmsec® Automated Haematology Analyzer KX-21N. **Results:** The result showed significant reduction in red blood cell count (RBC: $p = 0.000$), packed cell volume (PCV: $p = 0.000$) and haemoglobin concentration (HB: $p = 0.000$) of *P. berghei berghei* infected mice following *A. indica* leaf-extract administration. The mean cell haemoglobin (MCH: $p = 0.006$) and mean cell volume (MCV: $p = 0.024$) were significantly higher when compared to uninfected and untreated control mice. The red cell indices (RBC count, PCV and HB) animals that were infected without extract treatment were significantly higher than those of infected mice that received the extract. Extract treatment to uninfected mice, however, did not show any significant change in these parameters when compared with untreated control animals. Total white blood cell count (WBC: $p = 0.005$), neutrophils ($p = 0.024$) and lymphocytes ($p = 0.001$) were significantly higher in infected treated mice than those without extract treatment. Administration of extract to uninfected mice did not cause any significant change in total white blood cell counts and the differential counts. However, eosinophil ($p = 0.214$) and platelet ($p = 0.152$) counts of infected mice receiving extract treatment were not significantly different from those of uninfected control. **Conclusion:** We therefore conclude that treatment of malaria with extract of *A. indica*, though may be effective as anti-malaria and could prevent malaria associated platelet changes, may contribute to malaria associated anaemia. Care should, therefore, be taken to follow up the extract treatment with blood building agents.

Key words: *Azadirachta indica*, Red cell indices, White blood cells, Platelet counts, *Plasmodium* malaria mice.

INTRODUCTION

Changes in haematological indices resulting in diverse degrees of anaemia are common findings in many disease conditions including malaria, hookworm infestation, chronic inflammatory and immune disorders, cancer and chronic renal disease. The mechanisms of induction of anaemia vary with the disease conditions. The pathophysiologic mechanism for

malaria-induced anaemia and other complications involve destruction of infected erythrocyte, liberation of parasite and erythrocyte materials into circulation and the host reactions to these events¹. Once in the blood stream, plasmodium invades liver and red blood cells and makes more copies of itself. Eventually, as red cells break and free plasmodium to infect other red cells and as the body's immune system works to kill infected cells, the total number of red blood cells drops causing anaemia².

Also, plasmodium infected erythrocytes sequester in microcirculation of vital organs interfering with micro-circulatory flow and host tissue metabolism. The infected red blood cells rosette and stick to endothelial cells lining the blood vessels³. Several proteins including plasmodium falciparum erythrocyte membrane protein I (PFEMP1) form knobs on the surface of the infected red cells and bind to ligands including CD36 thrombospondin, VCAM-1, ICAM-1 and E-selections displayed on the surface of endothelial cells⁴. Ischaemia results from impaired circulatory blood flow in the affected vessels supplying vital organs, thus leading to most of the manifestations seen in severe malaria. Merozoite surface antigens released from infected red cells induce the production of TNF- α , interferon (IFN) and IL-1, by the host cells and these cytokines suppress the production of red cells, increase fever and induce nitric oxide syntheses leading to tissue damaged and increased expression of endothelial receptors for PfEMP-1⁵.

Azadirachta indica (Neem) a large green tree of the family maliaceal, is believed to originate from **Assan** and Burma in South Asia⁶. It is one of the most useful medicinal plants used traditionally as remedies for various diseases⁷. It has been speculated by traditional medicine practitioners to be effective for treatment of activities, leprosy, malaria, typhoid fever, respiratory disorders, constipation, cancer and chronic syphilis sores. Scientific studies have also demonstrated some specific pharmacological properties of neem leaf-extract; for instance,

hypolipadaemic, antioxidant and anti-diabetes activities of the leaves extract have been reported in diabetes rat models^{8,9,10}. Immunostimulatory^{11,12}; anti-inflammatory^{13,14} and hepatoprotective¹⁵ activities have also been shown in various experimental models. Traditional use of neem leaf-extract for the treatment of malaria had long been observed and some experimental evidences of its antimalarial activities have been documented^{16,17}.

Although vector control and use of chemoprophylates agents such as chloroquine, quinine, amodiaquine. Sulfadoxine, pyrimethamine and artemisinin, either individually or in combination, are common preventive/curative measures against malaria infection; attention is also being focused on the use of effective and safe natural remedies as alternative strategies. This has been encouraged with a view to bringing malaria treatment closer to the poor rural dwellers of the African societies who can hardly afford or access conventional anti-malaria drugs; and to cope the emergence of high rate of drug resistant strain of malaria parasite and cases of adverse drug reactions. *Azadirachta indica* (neem) is one of such popular natural remedies for malaria treatment in tropical regions of Africa, and it has been shown to be effective in this regard. However, the effects of the plant extracts on some blood parameters have not been closely examined. This study therefore is aimed at investigating the impact of *Azadirachta indica* leaf-extract on some haematological indices of *plasmodium berghei-berghei* infected mice model.

MATERIAL AND METHODS

Collection and preparation of plant materials

The fresh leaves of neem (*Azadirachta indica*) were harvested from the Endocrine Research farm, University of Calabar, Calabar. The samples were taken to the Department of Pharmacognosy, University of Uyo, Uyo, where it was authenticated. The fresh leaves of neem

were collected and selected to remove the dead ones. Leaves were washed with distilled water and sun-dried to remove the water. The leaves were blended using an electric blender and soaked in ethanol for 24 hours. The mixture was filtered and the residues were discarded. The filtrate was concentrated in a water bath at 50°C.

Determination of Extract Concentration by Gravimetric Method

The stock solution was stirred to mix. 2mls aliquot was drawn from the stock solution and oven dried in a beaker of known weight at 50°C. The beaker now containing the residue was then reweighed. The difference in weight of the empty beaker and when it contained the residue indicated the weight of the residue.

Experimental Animals

Thirty (30) mice were obtained from Pharmacology and Toxicology departmental animal house, University of Uyo. They were caged in a well ventilated room with 12 hours light-dark cycle. They were fed on commercial diet (Guinea Feed Ltd, Benin) and were also given tap water for 7 days during which they acclimatized.

Induction of Malaria and Parasite inoculation

Fifteen (15) mice weighing between 19kg and 31kg were randomly selected for induction of malaria, they were assigned to three groups with 5 animals per group. Chloroquine sensitive *plasmodium berghei-berghei* (animal parasite that causes the malaria) obtained from the National Medical Research Institutes, Abuja were maintained in mice. The parasite donor mice contained 5% (22.5×10^4) *plasmodium berghei-berghei* parasitized erythrocytes. The parasite donor blood was diluted with normal saline in proportion such that each mouse was inoculated on day 0, intraperitoneally about 1% (4.5×10^4) *plasmodium berghei berghei* parasitized red blood cells¹⁸. The animals were confirmed to be infected with malaria 72 hours after the inoculation.

111 **Experimental Groups**

112 Group 1: 5 infected animals receiving 50mg/kg of the extract orally.

113 Group 2: 5 infected animals receiving 75mg/kg of the extract orally.

114 Group 3: 5 malaria infected animals that did not receive any extract.

115 Group 4: 5 normal animals receiving 75mg/kg of the extract orally.

116 Group 5: 5 normal animals receiving no extract.

117 **Collection of samples for haematological analysis**

118 At the end of the 14 days, they were then euthanized under chloroform vapour and sacrificed.

119 Whole blood was collected via cardiac puncture using sterile syringes and needles. The whole

120 blood was emptied into EDTA containers preparatory to haematological (full blood count)

121 analysis. Full blood counts including PCV, Hb, RBC, WBC, platelet count, differential WBC

122 (lymphocytes and mixed), and red cell indices (MCHC, MCH and MCV), were estimated using

123 the Sysmex® Automated Haematology Analyzer KX-21N, Sysmex Corporation, Kobe-Japan.

124 The pre-diluted (PD) sample method was used where blood was diluted manually, and then fed

125 into the transducers. The transducer chamber has a minute hole called the aperture. On both sides

126 of the aperture, there are electrodes between which flows direct current. Blood cells suspended in

127 the diluted sample pass through the aperture, causing direct current resistance to change between

128 the electrodes. As direct current changes, the blood cell size is detected as electric pulses. Blood

129 cell count is then calculated by counting the pulses, and a histogram of blood cell sizes is plotted

130 by determining the pulse sizes. Also, analyzing a histogram makes it possible to obtain various

131 analysis data including differential whole blood count, red cell indices and derived values.

132

133

Statistical analysis:

Data are presented as Mean $\pm$ standard deviation. Differences between means were compared employing student's t-test and ANOVA post hoc, a probability of $p < 0.05$ was considered significant.

Results

Haematological changes associated with administration of *A. indica* leaf-extract to *plasmodium berghei berghei* infected mice were investigated. The effects of neem leaf - extract treatment of infected and uninfected mice on red blood cell indices are shown in table 1. The result showed that administration of *A. indica* leaf-extract to *P. berghei berghei* infected mice caused a significant reduction in some red blood cell indices namely: red blood cell count (RBC: $p = 0.000$), packed cell volume (PCV: $p = 0.000$) and haemoglobin concentration (HB: $p = 0.000$) while mean cell haemoglobin (MCH: $p = 0.006$) and mean cell volume (MCV: $p = 0.024$) were significantly higher when compared to uninfected and untreated control mice. Animals that were infected without receiving treatment with the extract also showed significant decrease in RBC count ($p = 0.004$), PCV ($p = 0.006$) and HB concentration ($p = 0.005$) but not in MCH ($p = 0.329$) and MCV ($p = 0.357$) when compared with control. However, the red cell indices (RBC count, PCV and HB) in this group were significantly higher than those of infected mice that received the extract. Extract treatment to uninfected mice did not show any significant change in these parameters when compared with untreated control animals.

Changes in white blood cell indices following *A. indica* leaf-extract treatment of infected and uninfected mice were presented in table 2. Total white blood cell count (WBC: $p = 0.005$),

neutrophils ($p = 0.024$) and lymphocytes ($p = 0.001$) were significantly higher in infected mice that received extract treatment than those without treatment. Administration of extract to uninfected mice did not cause any significant change in total white blood cell counts and the differential counts. However, eosinophil ($p = 0.214$) and platelet ($p = 0.152$) counts of infected mice receiving extract treatment were not significantly different from those of uninfected control but eosinophil ($p = 0.000$) was rather higher and platelet ($p = 0.001$) was lower in infected mice that received no treatment. Significant positive correlation existed between parasite-induced eosinophilia and total white blood cell count ($r = 0.993$, $p = 0.001$), neutrophil ($r = 0.898$, $p = 0.050$) and lymphocytes ($r = 0.994$, $p = 0.004$) in plasmodium infected mice that were not treated with extract. Significant negative correlation was also observed between eosinophil and platelet counts ($r = 0.818$, $p = 0.007$) in this group of mice.

Table 1: Changes in red cell indices of plasmodium infected and uninfected mice following treatment with *A. indica* leaf-extract.

Group	Pcv (%)	Hb g/l	RBC ($\times 10^{12}/L$)	MCHC (g/L)	MCH (pg)	MCV (fL)
1:Malaria + 50 mg/kg <i>A. indica</i>)	15.40 $\pm$ 2.23*	51.40 $\pm$ 1.12*	1.34 $\pm$ 0.11*	343.22 $\pm$ 0.38	38.82 $\pm$ 7.79*	113.14 $\pm$ 23.37*
2:Malaria infection + 75 mg/kg	15.0 00 $\pm$ 2.00*	50.40 $\pm$ 7.13*	1.34 $\pm$ 0.12*	335.89 $\pm$ 9.58	37.70 $\pm$ 4.53*	112.16 $\pm$ <u>13.69*</u>
3:Malaria infection only	24.00 $\pm$ 0.71 ^{*a}	79.20 $\pm$ 1.30 ^{*a}	2.66 $\pm$ 0.15 ^{*a}	330.28 $\pm$ 5.96	29.86 $\pm$ 1.60	90.54 $\pm$ 6.34
4:Normal mee + 75mg/kg <i>A.indica</i>	35.80 $\pm$ 3.84	119.40 $\pm$ 5.57	4.06 $\pm$ 0.24	333.72 $\pm$ 66.92	29.54 $\pm$ 1.35	88.40 $\pm$ 3.90
5:Normal mice (control)	37.60 $\pm$ 1.14	125.40 $\pm$ 3.91	4.42 $\pm$ 0.14	334.18 $\pm$ 1.23	28.46 $\pm$ 0.86	86.48. $\pm$ 2.31

Data are expressed as mean $\pm$ SD. n = 5 mice per group. * significant difference at $p < 0.05$ compared with control. ^a significant difference at $p < 0.05$ compared with extract treated infected mice.

176 Table 2: Changes in white blood cells and platelets of plasmodium infected and uninfected mice
177 following treatment with *A. indica* leaf-extract.

178

Group	WBC x 10 ³ /μL	Neut x 10 ³ /μL	Lymp x 10 ³ /μL	Eosin x 10 ³ /μL	Platelets x 10 ³ /μL
1: Malaria + 50 mg/kg <i>A. indica</i>)	22.54 ± 0.95*	4.56 ± 0.24*	18.88 ± 1.22*	0.24 ± 0.020	116.20 ± 8.14
2: Malaria infection + 75 mg/kg	28.04 ± 3.77*	6.16 ± 0.84*	21.88 ± 2.93*	0.25 ± 0.038	130.40 ± 6.58
3: Malaria infection only	14.80 ± 1.35* ^a	2.32 ± 0.25* ^a	12.42 ± 1.03* ^a	0.59 ± 0.056* ^a	95.20 ± 6.72* ^a
4: Normal mice + 75mg/kg <i>A.indica</i>	8.26 ± 1.29	1.88 ± 0.47	7.10 ± 1.07	0.21 ± 0.021	128.80 ± 6.22
5: Normal mice (control)	6.42 ± 1.78	1.24 ± 0.55	5.50 ± 2.42	0.20 ± 0.03	128.40 ± 6.19

179 Data are expressed as mean ± SD. n = 5 mice per group. * significant difference at p < 0.05
180 compared with control. ^a significant difference at p < 0.05 compared with extract treated infected
181 mice.

182
183 **DISCUSSION**

184
185 Unusual haemolytic reactions are known to occur in individual following administration
186 of different antimalarial drugs. Agents such as quinine, chloroquine, primaquine fansidar and
187 coartem are quite notable among others¹⁹. Other studies have also demonstrated dose dependent
188 increase in blood methaemoglobin concentration and increasing osmotic fragility of erythrocytes
189 following administration of fansidar, halofanthrine, quinine, chloroquine and coartem^{20,21}). All
190 these drugs were shown to generate rapid accumulation of reactive oxygen species (ROS) that
191 overwhelmed the antioxidant defense capacities of the erythrocyte and causes depletion of
192 erythrocyte glutathione concentration and red cell lysis. Also, plasmodium infected erythrocytes
193 are subsequently destroyed in the lymphoid tissues by cytotoxic T-lymphocytes and natural killer
194 cells as blood is passing through these organs thereby contributing to malaria associated
195 anaemia².

In this present study, we found that treatment of *Plasmodium berghei berghei* infected mice with leaf extract of *Azadirachta indica* caused significant reduction in red blood cell indices namely; total red cell count, PCV, HB concentration, and increase in MCH and MCV compared to uninfected mice that received no extract. Also infected mice that received no treatment showed a less severe anaemia than those treated with the extract while uninfected mice which were treated with the extract did not show significant change in the red cell indices compared to normal mice that received no treatment. These findings suggest that treatment of infected mice with *A. indica* leaves may enhance selective destruction of infected red blood cells in the affected mice groups while the uninfected red cells were spared.

A. indica is a well-known medicinal plant with scientific evidence of antimalarial activities¹⁷. The mechanism of its antiplasmodial activities is not fully understood, but had been shown to demonstrate significant immune-stimulatory and anti-inflammatory activities^{11,12}. Acetone-water neem leaf extract having antimalarial activity had been shown to inhibit the adhesion of malaria parasite-infected erythrocytes to endothelial cells¹⁷. Therefore, prevention of cytoadherence of infected erythrocyte to vascular endothelium increase the infected red cell contact with the cytotoxic T-lymphocyte and natural killer cells in the lymphoid tissue, thus resulting in increased destruction of parasitised cells and reduced parasite asexual reproduction in red cells. The extract is known to have significant anti-inflammatory activity as demonstrated by its ability in suppression of neutrophils and macrophages activation¹⁴. The anti-inflammatory effect can also be related to decreased expression of adhesion molecules by neutrophils and vascular endothelium and hence decreased cytoadherence.

The pathologic process of malaria infection and its complications involves cytoadherence and sequestration in micro circulation of vital organs of the body. Erythrocytes

220 containing mature form of plasmodium express several surface proteins (including PfEM-PI)
 221 which form kobs with ability to adhere to microvascular endothelium and thus disappear from
 222 the circulation, remaining stuck until they rupture at merogony (schizogony) to exacerbate the
 223 infection by infecting new red blood cells. The consequences of microcirculatory obstruction are
 224 activation of the vascular endothelial cells and reduced oxygen and substrate supply, leading to
 225 anaerobic glycolysis, acidosis and cellular dysfunction^{2,3}.

226 The mechanism by which *A. indica* leave extract causes severe anaemia in plasmodium
 227 infected mice, may therefore be due to decreased cytoadhesion of parasitized red cells to
 228 endothelium. This is particularly likely because administration of the extract to non-infected
 229 mice did not cause anaemia and the levels of anaemia in untreated plasmodium infected mice
 230 were not as severe as those treated with the extract. As indicated by the present study, the
 231 anaemia caused by malaria parasites alone without extract treatment is normochromic and
 232 normocytic in nature which is characteristic of mild haemolysis. However, treatment of the
 233 infected mice with *A. indica* leaf extract presented a macrocytic and slight hyperchronic picture
 234 as evident by increase MCV and MCH. This may be due to increased release into circulation of
 235 reticulocytes as the body is responding to the severe anaemic condition by increasing production
 236 of new red bloods cells.

237 We also observed that the white blood cells increased significantly in infected mice that
 238 were treated with the extract than in those that were not treated and those that were not infected
 239 by plasmodium but treated with the extract. This suggest an enhance activation and proliferation
 240 of the immune cells by the extract following antigence challenge such as plasmodium infection.
 241 Baral and Chattopadhyay¹² had shown that *Neem* leaf extract mediated higher immune
 242 activation. Activated lymphocyte (including T- lymphocyte and Natural killer cell) in the

lymphoid organ and in circulation could contribute to the excessive and selective destruction of red cells infected by the parasite. Most conventional anti-malarial drugs cause red cell destruction by increased generation of ROS and depletion of anti-oxidant defense molecules¹⁹, this mechanism might not generally be true for *A. indica* extract because earlier report had shown that the extract has significant anti-oxidant activities probably owing to its high concentration of polyphenols, such as flavonoids¹⁰, but may be related to its ability to inhibit cytoadherence of infected cells to endothelium.

The eosinophil counts of infected mice receiving extract treatment were not significantly different from those of control mice but the untreated infected mice should significantly higher eosinophil count relative to control. In the untreated mice, the increase in eosinophil may be a normal body's response against parasitic infection. However, the administration of neem leaf-extract ameliorates this effect by preventing activation and proliferation of the eosinophil through the same mechanism as indicated by Kaur and his colleagues¹⁴. This may further be supported by the significant positive correlation observed in this study between parasite-induced eosinophilia and total white blood cell counts, neutrophil and lymphocytes in plasmodium infected mice that were not treated with the extract.

Platelet activation and increased tendency for intravascular coagulation is characteristic of plasmodium malaria. Our studies showed increased platelet consumption in untreated infected mice as evident by significant decrease in platelet number compared to control mice. However, infected mice that were treated with the extract did not show significant reduction in platelet number. This finding suggests that the extract can prevent platelet activation and aggregation which can go a long way to reducing malaria associated coagulation complications. The anti-inflammatory property of the leaf extract may be responsible for this platelet affects.

We therefore conclude that treatment of malaria with extract of *A. indica*, though quite effective as anti-malaria and could prevent malaria associated bleeding disorders, may contribute to malaria associated anaemia and care should be taken to follow up the extract treatment with blood building agents.

References

1. David PH, Hommel M, Miller LH, Udeinya IJ, Oligino LD. Parasite sequestration in *Plasmodium falciparum* malaria: spleen and antibody modulation of cytoadherence of infected erythrocytes. *Proc Natl Acad Sci USA* 1983; 80: 5075–5079.
2. Flick K, Chen Q. var genes PfEMP1 and human host. *Mol Biochem Parasitol.* 2004; 134(1):3 – 9
3. Rowe JA, Moulds JM, Newbold CI, Miller LH. *Plasmodium falciparum* rosetting mediated by a parasite-variant erythrocyte membrane protein and complement-receptor 1. *Nature* 1997; 388: 292-295.
4. Baruch DI, Gormely JA, Ma C, Howard RJ, Pasloske BL. *Plasmodium falciparum* erythrocyte membrane protein 1 is a parasitized erythrocyte receptor for adherence to CD36, thrombospondin, and intercellular adhesion molecule 1. *Proc Natl Acad Sci USA* 1996; 93: 3497- 3502
5. McAdams AJ, Sharpe AH, Infectious Diseases: Malaria. In: *Pathologic Basis of Disease*. Kumar V, Abbas AK, Fausto N. (Eds.) 7th ed. Sounders Company, Philadelphia. 2004; pp 343-414.
6. National Research Council. *Neem: a tree for solving global problem* National Academy Press, Washington D.C., 1992.
7. Kausik BI, Ranajet KB. Biological activities and medicinal properties of neem plant. *Curr Sci.* 2002; 82: 1336–45.
8. Halim EM. Lowering of blood sugar by water extract of *Azadirachta indica* and *Abroma augusta* in diabetic rats. *Indian J. Exp. Biol.* 2003; 41(6): 636-640.
9. Gupta S, Kafaria M, Gupta PK, Murganandan S, Yashroy RC. Protective role of extracts of neem seeds in diabetes caused by streptozotocin in rats. *J. Ethnopharmacol* 2004; 90(2-3): 185-189.
10. Ekaidem IS, Akpan HD, Usoh IF, Etim OE, Ebong PE. Effects of ethanolic extract of *Azadirachta indica* leaves on lipid peroxidation and serum lipids of diabetic Wistar rats. *Acta. Biologica. Szegedensis* 2007; 51(1): 17-20.

- 307 11. Sen P, Medinata PK, Ray A. Immunostimulant activities of *Azadirachta indica*. Indian J.
308 Exp. Biol. 1992; 12: 1170-1175.
- 309
- 310 12. Baral RN, Chattopadhyay U. Neem (*Azadirachta indica*) leaf mediated immune
311 activation causes prophylactic growth inhibition of murine Ehrlich carcinoma and B16
312 melanoma. Int Immunopharmacol 2004; 4: 355-366. (PMID: 15037213)
- 313
- 314 13. Chattopadhyay RR, Chattopadhyay RN, Maitra SK. Possible mechanism of anti-
315 inflammatory activity of *Azadirachta indica* leaf extract. Indian J. Pharmacol. 1993; 25:
316 99-100.
- 317
- 318 14. Kaur G, Sarwar AM, Ather M. Nimbidin suppresses functions of macrophages and
319 neutrophils: relevance to its anti-inflammatory mechanisms. Phytother. Res. 2004; 18(5):
320 419-424.
- 321
- 322 15. John AA, Bamidele FP, oluwaseun OA. Some Protective Effects of Aqueous Leaf
323 Extract of *Azadirachta indica* on Paracetamol-induced Hepatotoxicity in Adult Wistar
324 Rats American J. Trop. Med.& Pub.Health 2011; 1(3): 97-106.
- 325
- 326 16. Omar S, Zhang J, MacKinnon S, Leaman D, Durst T, Philogene BJ, Arnason JT,
327 Sanchez-Vindas PE, Poveda L, Tamez, P.A., Pezzuto, J.M. Traditionally used
328 antimalarials from the Meliaceae. Curr Trop Med Chem. 2003; 3: 133-139.
- 329
- 330 17. Udeinya IJ, Mbah AU, Chijioke CP, Shu EN. An antimalarial extract from neem leaves is
331 antiretroviral. Trans. Res. Soc. Trop.Med. Hyg 2004; 98(7): 435-437.
- 332
- 333 18. de Ridder J. Malaria culture Manual, 2003
- 334
- 335 19. Anaba FC, Ahiane BO, Pepple DJ. In vitro hemolytic effect of
336 sulfadoxine/pyrimethamine and artemether/ lumefantrine on malaria parasitized
337 erythrocytes of female patients Pak. J. Pharm. Sci. 2012; 25 (4):851-855
- 338
- 339 20. Chikezie PC, Chikezie CM, Amaragbulem PI. Effect of antimalarial drugs on
340 polymerization of sickle cell hemoglobin (HbS). Turk. J. Biochem. 2010; 35: 41-44.
- 341
- 342 21. Chikezie PC, Uwakwe AA. Membrane stability of sickle erythrocytes incubated in
343 extracts of three medicinal plants: *Anacardium occidentale*, *Psidium guajava* and
344 *Terminalia catappa*. Pharm. Mag. 2011; 7: 121-125.
- 345