

**EVALUATION OF TOXICOLOGICAL EFFECTS OF ORAL ADMINISTRATION OF
CADMIUM NITRATE ON LIVER OF ADULT WISTAR RATS**

ABSTRACT

Available evidence suggest that adverse effect of health are caused by heavy metal and cadmium nitrate(CdN) has been shown to cause adverse health effect worldwide and liver is one of the organs mostly affected. The present study evaluates toxic effects of cadmium nitrate on liver of adult wistar rats.

Twenty (n=20) adult wistar rats of both sexes randomly divided into four groups (A-D) of five (n=5) rats each; Rats in group A served as control and were given 10ml/kg/day of distilled water for 21days. Rats in group B, C, and D (cadmium nitrate CdN group) were given 150mg/kg/bw of CdN, 225mg/kg/bw of CdN and 300mg/kg/bw of CdN administered orally through an orogastric cannula into the stomach via the esophagus, once a day respectively, for twenty one (21) consecutive days. The gross anatomical parameters of the liver and liver histology were assessed. An assessment of the histological profiles of the liver showed distortions of the liver cytoarchitecture following consumption of CdN. Body weights of experimental rats decreased as compared to the control group. The mean liver weight also decreased as compared to the control. The serum levels of AST and ALT were significantly increased in CdN group compared to the control group. It was conclude that cadmium nitrate induced liver toxicity in adult wistar rats.

KEY WORDS; Cadmium Nitrate, Liver, weight, Heavy metal, Wistar Rat

INTRODUCTION

Environmental pollution has many facets, and the resultant health risks include diseases in almost all organ systems [16][17]. Humans whose position in the food chain is at the top were exposed to various types of environmental contaminants at different stages of life, majority of which are harmful. In the environmental community, the notation of heavy metals implies stable high-density metals. These elements are natural constituents of the earth's crust. As a result of

anthropogenic activity, the input of heavy metal to the environment has increased sufficiently and resulted in the increase of their content in air, water, soil and tissues of living organisms. Heavy metals are mostly toxic but their exact toxicity varies depending on the chemical form of the metal, point of exposure and the degree of exposure to the metal [17]. Metals when concentrated can be quite toxic and can result in death of organisms. Numerous hazardous heavy metals are inhaled and absorbed by humans and animals every day [12]. The major concern with heavy metals is their ability to accumulate in the environment and thereby passing up to food chain [8].

Cadmium (cd) is a vital component of modern technology, with many applications in the electronics, communication, power generation and aerospace industries [18]. Cd compound are commercially produced. Production and consumption of these compounds are increasing particularly in china, Korea, and Japan. The worldwide production of cd was approximately 17,200 tons in 2004 [14]. Once taken up by the blood, the majority of cadmium is transported bound to proteins, such as Albumin and Metallothionein.[3].

The first organ reached after uptake into the GI-blood is the liver. Here cadmium induces the production of Metallothionein. After consecutive hepatocyte necrosis and apoptosis, Cd-Metallothionein complexes are washed into sinusoidal blood. From here, parts of the absorbed cadmium enter the entero-hepatic cycle via secretion into the biliary tract in form of Cadmium-Glutathione conjugates. Enzymatically degraded to cadmium-cysteine complexes in the biliary tree, cadmium re-enters the small intestines [19].

The main organ for long-term cadmium accumulation is the kidney [11]. Here the half-life period for cadmium is approx. 10 years. A life-long intake can therefore lead to a cadmium accumulation in the kidney, consequently resulting in tubulus cell necrosis.

The blood concentration of cadmium serves as a reliable indicator for a recent exposition; while the urinary concentration reflects past exposure, body burden and renal accumulation [7].

Excretion of Cadmium takes place via faeces and urine

Cadmium nitrate occurs as a colorless solid. It is very soluble in dilute acids and soluble in ethanol, acetone, water, diethyl ether, and ethyl acetate [23]. Cadmium nitrate is available in technical and reagent grades with a purity of 99% or higher [5]. Systemic toxicity offer a means

of understanding the mechanisms of free radical induced organ and or tissue damage. Among the tissues and organs in the mammalian body, liver and kidney seem to be the most sensitive predictor of chemical toxicity. This is due to their involvement in metabolism, detoxification, storage and excretion of xenobiotics and there metabolites, making them important target organs for xenobiotic induced injuries [13].

The present study was therefore designed to evaluate toxic effects of cadmium nitrate on histological damage of liver of adult wistar rats.

MATERIALS AND METHODS

CADMIUM NITRATE

Procedure

The cadmium nitrate in this research was purchase from pharmaceutical shop in Yaba Lagos State, Nigeria in January 6,2015 and it was taken to anatomy laboratory of department of anatomy, college of medicine, university Of Lagos, Idi-Araba Lagos State, Nigeria and were authenticated by a staff in the Department of Chemistry University of Lagos, Lagos state Nigeria.

Animals and diet

Twenty (n=20) adult wistar rats of both sexes were obtained from a breeding stock maintained in the animal house, the animal had free access to rat chow and tap water and they were randomly divided into four (4) groups (A-D) of five (n=5) rats each in a separate room at a constant temperature ($22.0 \pm 1.0^{\circ}\text{C}$) under a 12h light/dark cycle. Rats in group A which served as control were given 10ml/kg/day of distilled water for 21days. Rats in group B, C, and D (cadmium nitrate group) were given 150mg/kg/bw cadmium nitrate, 225mg/kg/bw of cadmium nitrate and 300mg/kg/bw of cadmium nitrate administered orally through an orogastric cannula into the stomach via the esophagus, once a day respectively, for Twenty one (21) consecutive days. The average daily oral intake of cadmium by non-smokers living in unpolluted areas was estimated to 10–25 μg [24] while in laboratory animals oral LD50 values for mice and rats range from 60 to over 5000 mg/kg of body weight[25]. All experimental investigations were done in compliance with humane animal as stated in the “Guide to the care and use of Laboratory

Animals Resources". National Research Council, DHHS, Pub.No NIH 86-23 [26] and in accordance with the guideline and approval of Nigeria Medical Ethical Association for Accreditation of Laboratory Animal Care.

Animal Sacrificed and Sample Extraction

Twelve hours after the administration of the last cadmium nitrate, the rats were at the time of sacrifice first weighed and then cervical dislocation was carried out following ethical humane animal euthanasia which was adopted with expertised cervical dislocation. The abdominal cavity of each rat was opened up through a midline abdominal incision to expose the liver. The liver was excised and weighed; the liver was weighed with an electronic analytical and precision balance. The liver of each animal was fixed in 10% formol-saline for histological examination. (BA 210S, d=0.0001- Sartoriusen GA, Goettingen, Germany).

Determination of Serum ALT, AST and ALP Assay

The degree of liver damage was evaluated by ALT, AST and ALP in serum using a commercially available kit. Detailed procedures for ALT, AST and ALP measurements were performed according to the kit manufacturer's instructions which was Sigma Aldrich products, made in Germany.

Histological Procedures and Analysis

This was done as described by Ogunlade *et al* [38]. Briefly, the organs were cut on slabs about 0.5cm thick and fixed in 10% formol saline for a day after which they were transferred to 70% alcohol for dehydration. The tissues were passed through 90% alcohol and chloroform for different durations before they were transferred into two changes of molten paraffin wax for 20min each in an oven at 570 C. Serial sections of 5µm thick were obtained from a solid block of tissue and were stained with haematoxylin and eosin stains, after which they were passed through a mixture of equal concentration of xylene and alcohol. Following clearance in xylene, the tissues were oven- dried. Photomicrographs were taken with a JVC colour video digital camera (JVC, China) mounted on an Olympus light microscope (Olympus UK

Ltd, Essex,UK) to demonstrate the liver damage

Statistical Analysis

All data were expressed as mean $\pm$ S.D. Differences between groups were analyzed using one-way analysis of variance (ANOVA). A value of $p < 0.05$ was considered to be statistically significant

RESULTS

TABLE 1. This table shows mean body weight and standard deviation of the experimental and control rats

GROUPS	DAY 7 (Mean $\pm$ S.D)	DAY 14 (Mean $\pm$ S.D)	DAY 21 (Mean $\pm$ S.D)
A(CONTROL)	118 $\pm$ 8.19	182 $\pm$ 9.11	202 $\pm$ 8.94
B	160 $\pm$ 62.60	150 $\pm$ 7.38*	156 $\pm$ 3.57
C	138 $\pm$ 5.39	162 $\pm$ 3.34	178 $\pm$ 5.90*
D	174 $\pm$ 67.00	136 $\pm$ 4.32	146 $\pm$ 5.21*

Value are expressed as n=5 mean $\pm$ SEM *p<0.005

TABLE 2: This table shows liver weight in experiment and control rats

Parameter	Group A(control)	Group B	Group C	Group D
Liver weight	1.21 $\pm$ 0.29	1.12 $\pm$ 0.01*	0.88 $\pm$ 0.32	1.03 $\pm$ 0.30*

Value are expressed as n=5 Mean $\pm$ SEM *p<0.005

Mean body weights, liver weights in experimental and control rats

As showed in table 2. Body weights of experimental rats decreased as compared to the control group. The mean liver weight also decreased as compared to the control.

TABLE 3: Results of acute toxicity test for cadmium Nitrate.

Test serial /No	Animal identity	Dose of cadmium nitrate mg/kg	Short term (72hours)	Long term (12days)
1	RET	400	+	+
2	LET	400	+	+
3	TC	400	+	+
4	RLT	400	+	+
5	I	400	+	+

+ -- Survival, REP—Right ear tagged, LEP—Left ear tagged, TP—Tail cut, RLT—Right leg tagged, I—Intact rat.

Acute oral Toxicity Studies

There was no death of rats at dose 400 mg/kg body weight of cadmium nitrate both within the short and long outcome of the limit dose test of Up and Down method (Table 3).The LD50 was calculated to be greater than 400 mg/kg body weight /orally

TABLE: 4 shows AST and ALT levels in control and experiment rats

GROUPS	AST(U/l)	ALT(U/l)
A(CONTROL)	26.0 ± 0.1	24.3 ± 0.2
B	110.9 ± 0.4**	130.4 ± 0.2**
C	110.2 ± 0.1**	129.2 ± 0.1
D	108.7 ± 0.5**	127.2 ± 0.1**

Values expressed as $n=5$ mean $\pm$ SD, ** $p < 0.001$ significant

AST and ALT levels

As showed in table 4, the serum levels of AST and ALT were significantly increased in cadmium nitrate group compared to the control group following the exposure to cadmium nitrate.

Liver Histology

Cross section of the livers of rats after the study showed Group A (control) had normal central vein (CV) with normal sinusoid (S) around the hepatocyte(H).[fig1] Group B show mild steatosis , moderate interstitial inflammation and mild periportal inflammation .[fig2] Group C show disseminated moderate macrovesicular steatosis and mild periportal inflammation . There is marked congestion of blood vessels and congestion of sinusoids with sinusoidal dilation .[fig3]

Group D show mild steatosis and moderate congestion of blood vessels. There are moderate periportal inflammation.[fig4]

PHOTOMICROGRAPY DEMOSTRATIONS

GROUP A (CONTROL)

Photomicrographs of the liver of experimental animal administered 10ml/kg/day of distilled water as control.




Fig.1. Histological demonstration of the liver using H & E staining techniques [X400] showing normal central vein (CV, black arrow) with normal sinusoid (S, yellow arrow around the hepatocyte(H, blue arrow).

GROUP B

Photomicrographs of the liver of experimental animal administered 300mg/kg/bw cadmium nitrate for twenty one days .





Fig.2. Histological demonstration of the liver using H & E staining techniques [X400] showing mild steatosis (black arrow), moderate interstitial inflammation (blue arrow) and mild periportal inflammation (yellow arrow).

GROUP C (225

Photomicrographs of the liver of experimental animal administered 225mg/kg/bw cadmium nitrate for twenty one days .



176


177 Fig.3. Histological demonstration of the liver using H & E staining techniques [X400] showing
 178 disseminated moderate macrovesicular steatosis (blue arrow) and mild periportal inflammation
 179 (black arrow). There is marked congestion (yellow arrow) of blood vessels and congestion of
 180 sinusoids with sinusoidal dilation (black arrow)

181

182 **GROUP D**

183 Photomicrographs of the liver of experimental animal administered 150mg/kg/bw cadmium
 184 nitrate for twenty one days .

185



186

187 Fig.4. Histological demonstration of the liver using H & E staining techniques [X400] showing
 188 mild steatosis and moderate congestion (yellow arrow) of blood vessels. There are moderate
 189 periportal inflammation (black arrow).

190 **DISCUSSION**

191 Water pollution has become a global problem. Heavy metals have long been recognized as
 192 serious pollutants of the aquatic environment. Health problems have been widely reported due
 193 long-term ingestion of contaminated drinking water with heavy metals. Pollution of water bodies
 194 with heavy metals from variety of sources is becoming a matter of global concern [2]. Though
 195 effects of chemical contamination of drinking water are not felt on short-term basis (except
 196 nitrate), their accumulation over a long period in the body has significant health effects [9].
 197 Present study demonstrates the toxic effect of administration of cadmium nitrate on the liver of
 198 experimental animals. Shows that exposure of liver to cadmium nitrate significantly increased
 199 serum intracellular enzymes such as aminotransferase (AST) and aspartate aminotransferase
 200 (ALT) compared to the control group. These are in accordance as reported by Tzirogiannis et al
 201 [20] that AST and ALT levels increased for 10 hours after injection CdL. It was supported that
 202 AST and ALT was increased five hours after injection. m.AST is an indicator of mitochondrial
 203 injury and is important in the clinical assessment of severe liver injury[21] . Osfor et al also
 204 reported that the levels of blood ALT, AST, ALP increased in Cd treated rats with several damages of
 205 liver tissues structure [22].

The histological evaluation can be used to shows the severity and toxicity of cadmium nitrate induced liver damage. It was observed in the control group that the normal hepatic cytoarchitecture was evident with visible terminal hepatic lobules consisting of terminal hepatic venules, hepatocyies with intervening sinusoidal spaces radially accentuated [Fig.1]. Cadmium nitrate Group B show mild steatosis , moderate interstitial inflammation and mild periportal inflammation [fig.2]. Group C show disseminated moderate macrovesicular steatosis and mild periportal inflammation. There is marked congestion of blood vessels and congestion of sinusoids with sinusoidal dilation [fig.3]. Group D show mild steatosis and moderate congestion of blood vessels, there are moderate periportal inflammations [fig.4]. This is in line with the report of Osfor et al that administration of Cd in rats cause several liver tissue structure damages [22]. we therefore deduced from our findings that oral administration of cadmium nitrate induced liver injury in adult wistar rats.

CONCLUSION

It can be therefore concluded from the present study that cadmium nitrate induced liver injury following the oral administration

ETHICAL APPROVAL

All the authors hereby declare that all the experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in line with the ethical procedure laid down by Nigeria Medical Ethical Association for Accreditation of Laboratory Animal Care.

REFERENCE

1. ATSDR Toxicological Profile for Cadmium. Agency for Toxic Substances and Disease RegistryU.S. Department of Health and Human Services, Atlanta, GA (2012)
2. Dike N.I, A.U. Ezealor, S.J. Oniye Concentration of Pb, Cu, Fe, and Cd during the dry season in river Jakau, Kano Nigeria Chem. Class. J., 1 (2004), pp. 78–81
3. Elinder CG, Lind B, Kjellstrom T, Linnman L, Friberg L. Cadmium in kidney cortex, liver, and pancreas from Swedish autopsies. Estimation of biological half time in kidney

cortex, considering calorie intake and smoking habits. Arch Environ Health. 1976;31:292–302.

4. Giles, M.A. Accumulation of cadmium by rainbow trout, *Salmo gairdneri*, during extended exposure. Can. J. Fish. Aquat. Sci. 1988, 45, 1045–1053.
5. IARC (1993). *Summaries & evaluations: Cadmium and cadmium compounds (Group 1)*. Lyon, International Agency for Research on Cancer, p. 119 (IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 58; <http://www.inchem.org/documents/iarc/vol58/mono58-2.html>)
6. IARC (in preparation). A review of human carcinogens. C: Metals, arsenic, dusts, and fibres. Lyon, International Agency for Research on Cancer (IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 100) [summary in Straif K et al. (2009). A review of human carcinogens—Part C: Metals, arsenic, dusts, and fibres. *The Lancet Oncology*, 10:453–454 ([http://www.thelancet.com/journals/lanonc/article/PIIS1470-2045\(09\)70134-2/fulltext](http://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(09)70134-2/fulltext)).
7. Jin T, Nordberg M, Frech W, Dumont X, Bernard A, Ye TT, Kong Q, Wang Z, Li P, Lundstrom NG, Li Y, Nordberg GF. Cadmium biomonitoring and renal dysfunction among a population environmentally exposed to cadmium from smelting in China (ChinaCad) Biometals. 2002;15:397–410. doi: 10.1023/A:1020229923095.
8. Leblanc J.C., Guérin T., Noël L., Calamassi-Tran G., Volatier J.L, Verger .P, Dietary exposure estimates of 18 elements from the 1st French Total Diet Study Food Addit. Contam., 22 (2005), pp. 624–641
9. Musa .H., I.A. Yakasai, H.H. Musa Determination of lead concentration in well and bore hole water in Zaria Nigeria. Chem. Class. J., 1 (2004), pp. 14–18
10. Ogunlade B, Saalu LC, Ogunmodede OS, Akunna GG, Adeeyo OA, Ajayi GO. The salutary role of *Allium cepa* extract on the liver histology, liver oxidative status and liver marker enzymes of rabbits submitted to alcohol- induced toxicity. American journal of Biochemistry and Molecular Biology. 2012;2(2):67-81.
11. Orłowski C, Piotrowski JK. Biological levels of cadmium and zinc in the small intestine of non-occupationally exposed human subjects. Hum Exp Toxicol. 2003;22:57–63. doi: 10.1191/0960327103ht326oa.

12. Pentyala .S., J. Ruggeri, A. Veerraju, Z. Yu, A. Bhatia, D. Desai, P. Vig Microsomal
Ca²⁺ flux modulation as an indicator of heavy metal toxicity Indian J. Exp. Biol., 48
(2010), pp. 737–743
13. Tayeb W, Nakbi A, Trabelsi M, Attia N, Miled A, Hammami M. Hepatotoxicity induced by
sub-acute exposure of rats to 2,4-Dichlorophenoxyacetic acid based herbicide “Desormone
Lourd”. *J Haz Mat.* 2010; 180: 225-233.
14. U.S Geological Survey. Mineral commodity Summaries, January 2005(online),available
from <http://mineral.vsgs.gov/minerals/pubs/commodity/cadmium>,(accessed 2006-01-01).
15. Wester RC, Maibach HI, Sedik L, Melendres J, DiZio S, Wade M. In vitro percutaneous
absorption of cadmium from water and soil into human skin. *Fundam Appl Toxicol.*
1992;19:1–5. doi: 10.1016/0272-0590(92)90021-9.
16. World Health Organisation (WHO) (1992). Environmental HealthCriteria 134-Cadmium
International Programme on Chemical Safety (IPCS) Monograph.
17. WHO, 1995. Environmental Health Criteria 2000, Copper. World Health Organization,
Geneva.
18. Yate EM. The world needs cadmium-a miner’s viewpoint . In cook ME,eds.
Cadmium92. London :cadmium Association,1992:1-7
19. Zalups RK AS. Molecular handling of cadmium in transporting epithelia. *Toxicol Appl
Pharmacol.* 2003;186:163–188. doi: 10.1016/S0041-008X(02)00021-2.
20. Tzirogiannis KN,Panoutsopoulos GI,Denonakou MD, Hereti RI, Alexandropoulos MG,
Bassyannis AC and Mykoniatis MG: Time-course of Cadmium-induced acute
hepatotoxicity in the rat liver. The role of apoptosis. *Arch Toxicol* 77,694-701 (2003)
21. Numakami k, Aoki Y, Ogawa Z and Itoh H: A kinetic model of mitochondrial aspartate
aminotransferase transmigration in hepatobiliary disorder. *Ann Clin Biochem* 36,226-232
(1999).
22. Osfor M.M.H, Ibrahim H.S., Mohamed Y.A., Ahmed A.M., Abd El Azeem A.S.,
Hegazy A.M. Effect of alpha lipoic acid and vitamin E on heavy metals intoxication in
male albino rats *J. Am. Sci.*, 6 (2010), pp. 56–63

- 290
291 23. James, D. W.; Carrick, M. T.; Leong, W. H. (1978). "Raman spectrum of cadmium
292 nitrate". Australian Journal of Chemistry **31** (6): 1189. doi:10.1071/CH9781189.
- 293 24. WHO (1992) *Cadmium*. Geneva, World Health Organization (Environmental Health
294 Criteria 134).
- 295 25. Krajnc EI et al. (1987) *Integrated criteria document. Cadmium — Effects. Appendix.*
296 Bilthoven, National Institute of Public Health and Environmental Protection (Report No.
297 758476004).
- 298 26. National Research Council (1985) "Guide to the care and use of Laboratory Animals
299 Resources". National Research Council, DHHS, Pub.No NIH 86-23