

## **Original Research Article**

### **Phytochemical Screening and Antimicrobial Activity of Leaf Extract of *Jatropha curcas***

#### **ABSTRACT**

As a measure of testing the medicinal properties of *Jatropha curcas*, this study screens aqueous, methanol and ethanolic extracts obtained from the leaf for phytochemicals and investigates the growth inhibition of *Escherichia coli*, *Staphylococcus aureus*, *Proteus spp*, *Klebsiella pneumonia* and *Pseudomonas aeruginosa* by methanolic, ethanolic and aqueous leaf extracts of *Jatropha curcas*. Tannins, saponins, flavonoids, alkaloids, oxalates and cyanogenic glycosides were the phytochemicals present. The test microorganisms were all significantly inhibited by the leaf extracts but the presence of oxalates and cyanogenic glycosides in the leaf renders it inedible and reduces its application to topical use only.

**Keywords:** *Jatropha curcas*, Phytochemical, antimicrobial

#### **INTRODUCTION**

Microbial infections are prima public health problems in the developed countries. Due to general use of commercial antibiotics, the relative incidence of multiple antibiotic resistances in human pathogens is not only large but also growing (Li and Krumholz, 2007). Intractable problems of resistance to antibiotics by microorganisms previously thought to have been brought under control have led to the resurface of interest in herbal products as source of noble compounds to suppress or possibly eradicate the ever increasing problems of emergence of newer diseases (Jain 1996, Mothanaa 2005). The World Health Organization estimated that about 80% of people still rely mainly on traditional remedies such as herbs for their medicines (WHO, 2003). Most of the plants used by the rural residential areas have biologically active compounds that have been proven by generations to be potent against specific disorders (Mark, 2002; Al-Sayed et al., 2014;

Eldahshan and Abdel-Daim, 2015). The prevention and treatment of diseases by the use of available and accessible medicinal plants in a particular locality will continue to play important roles in medical health care implementation in the developing countries as plants make up the primary source of new pharmaceuticals and health care products (Ivanona *et al.*, 2005). Natural products are therefore gaining attention as an alternative for antimicrobial agents (Helal *et al.*, 2015).

There have been major role played by *Jatropha curcas* in the treatment of different kinds of diseases ranging from bacterial to fungal infections (Thomas *et al.*, 2008). Glycosides, tannins, phytosterol, flavonoids and steroidal sapogenins are contained in different parts of the plants (leaves, fruits, latex and bark); a feature which enables it to demonstrate vast range of medicinal properties (Dalziel, 1955; Duke, 1988; Duke, 1994; Edeoga *et al.*, 2005; Agbogidi and Eruotor, 2012). The leaves have been reported to be a therapy for malaria, scabies, paralysis and hard tumor (Duke *et al.*, 2002; Aliyu, 2006). Also, because of the presence of ansovitexin, apigenin and vitexin in the leaves, they can serve as a cure for rheumatism and muscular pains (Rejore and Batra, 2003; El-Ekanali, 2010; Agbogidi and Ekeke, 2011). Antibiotic activities of *Jatropha curcas* seed have been demonstrated against *Staphylococcus aureus* and *Escherichia coli* (Thomas *et al.*, 2008). With the wealth of tradomedicinal claims surrounding the use of *Jatropha curcas* leaves (Esimone *et al.*, 2009), there is need for investigation of antibiotic potential of the leaves against some common pathogenic microorganisms. This research therefore investigates the growth inhibition of *E. coli*, *S. aureus*, *Proteus spp*, *Klebsiella pneumonia* and *Pseudomonas aeruginosa* by methanolic, ethanolic and aqueous leaf extracts of *Jatropha curcas*.

## MATERIALS AND METHODS

**Media Used:** Nutrient media (Micro master) and MacConkey agar (Lab M).

**Test Organisms:** While carrying out this work, the following bacteria were used as the test organisms: *Escherichia coli*, *Staphylococcus aureus*, *Proteus spp*, *Klebsiella pneumonia* and *Pseudomonas aeruginosa*. These organisms were obtained from the Microbiology Laboratory of Abia State University, Nigeria.

**Leave Extracts:** Fresh leaves of *Jatropha curcas* were collected from farm land in Agbani, Enugu State, Nigeria. The plant was identified and authenticated by the Department of Botany, University of Uyo, Nigeria. Voucher sample was prepared and deposited in the Herbarium for

reference. The leaves of *Jatropha curcas* were allowed to air-dry at room temperature for fourteen (14) days until the leaves became brittle, and ground into fine powder in a clean mortar using a clean pestle. Twenty grams of the leaves powder were weighed and poured into three (3) conical flasks; 100ml of distilled water, 70% ethanol, and 70% methanol. Ethanol and methanol used were of analytical grade. The preparations were allowed to stand undisturbed, but with occasional agitation, for 24 hours and filtered using Whatman No. 1 filter paper. The residues were discarded and the filtrates (extracts) were divided into two portions. One portion was refrigerated at 4°C until needed. The other was evaporated to dryness in a water bath at 100°C. To 1g each evaporated extract; 10ml of the initial solvents that is methanol, ethanol or water were added.

### **Phytochemical Analysis of the Plant Extracts**

The extracts were subjected to qualitative phytochemical tests for tannins, saponins, flavonoids, alkaloids, oxalates and glycosides in accordance with Trease and Evans (1989) and Harborne (1998) with little modification.

### **Antimicrobial Activity**

#### ***Preparation of Culture Media***

*Nutrient agar preparation:* The medium was prepared by suspending 2.8g of the agar powder into a 100ml of distilled water, in a conical flask and shaken very well to dissolve. It was then autoclaved at 121°C for 15mins and allowed to cool at 47°C before pouring into plates aseptically in the required amount. The medium was allowed to solidify (Cheesbrough, 2003).

*Macconkey agar preparation:* This medium was prepared by weighing 4.8g of agar powder into a conical flask; 100ml of distilled water was poured into the flask. The flask was corked, shaken and allowed to soak for about 10minutes. The medium was autoclaved at 121°C for 15minutes. After autoclaving, it was allowed to cool to 47°C before pouring into the plates aseptically in the required amount and allowed to solidify (Cheesbrough, 2003).

#### ***Incubation and Inoculation***

The agar well diffusion method was used in the assessment of the antimicrobial activity of the plant, this method was achieved by using a sterile cork borer to make six wells on each Nutrient and MacConkey agar plates, then plant extracts were dropped into wells and allowed to diffuse for 5 minutes. Tetracycline, a standard antibiotic, was used as control. 0.5ml of tetracycline was dropped in each hole. Plates were then kept to stand in the incubator for 24hours at 37°C (Igbinosa *et al*, 2009). Zones of inhibition were measured and recorded. The experiment was done in triplicate and the results are expressed as mean  $\pm$  standard deviation.

## RESULT AND DISCUSSION

**Table 1:** Phytochemical screening of methanolic, ethanolic and aqueous extract of *Jatropha curcas* Leaves

| Phytochemicals        | Methanolic extract | Ethanolic extract | Aqueous extract |
|-----------------------|--------------------|-------------------|-----------------|
| Saponin               | -                  | -                 | +++             |
| Tannins               | -                  | ++                | +               |
| Flavonoids            | +++                | +                 | -               |
| Cyanogenic glycosides | +++                | ++                | ++              |
| Alkaloids             | -                  | +                 | +++             |
| Oxalate               | ++                 | +++               | -               |

+++Highly present, ++Moderately present, +Present, -Absent

Phytochemical screening of *Jatropha curcas* leaf, presented in Table 1, revealed the presence of tannins, saponins, flavonoids, alkaloids, oxalates and cyanogenic glycosides. These phytochemicals are biologically active and can be said to responsible for the antimicrobial activity of the plant. The mechanisms through which these secondary metabolites exert their antimicrobial activities differ. Tannins have been reported to form irreversible complexes with proline rich protein (Shimada, 2006), thereby inhibiting cell protein synthesis. (Parekh and Chanda, 2007). Intestinal disorders such as diarrhea and dysentery have been treated with herbs that have tannins as their main components. (Dharmananda, 2003).

Another secondary metabolite observed in the leave extract of *J. curcas* was alkaloid. Alkaloids have analgesic effects (Sutradhar *et al.*, 2007) and have been used in pain killer medications

(Kam and Liew, 2002). Alkaloids have been acclaimed for their antimicrobial activities, especially against gram negative bacteria (Cushnie *et al.*, 2014). Other secondary metabolites present in *J. curcas* whose antimicrobial activities have been documented are flavonoids and saponins (Hodek *et al.*, 2002).

Though traditionally used in the fermentation of cassava, *J. curcas* is inedible. This is due to the presence of oxalate and cyanogenic glycosides in it as presented in Table 1. Oxalate is an antinutrient and an acute toxin (Munro and Bassir, 1996). Cyanogenic glycosides are not just distasteful but also toxic, as they release hydrogen cyanide which generates free radicals and causes oxidative stress (Gleadow and Woodrow, 2002).

**Table 2:** Antibacterial Activity of *Jatropha Curcas* Leaves Extracts Using Each Solvent (Water, Ethanol And Methanol) at Different Concentrations.

| Organisms                     | Zone of Inhibition (mm) |                |                |                |                |                |         | AI   |
|-------------------------------|-------------------------|----------------|----------------|----------------|----------------|----------------|---------|------|
|                               | W <sub>1</sub>          | W <sub>2</sub> | E <sub>1</sub> | E <sub>2</sub> | M <sub>1</sub> | M <sub>2</sub> | Control |      |
| <i>Escherichia coli</i>       | 7.0±1.4                 | 7.0±1.4        | 9.0±1.4        | 10.0±2.8       | 11.0±4.2       | 9.0±1.4        | 8       | 7.25 |
| <i>Staphylococcus aureus</i>  | 8.5±0.71                | 8.0±0.17       | 8.0±1.4        | 8.0±1.4        | 7.5±2.12       | 8.5±0.7        | 9       | 4.9  |
| <i>Proteus spp</i>            | 6.5±0.71                | 6.5 ± 0.71     | 7.0±0          | 6.5±0.71       | 8.0±1.4        | 7.0±0          | 7       | 5.9  |
| <i>Klebsiella pneumonia</i>   | 6.5±2.12                | 6.5±2.12       | 8.0±1.4        | 8.0±1.4        | 11.5±3.5       | 10.5±2.12      | 9       | 5.8  |
| <i>Pseudomonas aeruginosa</i> | 7.5±0.71                | 7.0±0          | 7.5±0.71       | 7.0±0          | 8.0±1.4        | 8.0±1.4        | 7       | 6.8  |

Means±Standard Deviation

W<sub>1</sub> = Water extract (refrigerated) 4°C

W<sub>2</sub> = Water extract (evaporated) 100°C

E<sub>1</sub> = Ethanolic extract (refrigerated) 4°C

E<sub>2</sub> = Ethanolic extract (evaporated) 100°C

M<sub>1</sub> = Methanolic extract (refrigerated) 4°C

M<sub>2</sub> = Methanolic extract (evaporated) 100°C

AI (Activity Index) =  $\frac{\text{Mean of zone of Inhibition by Extract}}{\text{Zone of Inhibition by Antibiotic Drug}}$

As presented in Table 2, all the extracts demonstrated significant inhibition against the test microorganisms. Ethanolic extract however demonstrated the greatest inhibition. This is

consistent with the high level of phytochemicals in the ethanolic extract. The inhibition was most effective against *E coli* with mean inhibition zone of 8.33mm. This is consistent with the inhibition by stem bark extracts of *Jatropha curcas* (Igbinosa *et al.*, 2009). The use of *J. curcas* as a medication for diarrhea is however not feasible because of the presence of oxalate and cyanogenic glycosides in it. The extracts are also active against *Staphylococcus aureus* and *Klebsiella pneumonia*. This agrees with the report of Ekundayo and co workers (2011) on the antimicrobial activities of the nut extracts of *J. curcas*. *Staphylococcus aureus* is the causative organism for skin infections such as abscesses, respiratory infections such as sinusitis, and food poisoning (Curran *et al.*, 1980). *Klebsiella pneumonia* infections are seen mostly in people with a weakened immune system. *Klebsiella* infections are often nosocomial. The most common condition caused by *Klebsiella* outside the hospital is pneumonia (Brooks, *et al.*, 2004). *Klebsiella* is also implicated in urinary tract, lower biliary tract, and surgical wound sites infections (Ryan and Ray, 2004). The inhibition of *S. aureus* and *K. pneumonia* by the leave extract substantiates the trado-medicinal use of *J. curcas* in wound healing, scabies and other skin infections (Thomas *et al.*, 2008).

Other microorganisms inhibited include *Proteus spp* and *Pseudomonas aeruginosa*. Similar inhibitions by the nuts and stem bark of *J. curcas* have been reported (Igbinosa *et al.*, 2009; Ekundayo *et al.*, 2011). *Pseudomonas aeruginosa* is a ubiquitous multi drug resistant pathogen that is associated with serious illnesses - especially nosocomial infections such as ventilator-associated pneumonia and various sepsis syndromes as well as hot-tub rash (Balcht and Smith, 1994). *Proteus spp* is implicated in urinary, septic and wound infections, often nosocomial (Guentzel, 1996). However, it is opportunistic and only affects immunosuppressed individuals (Brooks, *et al.*, 2004). The inhibition of *Proteus spp* by *Jatropha curcas*, as shown in Table 2, substantiates the reported wound healing ability of *Jatropha curcas* (Shetty *et al.*, 2006; Esimone *et al.*, 2008). There were no significant differences between the inhibitions by the refrigerated and evaporated extracts. This demonstrates that the preservation methods have no significant effect on the antimicrobial properties of the extracts.

## CONCLUSION

Phytochemical screening of *Jatropha curcas* leaf revealed the presence of several biological active compounds including tannins, saponins, flavonoids, alkaloids, oxalates and cyanogenic

glycosides; which can be responsible for the observed antimicrobial action of the leave. The leaf demonstrated considerable inhibition zone against the test microorganisms but can only be used for topical treatment because of its inedibility.

## REFERENCES

Agbogidi OM, Ekeke EA. (2011). *Jatropha curcas*: Linn an important but neglected plant species in Nigeria. *J Biol Chem Res.* **281** : 52-62.

Agbogidi OM, Eruotor PG. (2012). Morphological Changes due to spent engine oil contamination and its heavy metal components of *Jatropha curcas*. In: Baby, S. and Sandhu, P.S. eds. . Proceedings of the International Conference on Bioscience, Biotechnology and Health Sciences ICBBHs' 2012 organized by Planetary Science Centre Research December 14 and 15, 2012 in Singapore. Pp. 88-93.

Aliyu BS. (2006) . Some ethnomdicinal plants of the Savannah regions of West Africa: description and phytochemicals. Triumph Publishing Company Ltd, Kano WHO (World Health Organization) (2003). Traditional Medicine, Fact sheet No 134.

Al-Sayed E, Martiskainen O, Seif el-Din SH, Sabra AA, Hammam AO, El-Lakkany NM, and Abdel-Daim MM. (2014). Hepatoprotective and Antioxidant Effect of *Bauhinia hookeri* Extract against Carbon Tetrachloride-Induced Hepatotoxicity in Mice and Characterization of Its Bioactive Compounds by HPLC-PDA-ESI-MS/MS. *BioMed Research International*. Article ID 245171. doi:10.1155/2014/245171.

Balcht A. and Smith R. (1994). *Pseudomonas aeruginosa*: Infections and Treatment. Informa Health Care. pp. 83–84. ISBN 0-8247-9210-6.

Brooks, G. F., Butel, J.S., Morse, S. A., Carroll, K. C., and Mietzner, T. A. (2004). Medical

Cheesbrough M. (2003). Medical Laboratory Manual. Tropical Health Technology, Low Priced Edition. Doddington, Cambridge shire, England. Pp. 132-143.

Curran JP, Al-Salihi FL; Al-Salihi (1980). "Neonatal staphylococcal scalded skin syndrome: massive outbreak due to an unusual phage type". *Pediatrics* **66** (2): 285–90. PMID 6447271.

189 Cushnie TPT, Cushnie B, Lamb AJ (2014). Alkaloids: An overview of their antibacterial,  
 190 antibiotic-enhancing and antivirulence activities. *International Journal of Antimicrobial*  
 191 *Agents*: **44**(5):

192 Dalziel JM (1955). The Useful Plants of West-Tropical Africa. Crown Agents for Oversea  
 193 Governments and Administration, London, P.147

194 Dharmananda S (2003). Gallnuts and the uses of Tannins in Chinese Medicine. In: Proceedings  
 195 of Institute for Traditional Medicine, Portland, Oregon.

196 Duke JA (1988). CRC Handbook of Medicinal Herbs. CRC Press, Boca Raton, Florida, pp. 253–  
 197 254.

198 Duke JA, Bogenschutz-Godwin MJ, DuCellier J, Duke PAK. 2002 . Handbook of medicinal  
 199 herbs. 2nd Edition, Boca Raton, CRC Press.

200 Duke JA. 1994 . Biologically active compounds in important species. In: Charalambous, E. ed. .  
 201 Spices, herbs and edible fungi. Elsevier Science Oxford. Pp 225-250.

202 Edeoga HO, Okwu DE, Mbaebre BO. 2005 . Phytochemical constituents of some Nigerian  
 203 plants. *African Journal of Biotechnology* 44 7 :685-688.

204 Ekundayo FO, Adeboye CA and Ekundayo EA (2011). Antimicrobial activities and  
 205 phytochemical screening of pignut (*Jatropha curcas* Linn.) on some pathogenic bacteria.  
 206 *J. Med. Plant. Res.* **5**(7): 1261-1264.

207 Eldahshan OA and Abdel-Daim MM. (2015). Phytochemical study, cytotoxic, analgesic,  
 208 antipyretic and anti-inflammatory activities of *Strychnos nux-vomica*. *Cytotechnology*.  
 209 **67**(5):831-44. doi: 10.1007/s10616-014-9723-2.

210 Esimone CO, Nworu CS and Jackson CL. (2008). Cutaneous wound healing activity of a  
 211 herbal ointment containing the leaf extract of *Jatropha curcas* L. (Euphorbiaceae).  
 212 *IJARNP*. **1**(4): 1-4.

213 Esimone CO, Nworu CS and Jackson CL. (2009). Cutaneous wound healing activity of a herbal  
 214 ointment containing the leaf extract of *Jatropha curcas* L. (Euphorbiaceae). *International*  
 215 *Journal of Applied Research in Natural Products*. **1**(4), pp. 1-4.

216 Gleadow RM and Woodrow IE (2002). Mini-Review: Constraints on Effectiveness of  
 217 Cyanogenic Glycosides in Herbivore Defense. *Journal of Chemical Ecology*. **28**(7):  
 218 1301-1313.

219 Gübitz GM, Mittelbach M, Trabi M (1999) Exploitation of the tropical oil seed plant *Jatropha*  
 220 *curcas* L. *Bio Tech.*, 67: 73–82.

221 Guentzel MN (1996). *Escherichia*, *Klebsiella*, *Enterobacter*, *Serratia*, *Citrobacter*, and *Proteus*.  
 222 In: Barron's Medical Microbiology (Barron S *et al.*, eds.) (4th ed.). Univ of Texas  
 223 Medical Branch. ISBN 0-9631172-1-1. (via NCBI Bookshelf)

224 Harborne JB (1998). *Phytochemical Methods - A Guide to Modern Techniques of Plant*  
 225 *Analysis*. Chapman and Hall, London. pp. 182- 190

226 Helal MS, Youssef FM, Moursi MK, Khalil WF, Abdel-Daim MM. (2015). Effectiveness of  
 227 prebiotic as an alternative to the antimicrobial growth promoter on growth performance,  
 228 blood constituents, intestinal healthiness and immunity of broilers. *AJVS*. **45**(1): 13-25.  
 229 doi:10.5455/ajvs.179869

230 Igbinosa, O. O., Igbinosa, E.O. and Aiyegoro, O. A. (2009). Antimicrobial activity and  
 231 Phytochemical screening of stem bark extracts from *Jatropha curcas*. *African Journal of*  
 232 *Pharmacology*. **3** (2): 58-62.

233 Igbinosa, O. O., Igbinosa, E.O. and Aiyegoro, O. A. (2009). Antimicrobial activity and  
 234 Phytochemical screening of stem bark extracts from *Jatrophacurcas*. *African Journal*  
 235 *of Pharmacology*. **3** (2): 58-62.

236 Ivanova, D., D. Gerova T. Chervenkov and T. Yankova. Polyphenols and antioxidant capacity of  
 237 Bulgarian medicinal plants. *J.Ethnopharmacol.*, 2005;96:145-150.

238 Jain, S. K. (1996). *Medicinal plants*, National Book Trust. India, pp. 3.

239 Kam PCA, Liew S (2002). Traditional Chinese herbal medicine and anaesthesia. *Anaesthesia*  
 240 **57**(11): 1083-1089.

241 Kumar A, Sharma S (2008). An evaluation of multipurpose oil seed crop for industrial uses  
 242 (*Jatropha curcas* L.): A review. *Bios Eng.*, 97: 201-207.

243 Li X. and Krumholz L. R. (2007). Regulation of arsenate resistance in *Desulfovibrio*  
 244 *desulfuricans* G20 by an *arsRBCC* operon and an *arsC* Gene. *J. Bacteriol.* :**189**(10):  
 245 3705-11.

246 Mark M. S. (2002). *Phytochemicals in Nutrition and Health*. CRC Press. p. 123. ISBN 9781587160837.

247 Martínez-Herrera J, Siddhuraju P, Francis G, D'ávila-Ortiz G, Becker K (2006). Chemical  
 248 composition, toxic/antimetabolic constituents, and effects of different treatments on their  
 249 levels, in four provenances of *Jatropha curcas* L. from Mexico. *Food Chem.*, 96: 80–89.

250 Microbiology. (25<sup>th</sup> edn.). New York: Mc Graw-Hill. Pp. 185-213.

251 Mothanaa, R. A. and Lindequist U. (2005). Antimicrobial activity of some medicinal plants.  
 252 *Journal of Ethnopharmacology*. **96**(1): 177-181.

253 Munro, A. and Bassir, O. (1996). Oxalates in Nigerian Vegetables. *West African Journal of*  
 254 *Biological and Applied Chem.* **12**:14-18.

255 Hodek P, Trefil P, Stiborova M (2002). Flavonoids - Potent and versatile biologically  
 256 active compounds interacting with cytochrome P450. *Chemico-Biol. Intern.*, **139**(1): 1-  
 257 21.

258 Parekh J, Chanda S (2007). *In vitro* antibacterial activity of crude methanol extract of *Wood for*  
 259 *dia fruticosa* Kurz flower (Lythaceae). *Braz. J. Microbiol.* 38: 2.

260 Rajore S, Batra A. 2003 . *Jatropha curcas*: a plant of immense potential value. *Journal of*  
 261 *Ecology, Taxonomy and Botany* 27:36-41.

262 Ryan KJ and Ray CG (2004). *Sherris Medical Microbiology* (4th ed.). McGraw Hill. ISBN  
 263 0-8385-8529-9.

264 Schmook B, Seralta-Peraza L (1997). *J. curcas*: distribution and uses in the Yucatan Peninsula  
 265 of Mexico. In: *Biofuels and industrial products from Jatropha curcas*. (Gübitz G M;  
 266 Mittelbach M; Trabi M, eds), DBV Graz, pp. 53–57

267 Shetty S, Udupa SL, Udupa AL and Vollala VR (2006). Wound healing activities of Bark  
 268 Extract of *Jatropha curcas* Linn in albino rats. *Saudi Med J.* **27**(10):1473-6.

269 Shimada T (2006). Salivary proteins as a defense against dietary tannins. *J. Chem. Ecol.* **32** (6):  
270 1149-1163.

271 Srivastav, J., Lambert, J., and Vietmeyer, N. (1996). Medicinal plants: An expanding role in  
272 development. *World Bank Technical Paper* No. 320.

273 Sutradhar RK, Rahman AM, Ahmad M, Bachar SC, Saha A, Roy TG (2007). Anti-inflammatory  
274 and analgesic alkaloid from *Sida cordifolia* linn. *Pak J Pharm Sci.* **20**(3):185-8.

275 Thomas T., Sah N. K. and Sharma P. B. (2008). Therapeutic Biology of *Jatropha curcas*: A Mini  
276 Review. *Curr Pharm Biotechnol.* **9**(4):315-24.

277 Trease GE, Evans WC (1989) Textbook of Pharmacognosy. 12th Edn. Balliere, Tinadl London