

# The Nutraceuticals: A Voluminous Torrent in Pharmaceuticals- Coupling Health & Drugs

## ABSTRACT

The present review is focused on the nutraceuticals which are present in our surroundings having an excellent impact over the health of humans but are not known for their pharmaceutical use. Nutraceuticals include vitamins, minerals and other dietary supplements, which may be herbs, enzymes, animal extracts etc. Various researchers have proved that nutraceuticals are having an important role in reducing the risk of various diseases such as diabetes, cardiovascular diseases, Parkinson's disease etc. with very few or negligible side effects.

*Keywords: Nutrients, Drugs, Dietary supplements, Functional foods.*

## 1. INTRODUCTION

"Let food be your medicine and medicine be your food" the great line said by great scientist Hippocrates is all related with the chemicals having nutritional and therapeutic value i.e. Nutraceuticals [1].

Nutraceuticals are a heterogenic products category which has a number of synonyms that are used internationally. The term "Nutraceutical" was given by Stephen De Felice who was the founder and chairman of the Foundation for Innovation in Medicine. This term has been part of the industry lexicon for almost a decade [2]. The term "Nutraceutical" is derived from the combination of two words, "nutrient" which means a nourishing food or food component and second "pharmaceutical" which means a medical drug. Nutraceuticals may contain substances that are "natural", intended to treat or prevent a number of diseases, but may not be generally recognized as safe. Hence these are the food products intended for health and medical benefit [3, 4]. It has been proved by research that the nutraceuticals are useful in providing protection from a number of diseases like diabetes, cancer, cardiac disease, hypertension etc., for example, carotenoids and anti-oxidants found in carrots help in avoiding chronic diseases, by preventing free radical damage [5]. Nutraceuticals have become more popular in modern society because of negligible chances of adverse effects of pharmaceuticals, ease of self-medication and aging in population.

## 2. TYPES OF NUTRACEUTICALS

### 2.1 Dietary Supplements

Dietary supplements (DS) include herbs, minerals, vitamins, or products obtained from plant sources, animal sources such as yeasts, fungus, algae, seafood and many more for example, energy bars, amino acids, and liquid supplements. They are not consumed

in large quantities but have the basic objective to provide nutrition. The United States authorities say that dietary supplements may be regarded as foods, while elsewhere they are classified as drugs or other products [6, 7].

## 2.2 Functional Foods

Japan introduced the concept of functional food in 1980s, to promote health or reduce the risk of diseases. The functional food include those food items which are advised to be consumed as part of the normal diet, they contain biologically active constituents offering the potential to enhance health or reduce risk of various diseases. Among these foods are those that contain fatty acids, vitamins, specific minerals or dietary fibers, foods with added biologically active substances such as phytochemicals or other antioxidants and probiotics that have beneficial live cultures [8]. Some examples of functional food products are; milk, cheese and eggs (enriched with omega-3 fatty acids); yogurt enhanced with live active cultures (probiotics); fruit juices and drinks (having antioxidant properties or containing antioxidants); cereals and grains such as wheat, oat, barley (having enriched amounts of dietary fibre); modified fatty acid vegetable oils; and soy, canola and hemp (vegetable proteins) and legumes [9, 10].

According with the established requirement for the functional food in Japan, the functional food should be consumed:


- a). In its natural form, rather than a prepared dosage form like capsule, tablet, or powder;
- b). Daily, in sufficient quantity; and
- c). In the right way to it can regulate a biological process, in order to prevent or cure a disease [11].

## 2.3 Dietary supplements and Food additives

Dietary supplements provide nutrients that may otherwise not be consumed in sufficient quantities. Supplements are generally like vitamins, minerals, fatty acids, fiber or amino acids etc. According to United State authorities dietary supplements are foods [12].

Food additives like the dietary suppliments, are any substances that are either deliberately added to food to enhance its shelf-life, nutrition, texture, or other quality aspect, or which unintentionally contaminates food (indirect additive) (Figure 1) [13].

Nutrients are the nutritive constituents present in food that a person consumes for his/her survival and growth. Macronutrients provide the bulk energy required for functioning of metabolic system, whereas micronutrients are helpful in providing the necessary co-factors for metabolism. Plenty of these nutrients are available in the environment [14]. Inorganic chemical compounds like water, minerals, and oxygen should also be considered as nutrients [15].



**Figure 1: Diagrammatic Representation of Nutraceuticals**

## 2.4 Herbals

In ancient time; a large number of herbs were used to prevent and treat many diseases. A plant containing non-nutritive phyto-chemicals, provides health benefits if included in diet [16, 17,18]. Nutraceuticals as herbals are big boon to human being in aspect of improving their health and to prevent them from chronic diseases e.g. willow bark (Salix Nigra), helps as anti-inflammatory, analgesic, anti-arthritis, astringent as well as antipyretic [19].

### *Phyto-chemicals*

Phyto-chemicals are the plant components having bio-activities which are used to get health benefits. But their use always requires to be defended with some scientific rational in food, as potential nutraceutical. Phyto-chemicals having following health benefits:

- (1) These are used as substrates for biochemical reactions.
- (2) These are used as cofactors of enzymatic reactions.
- (3) These act as ligands which agonize or antagonize cell surface or intracellular receptors.
- (4) These are used as scavengers of various reactive or toxic chemicals.
- (5) These are used as compounds to increase the absorption and or improve stability of many essential nutrients.
- (6) These work as selective growth factors for gut friendly bacteria.

(7) Fermented phyto-chemicals are beneficial for non-pathogenic bacteria found in GI tract.

(8) These are selective inhibitors of deleterious intestinal bacteria. Phyto-chemicals like terpenoids, phenolics, alkaloids and fiber, are extensively examined and used for their ability to get health advantage [20].

## 2.5 Probiotic/ Prebiotics

Probiotic bacteria are “living microorganisms that taken in tolerable quantity, provide the following benefits to host health”. These may interact with commensal bacteria to have a direct impact on the host. [21] Metchnikoff was the first person who successfully revamped the toxic flora of the large intestine into a host-sympathetic colony of *Bacillus bulgaricus*, found by Hord [24, 23]. Hence probiotics are gut friendly bacteria which aids in digestion and absorption of some nutrients. They act by eliminating the disease causing pathogens, like yeasts, other bacteria and viruses which mutually develop advantageous symbiosis within gastrointestinal tract (Table 1) [24]. The Japanese were the first to recognize the value of non-digestible oligosaccharides, and added these in feed of piglets to relieve and prevent from the diarrhoea. It was observed that fructo-oligosaccharides and galacto-oligosaccharides cause an increase in intestinal bifido-bacteria which stimulated their growth in the human gut. Hence a prebiotic is “a selectively fermented ingredient that causes changes in the activity of the gastrointestinal microbiota that are beneficial for human health” [25]. Some people use *Lactobacillus* for irritable bowel syndrome; Crohn's disease; inflammation of the colon; necrotizing enterocolitis in babies born prematurely, infection with *Helicobacter pylori*, urinary tract infections, vaginal yeast infections, in prevention of common cold in adults, and to prevent respiratory infections in children attending daycare centers. It is also being tested to prevent serious infections in people on ventilators [26-31].

A number of marketed nutraceuticals are available in market as mentioned in Table 2.

## 3. PHARMACOLOGICAL USE OF NUTRACEUTICALS

Better life quality is achieved with food items filled with nutrient values and remarkably potent in preventing diseases which may be cancer, diabetes, heart diseases, hypertension etc (Table 3 & Figure 2). Such products may be dietary supplements, food additives, phyto-chemicals, prebiotics, probiotics, genetically modified food, processed food and plant products.

137

138

**Table 1: Different species of microbes used as Probiotic**

<b>S.No.</b>	<b>Genus</b>	<b>Species</b>
1	Lactobacillus	Acidophilus Delbrueckia Brevis Fermentum Gasseri Johnsonii Paracasei Plantarum Reuteri Rhamnosus Salivarius
2	Bifidobacterium	Adolescentis Animalisb Breve Bifidum Infantis Longum
3	Streptococcus	Thermophilus Salivarius
4	Saccharomyces	Cerevisiae
5	Escherichia	Coli
6	Enterococcus	Faecium
7	Bacillus	Coagulansic Clausii

139

140

141


142

143

144

145

146



**Figure 2: Pharmacological Uses of Nutraceuticals**

### 3.1 In cardiovascular disease

It is not easy to set up a clear impact of nutrition/physical exercise on major cardiovascular diseases because history of cardiovascular diseases is too long [32]. The effect of calcium on hypertension and pre-eclampsia (a condition in pregnancy characterized by high blood pressure, sometimes with fluid retention and proteinuria) is unpredictable as well as ambiguous. It is supposed that high levels of intracellular calcium may increase vascular smooth muscle tone, peripheral vascular resistance, and responsiveness blood pressure. Treatment with vitamin C and selenium need further study to observe its effect on mortality [33-34].

It has been observed that some nutraceuticals may be useful to prevent the risk of thrombosis in women with thrombophilic gene mutations like vitamin E causes inhibition of platelet aggregation by a protein kinase C-dependent pathway [35-36]. Nutraceuticals like vitamins, minerals, omega-3 poly-unsaturated fatty acids (n-3 PUFAs), dietary fibers and antioxidants, in addition to physical exercise, are recommended to prevent and treat cardiovascular diseases. Researchers have proved that polyphenols found in grapes and in wine are helpful in reducing arterial disease by altering cellular metabolism and signalling [37].

Onion, black grapes, cherries, cruciferous vegetables, grapefruits, red wine, apples and berries are good sources of flavonoids [38] and also available as flavones and flavonols which are beneficial for the treatment of cardiovascular diseases [39-41]. Flavonoids block the angiotensin-converting enzyme, a key moiety that causes raise in blood pressure [42]. Antioxidant activity of ascorbic acid, alpha-tocopherol, and beta-carotene as has been studied and reviewed [43].

### 3.2 In management of diabetes

Various *in-vitro* and *in-vivo* studies (animal) have proved that plant polyphenols including phenolic acids, stilbenes, lignans and flavonoids are effective nutraceuticals in diabetes and in prevention. Although human clinical trials are required to check the efficacy of poly-phenol compounds in treatment of diabetes [44].

Vitamin C (ascorbic acid) is a chain-breaking antioxidant which prevents the propagation of chain reactions that may lead to a reduction in protein glycation. It has been reported that ascorbic acid helps in reducing diabetes-induced sorbitol in animals [16]. Ascorbic acid (800 mg/day) partially replenishes ascorbic acid levels in patients with type 2 diabetes but endothelial dysfunction or insulin resistance is not effected at all [17].

N. Bunyapraphatsara *et al* (1996) observed a combined effect of Aloe vera juice with glibenclamide in diabetic patients and concluded that glibenclamide alone did not show any effect while Aloevera juice showed significant reduction in fasting blood glucose level and triglycerides within two weeks and four weeks respectively. Although it did not show any effect on level of cholesterol but aloe vera was found to be effective in the treatment of diabetes [45]. Acacia is non-starch polysaccharides which is not digested in the intestine, but generate short chain fatty acids in large bowel; hence, it produces extensive biological effects. Philips AO *et al* (2011) conducted study over the extract of Acacia Arabica and confirmed the anti-diabetic effect of acacia resulted by increasing the insulin release [46]. Hou *et al* (2003) demonstrated significant hypoglycemic effects of Acacia Arabica powder in healthy rabbits, whereas no significant decrease in blood sugar was observed in the alloxan-induced diabetic rabbits was observed [47]. Wadood *et al* concluded that Acacia arabica initiates the release of insulin from pancreatic beta cells of normal rabbits [48]. Antidiabetic activity of glycerrhiza in non insulin dependent diabetic model was observed by Takii [49].

High intake of isoflavone (20–100 mg/day) is helpful in lowering of rate of mortality in diabetes of type 2, osteoporosis, cardiac disease and certain cancers [50]. Docosahexaenoic acid is vital for neurovisual development which helps in modulating insulin resistance and nurture the advocacy for essential fatty acids in pregnancy in women with gestational diabetes mellitus [51].

Omega-3 fatty acids helps in reducing blood glucose tolerance in patients predisposed to diabetes. Insulin is needed to synthesize long chain n-3 fatty acids; thus heart may be victim to their depletion in case of diabetes. Hence it is concluded that ethyl esters of n-3 fatty acids may be advantageous in diabetic patients [52]. Lipoic acid is a well known antioxidant, and in Germany being used to cure diabetic neuropathy. Lipoic acid; as a long-term dietary supplement is aimed at the prophylactic protection of diabetics from complications [53].  $\alpha$ -Lipoic acid enhances insulin sensitivity by approximately 18–20% in patients suffering from type 2 diabetes [54]. Clinical trial studies on  $\alpha$ -lipoic acid reported advantageous in the treatment of diabetic neuropathy [55].



Dietary fibers obtained from psyllium have been used to reduce weight and lipid levels in hyperlipidemia as dietary supplement [56]. Intake of Chromium supplements may be helpful to enhance sensitivity to insulin and boost glucose tolerance in type II diabetic patient [48]. Magnesium-rich diet intake may reduce risk of diabetes by improving in insulin sensitivity [57]. Diabetes management is supported by one of the nutraceuticals Biotin which increase insulin production and stimulates liver glucokinase activity, thus improves the uptake of glucose in muscle cells [58-60]. Pharmacological effect of epinephrine can be inhibited by Azadirachta Indica which results in enhanced utilization of peripheral glucose [61-62] and reduce hypo-glycaemic activity without change in the serum cortisol level [63-64].

Kernels of *Eugenia jambolana* (*Syzygium cumini*) are useful in diabetes management; their aqueous/alcoholic extract shows hypoglycemic effect [65]. Green Tulsi (*Ocimum sanctum*) leaves extract also reduces blood sugar significantly by cortisol inhibiting potency as proved in both normal and alloxan induced diabetic rats [66-67].

### 3.3 In Parkinson's disease

Latif S *et al* concluded that diet enriched with vitamin E may decreases the chances of Parkinson's disease [68] while Brower V reported that creatine is helpful in management of Parkinson's disease by decreasing the clinical symptoms [69].

Antioxidant vitamin supplements such as tocopherol, ascorbic acid and beta-carotene are the abundantly occurring nutraceuticals. As per various earlier literature vitamin E supplements are becoming popular in treatment of Parkinson's disease, whereas epidemiological studies reported that vitamins C and E rich diets are associated with decreased risk of Parkinson's disease [70-71].

### 3.4 In Alzheimer's disease

Literatures have been reported that fulvic acid, an active principle of Shilajit is highly effective against brain disorders exclusively and in combination with vitamin B complex. [72] Patients suffering from Alzheimer's disease treated with donepezil and vitamin E found effective although future study was suggested to check and compare additive as well as individual effect. [73] Wettstein A. *et al* reported that mild to moderate Alzheimer's dementia could be treated with metrifonate, donepezil, rivastigmine which are second-generation cholinesterase inhibitors [74].

Hager *et al* found stable cognitive function especially in those patients who were administered with 600 mg Alpha-lipoic acid along with acetyl-cholinesterase inhibitors, in comparison to those patient who only received therapy of of standard acetyl-cholinesterase inhibitors since last 337 days [75]. Huperzine-alpha is a plant alkaloid derived from club moss plant (*Huperzia serrata*), which is a member or the Lycopodium species. Huperzine-alpha is in phase III clinical trial in the USA and is available as a dietary supplement [76]. The meta-analysis of Huperzine A reported here highlights that this treatment has certain significant improvement for patients with Alzheimer's disease



and Vascular Dementia, and longer durations may result in better efficacy for patients with Alzheimer's disease [77].

Literature survey supports that Lipoic acid also helps to improve potential of mitochondrial membrane, memory loss due to ageing and brain ailments as well as in patient suffering from Parkinson's and Alzheimer's disease [78].

### 3.5 In Erectile Dysfunctioning

L-arginine in combination with **pycnogenol, a product obtained from the pine bark (*Pinus pinaster*)**, is found safe and effective in mild to moderate erectile dysfunction in Japanese patients [79]. When Patients suffering from moderate to severe ED and dyslipidemia were kept on Niacin rich diet, significant improvement was observed in patients [80].

***Kaempferia Parviflora*** Wall. Ex. Baker (KP), which is a Thai plant with name, Kra-Chai-Dum and rhizomes of which are used as a traditional medicine to alleviate male impotency, improve male libido, as energizer, control blood pressure and also reduce stomachache. Study reveals that KP is a potential nutraceutical compound effective in male erectile dysfunction caused due to ageing [81].

### 3.6 Osteoarthritis

Chondroitin sulfate and Glucosamine both are commonly used to **alleviate/reduce** the symptoms of osteoarthritis. Chondroitin sulfate and Glucosamine both act as nutraceuticals and possible mechanism of their anti-inflammatory activity may be due to synthesis of NO and PGE2 [82]. **Capsaicin reduces pain and stiffness and increases joint functioning by acting as agonist for transient receptor potential vanilloid 1 (pain receptor) [82]. Boswellia serrata relieved joint pain, reduced joint swelling, and stiffness by Inhibiting TNF- $\alpha$ -induced MMP-3 expression and protected against IL-1 $\beta$ -induced chondrocyte death [83-85]. Capsaicin reduces pain and stiffness to increased joint function by agonising transient receptor potential vanilloid 1 (pain receptor) while prolonged exposure of capsaicin leads to desensitization of this pain pathway [86-87].**

**Cat's claw reduce osteoarthritis associated pain by Inhibiting lipo-polysaccharide induced PGE2 production and activation of TNF- $\alpha$ . Avocado/soybean unsaponifiables reduced pain in osteoarthritis patients and reduces NSAID consumption by suppressing TNF- $\alpha$ , IL-1 $\beta$ , COX-2, and iNOS in LPS-activated chondrocytes [88-89]. Collagen hydrolysates alleviate osteoarthritis related pain by stimulating regeneration of type II collagen and by increasing biosynthesis of proteoglycans [90].**

### 4. Nutraceuticals tailoring genes- Nutrigenomics

Nutrigenomics is an emerging field to know interactions between food and genes, due to manipulated diet [91]. Nutrigenomics is the study of the use of functional genomic tools to probe a biological system followed by a nutritional stimulus to understand how nutritional molecules may affect metabolic pathways and homeostatic control [92]. Study of effect of nutrigenomics also becomes necessary to find out the effect of foods

on that may interact with particular genes to increase risk of diseases like diabetes mellitus, obesity, cardiovascular diseases etc [93]. Hence it is a vast field to describe the impact of food on genes of human being and requires lot of study to be done.

## Conclusion

In the present scenario nutraceuticals have become more popular in modern society and became important due to increasing applications of Nutraceuticals which serves as a part of growing pharmaceutical industry. Modern society now is being aware about the food products that are beneficial for them in aspects of health and nutrition owing to very few or no side effects. Nutraceuticals are being used to treat fatal diseases like cancer, diabetes, cardiovascular diseases; as well as Parkinson's disease, osteoarthritis etc.

**Table 2: List of Marketed Nutraceuticals**

<b>Products</b>	<b>Category</b>	<b>Contents</b>	<b>Manufacturer</b>
Alamin SE	Protein supplement	L-Arginine & other Protein	Albert David Ltd., India
Albumen Care	Protein supplement	L-arginine, multivitamins & minerals	B.V. Bio-Corp Pvt. Ltd., India
Arginine	Protein Supplement	L-arginine, proantho- cyanidis	Manikind Pharma Pvt. Ltd, India
Appetite Intercept™	Appetite suppressant	Caffeine, tyrosine and Phenylalanine	Natrol, Chatsworth, CA, USA
Betafactor™	Immune supplement	Beta glycan	Ameridan International Inc. USA
Brainspeed Memory®	Brain Health supplement	Vitamin and minerals	Natrol, Chatsworth, CA, USA
Biovinca™	Neurotonic	Vinpocetine	Cyvex nutrition, Irvine, USA
Coral calcium	Calcium supplement	Calcium and trace minerals	Nature's answer, Hauppauge, NY, USA
Calcirol D-3®	Calcium supplement	Calcium and vitamins	Cadilla healthcare, India
Chaser™	Hangover supplement	Activated calcium carbonate, and vegetable carbon	Living essentials, Walled lake MI, USA
GRD®	Nutritional supplement	Proteins, vitamins, minerals and carbohydrates	Zydus Cadila Ltd. Ahmedabad, India
Weight smart™	Nutritional supplement	Vitamins and trace elements	Bayer corporation, Morristown, NL, USA
Yakult	Probiotic dairy product	Skimmed milk, <i>Lactobacilli casai Shirota</i>	Danone india Ltd. India
ImmunAge	Fermented papaya Preparation	Papaya, yeast, dextrose	Osato laboratory Inc. USA
Glowelle®	Beauty drink	Antioxidants, vitamins and fruit extracts	Nestle, India
HiOwna	Nutriional supplement	Protein, multivitamins, minerals and antioxidant	Himalaya herbal ltd, India
PediaSure®	Nutritional supplement	Protein, multivitamins and antioxidant	Abbott India ltd, India
Revital®	Health suppliment	Ginseng, vitamin and minerals	Ranbaxy, India
Proteinex®	Protein supplement	Predigested proteins, vitamins, minerals and carbohydrates	Pfizer Ltd., Mumbai, India
Rox® Glucon-D Glucose-D	Energy drink	Taurine, caffeine and glucuronolactone Glucose	Rox America, Spartanburg, SA, USA Dabur
Omega woman	Immune supplement	Antioxidants, vitamins and. phytochemicals (eg. Lycopene, and resveratrol)	Wassen, Surrey, U.K
Mushroom optimizer™	Immune supplement	Mushrooms polysaccharides and Folic acid	Jarrow formulas, Los Angeles,CA, USA
Proplus®	Nutritional supplement	Soy proteins	Campbell soup company, Camden, NJ, USA
Snapple-aday™	Meal replacement beverage	Vitamins and minerals	Snapple beverage group, White Plains, NY, USA
WelLife®	Amino acid supplement	Granulated-L-glutamine	Daesang America Inc., Hackensach, NJ, USA
Olivinol™	Dietary supplement	Natural antioxidant, Hydroxytyrosol	Cre Agri, Hayward, CA, USA
Threptin®	Diskettes Protein supplements	Proteins and vitamin B	Raptakos, Brett & Co. Ltd., Mumbai, India
Red bull®	Energy drink	Taurine, Caffeine, Glucuronolactone, b- group vitamins	Austrian red bull GmbH

**Table 3: Detailed Review on Various Nutraceuticals**

<b>Types of Nutraceuticals</b>	<b>Sources</b>	<b>Active Constituents</b>	<b>Applications</b>
<b>Dietary Fibre</b>	Whole grain foods wheat and corn bran, nuts	Insoluble Fibre	Reduce chances of colon or breast cancer (anticancer) [94, 95], maintain health of digestive tract [96]
	Oats, barley	Beta-Glucan	Reduce risk of cardiovascular disease, lower down Low Density Lipids and total cholesterol [97,98]
	Beans e.g. Legumes, oats, barley and some fibrous fruits	Soluble Fibre	Anticancer (Colon Cancer), Digestive [99,100]
<b>Fatty Acids</b>	Salmon and other fish oils	Long chain omega-3 Fatty Acids-DHA/EPA	Reduce risk of CVD, Improve mental, visual functions[101-103]
	Cheese, meat products	Conjugated Linoleic Acid	Improving of body composition, Decreases chances of certain cancers[104-106]
	Fruits	Anthocyanidins	Antioxidant ; reduce risk of cancer[87-109]
	Green Tea	Catechins	Antitumor [110]
<b>Phenolics</b>	Citrus	Flavonoids	Antioxidative activity, Prevention of coronary heart disease, hepato-protective, Effective in inflammation and cancer [111]
	Cocoa, Chocolate, Cranberries & cranberry	Tannins	Anti-microbial, Reduce risk of cardiovascular disease [112]
	Corn, soy, wheat, wood oils	Plant Sterols, Stanol ester	Lower blood cholesterol levels by inhibiting cholesterol absorption [113]
	Tomatoes	Lycopene	Antioxidant, protect against prostate cancer [114]
<b>Carotenoids</b>	Corn, various fruits, egg yolk, spinach	Lutin	Antioxidant, Muscle regeneration, anti cancer activity, protect eyes against age related muscular degenerations, cataract [115, 116]
	Carrots, various fruits (Guava, papaya, Water melon etc) and vegetables (tomatoes etc).	Beta carotene	Antioxidant, protection of cornea against UV light.
	Soya beans	Saponins	Effective against colon cancer, reduces cholesterol level [117]
<b>Probiotics/ Prebiotics</b>	Curd	Lactobacillus	Antibacterial, acute diarrhea [118]
	Whole grains, onions, combination of Pro & Prebiotics	Fructo– oligosaccharides	Improve GI health, restore gut flora [119]
	Grains	Tocotrienols and tocopherols	The growth of diverse tumors cell lines was suppressed via initiation of apoptosis and concomitant arrest of cells in the G1 phase of the cell cycle [120]
<b>Phytochemical</b>	Cereal grain, dairy & egg products and plants oil	Phytosterols	Exhibit antioxidant, anti-inflammatory, anti-neoplastic, anti-pyretic & immune-modulating activity, decrease cholesterol [121-123]
	Various plants, whole grain	Phenolic constituents	Antioxidants, Anti-hyperglycemic, and anti hypertensive [124]
	Grapes, berries, cocoa, green tea, acacia spp.	Catechin & gallic acids	Antioxidants, Antiradical property, cyto-protective.[ 125, 126]
	Soybeans	Isoflavonoids	Treating cancers & attenuates bone loss [127, 128]

## References

1. Bagchi D. Nutraceuticals and functional foods regulations in United States and around the world. *Toxicol.* 2006; 221: 1-3.
2. Granato H. Regulatory Concerns Cloud Functional Food, Nutraceutical Markets. c 2009  
<http://www.naturalproductsinsider.com/articles/2000/11/regulatory-concerns-cloud-functional-food-nutraceutical.aspx>
3. Ross S. Functional foods: the Food and Drug Administration perspective. *Am J Clin Nutr.* 2000; 71: 1735-38.
4. Palthur, MP, Palthur SSS, Chitta SK. Nutraceuticals: Concept and regulatory scenario. *Int J Pharm Pharm Sci.* 2010; 2: 14-20.
5. Cindy, Oliveri S. Nutraceuticals, Phytochemicals, and Antioxidants-What Are They All About. OSU Extension Fact Sheet. 2003; 5051-98
6. Dietary Supplements What You Need to Know, USFDA, May 2006  
<http://www.fda.gov/downloads/Food/DietarySupplements/UCM240978.pdf> (last cited on Jan 20, 2016).
7. Dietary supplement health and education act 1994, public law 103-417, 103<sup>rd</sup> congress [http://ods.od.nih.gov/pdf/About/DSHEA\\_Wording.aspx](http://ods.od.nih.gov/pdf/About/DSHEA_Wording.aspx) (last cited on Jan 20, 2016).
8. <http://www.eufic.org/article/en/expid/basics-functional-foods/> (last cited on Jan 20, 2016).
9. Hathcock J. Dietary supplements: How they are used and regulated. *J Nut.* 2001; 131: 1114-7.
10. Dureja H, Kaushik D, Kumar V. Developments in nutraceuticals. *Ind J Pharmacol.* 2003; 35: 363-72.
11. Hardy G. Nutraceuticals and functional foods: introduction and meaning. *Nutrition.* 2000; 16(7-8): 688-9.
12. Grace, Emily. How to choose the best supplement. Health Beacon. Accessed on 3 march 2016.
13. Camire ME, Kantor MA. Dietary Supplements: Nutritional and Legal Considerations, *Food technol.* 1999; 53(7): 87-96.
14. Whitney, Elano, Sharo R. Understanding nutrition. 10th ed. Thomson-Wadsworth; 2005.

- 361 15. Frances S, Ellie W. Nutrition: concepts and controversies. 1<sup>3th</sup> ed. Cengage learning  
362 2007.
- 363 16. Hamid AA, Luan YS. Functional properties of dietary fiber prepared from defatted  
364 rice bran. Food Chem. 2000; 68: 15 - 9.
- 365 17. Mermel V. Old paths new directions: use of functional foods in the treatment of  
366 obesity. Trends Food Sci Tech. 2004; 15: 532-40.
- 367 18. Dulloo AG, Duret C, Rohrer D. Efficacy of a green tea extract rich in catechin  
368 polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in  
369 humans. Am J Clin Nutr. 1999; 70: 1040-5.
- 370 19. Ehrlich SD. (Willow bark), private practice specializing in complementary and  
371 alternative medicine Phoenix AZ Review. VeriMed Healthcare Network, 2008.
- 372 20. Cora JD, German JB. Phytochemicals: nutraceuticals and human health. J Sci Food  
373 Agric, 2000; 80:1744-56.
- 374 21. FAO/WHO (2001) Joint FAO/WHO Expert Consultation on evaluation of health and  
375 nutritional properties of probiotics in food including powder milk with live lactic acid  
376 bacteria Cordoba, Argentina, October 2001.
- 377 22. Hord NG. Eukaryotic microbiotic crosstalk: potential mechanisms for health benefits  
378 of prebiotics and probiotics. Annu Rev Nutr. 2008; 28: 215–31.
- 379 23. Michail S, Sylvester F, Fuchs G, Issenma R. Clinical efficacy of probiotics: Review of  
380 the evidence with focus on children, clinical practice guideline. J Pediatr  
381 Gastroenterol Nutr. 2006; 43(4):550-7.
- 382 24. Holzapfel WH, Haberer P, Geisen R, Bjorkroth J, Schillinger U. Taxonomy and  
383 important features of probiotic microorganisms in food and nutrition. Am J Clin Nutr.  
384 2001; 73: 365S-73S.
- 385 25. Gibson GR. Dietary prebiotic: current status and new definition. IFIS Func Foods  
386 Bull. 2011; 7: 1–19.
- 387 26. Nikfar S, Rahimi R, Rahimi F, et al. Efficacy of probiotics in irritable bowel syndrome:  
388 a meta-analysis of randomized, controlled trials. Dis Colon Rectum. 2008;  
389 51:1775–80.
- 390 27. McFarland LV, Dublin S. Meta-analysis of probiotics for the treatment of irritable  
391 bowel syndrome. World j Gastroenterol. 2008; 14: 2650-61.
- 392 28. AlFaleh K, Anabrees J. Probiotics for prevention of necrotizing enterocolitis in  
393 preterm infants. Cochrane Database Syst Rev. 2014; 10; 4.

- 394 29. Hamilton-Miller J.M.T. The role of probiotics in the treatment and prevention of  
395 *Helicobacter pylori* infection. Int J Antimicrob Agent. 2003;22 (4): 360–66.
- 396 30. Stapleton AE, Au-Yeung M, Hooton TM. Randomized, Placebo-Controlled Phase 2  
397 Trial of a *Lactobacillus crispatus* Probiotic Given Intravaginally for Prevention of  
398 Recurrent Urinary Tract Infection. Clin Infect Dis. 2011; 52 (10):1212-17.
- 399 31. Strus M, Kucharska A, Kukla G, Brzychczy-Włoch M. The in vitro activity of  
400 vaginal lactobacillus with probiotic properties against candida. Inf Dis Obstet  
401 Gynecol  
402 2005; 13(2): 69-75.
- 403 32. Ignarro LJ, Balestrieri ML, Napoli C. Nutrition, physical activity, and cardiovascular  
404 disease: an update. Cardio Res. 2007; 73: 326-40.
- 405 33. Trumbo PR, Ellwood KC. Supplemental calcium and risk reduction of hypertension.  
406 Nutr Rev. 2007; 65: 78-87.
- 407 34. Hatton DC, Yue Q, McCarron DA. Mechanisms of calcium's effects on blood  
408 pressure. Semin Nephrol. 1995;15(6):593-602.
- 409 35. Ravi SMT. Nutrigenetics and nutraceuticals: the next wave riding on personalized  
410 medicine. Trans Res. 2007; 149:55-61.
- 411 36. Freedman JE, Farhat JH, Loscalzo J, *et al.* Alpha-tocopherol inhibits aggregation of  
412 human platelets by a protein kinase C-dependent mechanism. Circulation 1996;  
413 94:2434–40.
- 414 37. German JB, Walzem RL. The health benefits of wine. Ann Rev Nutr. 2000; 20: 561-  
415 93.
- 416 38. Hollman PCH, Hertog MGL, Katan MB. Analysis and health effects of flavonoids.  
417 Food Chem. 1996; 57:43- 6.
- 418 39. Yao LH, Jiang YM, Shi J, Tomás-Barberán FA, Datta N, Singanusong R, Chen SS.  
419 Flavonoids in food and their health benefits. Plant Foods Hum Nutr. 2004; 59(3):  
420 113-22.
- 421
- 422 40. Cook NC, Samman S. Flavonoids - Chemistry, metabolism, cardio-protective effects,  
423 and dietary sources. J Nutri Biochem. 1996;7:66-76.
- 424 41. Hollman PCH, Feskens EJ, Katan MB. Tea flavonols in cardiovascular disease and  
425 cancer epidemiology. Proc Soc Exper Biol Med. 1999; 220:198-202.
- 426 42. Rajasekaran A, Sivagnanam G, Xavier R. Nutraceuticals as therapeutic agents: A  
427 Review. Res J Pharm Tech. 2008; 1(4):328-40.



- 428 43. Niki E, Noguchi N, Tsuchihashi H, Gotoh N. Interaction among vitamin C, vitamin E,  
429 and beta-carotene. *Am J Clin Nutr.* 1995; 62(6Suppl):1322S-6S.
- 430 44. Zahra B, Parvin M, Fereidoun A. Dietary polyphenols as potential nutraceuticals in  
431 management of diabetes: a review. *J Diabetes Metab Disord.* 2013; 12(1):43.
- 432 45. Bunyapraphatsara N, Yongchaiyudha S, Rungpitarangsi V, Chokechaijaroenporn O.  
433 Antidiabetic activity of Aloe vera L juice II Clinical trial in diabetes mellitus patients in  
434 combination with glibenclamide. *Phytomed.* 1996;3(3): 245-8.
- 435 46. Philips AO, Philips GO. Biofunctional behavior and health benefits of a specific Gum  
436 Arabic. *Food hydrocoll.* 2011; 25(2): 165-9.
- 437 47. Hou CC, Lin SJ, Cheng JT, Hsu FL. Antidiabetic dimeric guianolides and a lignan  
438 glycoside from *Lactuca indica*. *J Nat Prod.* 2003; 66(5):625-9.
- 439 48. Wadood A, Wadood N, Shah SA. Effects of *Acacia arabica* and *Caralluma edulis* on  
440 blood glucose levels of normal and alloxan diabetic rabbits. *J Pak Med Assoc.* 1989;  
441 39(8): 208-12.
- 442 49. Takii H, Kometani T, Nishimura T, Nakae T, Okada S, Fushikii T. Antidiabetic effect  
443 of glycyrrhizin in genetic diabetes *Licorice* in diabetes. *Biol Pharma Bull.* 2000; 24:  
444 484-7.
- 445 50. Brouns F. Soya isoflavones: a new and promising ingredient for the health foods  
446 sector. *Food Res Int.* 2002; 35: 187-93.
- 447 51. Thomas B, Ghebremeskel K, Lowy C, Crawford M, Shore R N. Nutrient intake of  
448 women with and without gestational diabetes with a specific focus on fatty acids.  
449 *Nutrition.* 2006; 22: 230-6.
- 450 52. Sirtori CR, Galli C. Fatty acids and the Omega 3. *Biomed Pharmacother.* 2002; 56:  
451 397-406.
- 452 53. Coleman MD, Eason RC, Bailey CJ. The therapeutic use of lipoic acid in diabetes: a  
453 current perspective. *Envir Toxi Pharmacol.* 2001; 10: 167-72.
- 454 54. Singh U, Jialal L. Alpha-lipoic acid supplementation and diabetes. *Nutr Rev.*  
455 2008;66: 646–57.
- 456 55. Ziegler D, Reljanovic M, Mehnert H, Gries FA. Alpha-lipoic acid in the treatment of  
457 diabetic polyneuropathy in Germany: Current evidence from clinical trials. *Exp Clin*  
458 *Endo Dia.* 1999;107: 421–30.
- 459 56. Baljit S. Psyllium as therapeutic and drug delivery agent. *Int J Pharm.* 2007; 334:1-  
460 14.

- 461 57. Lau FC, Bagchi M, Sen CK, Bagchi D. Nutrigenomic basis of beneficial effects of  
462 chromium (III) on obesity and diabetes. *Mol Cell Biochem.* 2008; 3:171–10.
- 463 58. McCarty MF. Nutraceutical resources for diabetes prevention—an update. *Med*  
464 *Hypo.* 2005; 64:151–8.
- 465 59. Dakshinamurti K, Cheah-Tan C. Biotin-mediated synthesis of hepatic glucokinase in  
466 the rat. *Arch Biochem Biophys.* 1968; 127: 17-21.
- 467 60. Furukawa Y. Enhancement of glucose-induced insulin secretion and modification of  
468 glucose metabolism by biotin. *Nippon Rinsho.* 1999; 57: 2261-9.
- 469 61. Wang ZQ, Zhang XH, Cefalu WT. Chromium picolinate and biotin enhance glycogen  
470 synthesis and glycogen synthase gene expression in human skeletal muscle culture.  
471 *Diab Res Clin Pract.* 2000; 50: 395.
- 472 62. Chattopadhyay RR, Chattopadhyay RN, Nandy AK, Poddar G, Maitra SK. The effect  
473 of fresh leaves of *Azadirachta indica* on glucose uptake and glycogen content in the  
474 isolated rat hemi diaphragm. *Bull Cal Sch Trop Med.* 1987; 35: 8–12.
- 475 63. Chattopadhyay RR. Possible mechanism of antihyperglycemic effect of *Azadirachta*  
476 *indica* leaf extract part IV. *Gen Pharmacol.* 1996; 27:431–4.
- 477 64. Chattopadhyay RR. A comparative evaluation of some blood sugar lowering agents  
478 of plant origin. *J Ethnopharmacol.* 1999;67:367–72.
- 479 65. Gholap S, Kar A. Hypoglycaemic effects of some plant extracts are possibly  
480 mediated through inhibition in corticosteroid concentration. *Pharmazie.*  
481 2004;59:876–8.
- 482 66. Sharma SB, Nasir A, Prabhu K, Murthy PS, Dev G. Hypoglycaemic and  
483 hypolipidemic effect of ethanolic extract of seeds of *Eugenia jambolana* in alloxan-  
484 induced diabetic rabbits. *J Ethnopharmacol.* 2003;85:201–6.
- 485 67. Rai V, Iyer U, Mani UV. Effect of *Tulsi* (*Ocimum sanctum*) leaf powder  
486 supplementation on blood sugar levels serum lipids and tissue lipids in diabetic rats.  
487 *Plant Foods Hum Nutri.* 1997; 50:9–16.
- 488 68. Latif S, Anwar F, Ashraf M, Gilani AH. *Moringa oleifera*: a food plant with multiple  
489 medicinal uses. *Phytother Res.* 2007; 21:17-25.
- 490 69. Brower VA. Nutraceutical a day may keep the doctor away. *EMBO reports.* 2005;  
491 8:708- 11.
- 492 70. Maxwell SRJ. Antioxidant vitamin supplements: update of their potential benefits and  
493 possible risks. *Drug Saf.* 1999; 21:253–66.

- 494 71. Anderson C, Checkoway H, Franklin GM et.al, Dietary factors in Parkinson's  
495 disease: the role of food groups and specific foods. *Mov Disord.* 1999; 14:21–7.
- 496 72. Carlos C, Gonzalo F, Patricio F, Fernando C, Ricardo M. Can Nutraceuticals  
497 Prevent Alzheimer's Disease? Potential Therapeutic Role of a Formulation  
498 Containing Shilajit and Complex B Vitamins. *Arch Med.* 2012; 43(8): 699–704.
- 499 73. Klatte ET, Scharre DW, Nagaraja HN, Davis RA, Beversdorf DQ. Combination  
500 therapy of donepezil and vitamin E in Alzheimer disease *Alzheimer. Dis Asso*  
501 *Disord.* 2003; 17: 113-6.
- 502 74. Wettstein A, Stadtärztllicher DZ. Cholinesterase inhibitors and Ginkgo extracts--are  
503 they comparable in the treatment of dementia? Comparison of published placebo-  
504 controlled efficacy studies of at least six months' duration. *Phytomedicine.* 2000;  
505 6(6): 393-401.
- 506 75. Hager K, Marahrens A, Kenklies M, Riederer P, Munch G. Alpha lipoic acid as a new  
507 treatment option for Alzheimer type dementia. *Arch Gerontol Geriatr.* 2001;32:275-  
508 82.
- 509 76. Grunwald J, Raveh L, Doctor BP. Huperzine A as a pretreatment candidate drug  
510 against nerve agent toxicity. *Life Sci.* 1994; 54: 991-7.
- 511 77. Shu-huai X, Chun-Xiao Z, Rui Z, Li An. Huperzine A in the Treatment of Alzheimer's  
512 Disease and Vascular Dementia: A Meta-Analysis. *Evid Based Complement Alternat*  
513 *Med.* 2014 Article ID 363985 10 <http://dxdoiorg/101155/2014/363985>
- 514 78. Kramer K, Elkins MHR. Alpha lipoic acid: Nature's supreme Antioxidant. In: Elkins  
515 MHR. USA: Wood land Publishing; 1998.
- 516 79. Aoki H, Nagao J, Ueda T, Strong JM, Schonlau F, Yu-Jing S, Lu Y, Horie S. Clinical  
517 assessment of a supplement of Pycnogenol and L-arginine in Japanese patients  
518 with mild to moderate erectile dysfunction. *Phytother Res.* 2012; 26(2): 204-7.
- 519 80. Ng CF, Lee CP, Ho AL, Lee VW. Effect of niacin on erectile function in men suffering  
520 erectile dysfunction and dyslipidemia. *J Sex Med.* 2011; 8(10): 2883-93.
- 521 81. Yenjai CK, Prasanphen S, Daodee V, Wongpanich, Kittakoo P. Bioactive  
522 flavonoids from *Kaempferia parviflora*. *Fitoterapia.* 2004; 75(1):89-92.
- 523 82. Kalioraa AC, Dedoussisa GVZ, Schmidt H. Dietary antioxidants in preventing  
524 atherogenesis. *Atherosclerosis.* 2006; 187(1):1-17.
- 525 83. Krishnaraju, AV, Sundararaju D, Vamsikrishna U, Suryachandra R, Machiraju G,  
526 Sengupta, K, Trimurtulu G. Safety and toxicological evaluation of Aflapin: A novel  
527 *Boswellia*-derived anti-inflammatory product. *Toxicol Mech Method.* 2010; 20:556–  
528 63.

- 529 84. Kimmattkar, N, Thawani V, Hingorani, L, Khiyani R. Efficacy and tolerability of  
530 *Boswellia serrata* extract in treatment of osteoarthritis of knee—A randomized  
531 double blind placebo controlled trial. *Phytomedicine*. 2003; 10:3–7.
- 532 85. Sengupta, K, Kolla, JN, Krishnaraju AV, Yalamanchil N, Rao CV, Golakoti T,  
533 Raychaudhuri S, Raychaudhuri SP. Cellular and molecular mechanisms of anti-  
534 inflammatory effect of Aflapin: A novel *Boswellia serrata* extract. *Mol. Cell Biochem*.  
535 2011, 354, 189–97.
- 536 86. Kosuwon, W, Sirichatiwapee, W, Wisanuyotin T, Jeeravipoolvarn P,  
537 Laupattarakasem W. Efficacy of symptomatic control of knee osteoarthritis with  
538 0.0125% of capsaicin versus placebo. *J. Med. Assoc.* 2010; 93:1188–95.
- 539 87. Remadevi R, Szallisi A. Adlea (ALGRX-4975), an injectable capsaicin (TRPV1  
540 receptor agonist) formulation for longlasting pain relief.. *Drugs*. 2008; 11:120–32.
- 541 88. Christensen R, Bartels EM, Astrup A, Bliddal H. Symptomatic efficacy of avocado-  
542 soybean unsaponifiables (ASU) in osteoarthritis (OA) patients: A meta-analysis of  
543 randomized controlled trials. *Osteoarthr. Cartilage*. 2008; 16: 399–408.
- 544 89. Au RY, Al-Talib TK, Au AY, Phan PV, Frondoza CG. Avocado soybean  
545 unsaponifiables (ASU) suppress TNF-alpha, IL-1beta, COX-2, iNOS gene  
546 expression, and prostaglandin E2 and nitric oxide production in articular  
547 chondrocytes and monocyte/macrophages. *Osteoarthr. Cartilage*. 2007; 15:1249–  
548 55.
- 549 90. Bello AE, Oesser S. Collagen hydrolysate for the treatment of osteoarthritis and  
550 other joint disorders: A review of the literature. *Curr Med Res Opin*. 2006; 22: 2221–  
551 32.
- 552 91. Neeha VS, Kint P. Nutrigenomics research: a review. *J Food Sci Technol*. 2013;  
553 50(3):415–28.
- 554 92. Mutch DM, Wahli W, Williamson G. Nutrigenomics and nutrigenetics: the emerging  
555 faces of nutrition. 19 2005. *FASEB Journal*. 1602-16.
- 556 93. Nutrigenomics focuses on the effect of nutrients on the genome, proteome, and  
557 metabolome. Ordovas J & Mooser M. (2004) *Current Opinion in Lipidology* 15:101-8.
- 558 94. Adom KK, Liu RH. Antioxidant activity of grains. *J Agric Food Chem*. 2002; 50 (21):  
559 6182–7.
- 560 95. Ferguson LR, Harris P. Protection against cancer by wheat bran: role of dietary fibre  
561 and Phytochemicals. *Euro J Can Pre*. 1999; 8(1): 17-25.
- 562 96. Engberg RM, Hedemann MS, Steenfeldt S, Jensen BB. Influence of Whole Wheat  
563 and Xylanase on Broiler Performance and Microbial Composition and Activity in the  
564 Digestive Tract. *Poul Sci*. 2004; 83(6): 925-38.

- 565 97. Keogh GF, Cooper GJS, Mulvey TB, McArdle BH, Coles GD, Monro JA, Poppitt SD.  
566 Randomized controlled crossover study of the effect of a highly  $\beta$ -glucan-enriched  
567 barley on cardiovascular disease risk factors in mildly hypercholesterolemic men.  
568 *Am J Clin Nutr.* 2003; 78(4): 711-8.
- 569 98. Theuwissen E, Mensink RP. Water-soluble dietary fibers and cardiovascular  
570 disease. *Physiol Behav.* 2008;94(2):285-92.
- 571 99. McIntyre A, Gibson PR, Young GP. Butyrate production from dietary fibre and  
572 protection against large bowel cancer in a rat model. *Gut.* 1993; 34: 386-91.
- 573 100. Malkki Y, Virtanen E. Gastrointestinal effect of oat bran and oat gum: a review.  
574 *LWT- Food Sci Technol.* 2001;34(6):337-47.
- 575 101. Popp-Snijders C, Schouten JA, Heine RJ, Vander MJ, Vander VE. A Dietary  
576 supplementation of omega-3 polyunsaturated fatty acids improves insulin sensitivity  
577 in non-insulin-dependent diabetes. *Diabetes Res.* 1987; 4(3):141-7.
- 578 102. Connor WE, Importance of n-3 fatty acids in health and disease. *Am J clin Nutr.*  
579 2000; 71(1): 171S-5S.
- 580 103. Young G, Conquer J. Omega-3 fatty acids and neuropsychiatric disorders.  
581 *Reprod Nutrition Dev.* 2005; 45(1): 1-28.
- 582 104. Lee KN, Kritchevsky D, Pariza MW. Conjugated linoleic acid and  
583 atherosclerosis in rabbits. *Atherosclerosis.* 1994;108(1):19-25.
- 584 105. Clement I, Banni S, Angioni E, Carta G, McGinley J, Thompson HJ, Barbano  
585 D, Bauman D. Conjugated Linoleic Acid-Enriched Butter Fat Alters Mammary Gland  
586 Morphogenesis and Reduces Cancer Risk in Rats. *J Nutr.* 1999;  
587 129(12): 2135-42.
- 588 106. Shultz TD, Chewab BP, Seamana WR, Luedeck LO. Effect of conjugated linoleic  
589 acid on body composition in mice. *Cancer Lett.* 1992; 63(2):125-33.
- 590 107. Wang H, Cao G, Prior RL. Oxygen Radical Absorbing Capacity of Anthocyanins.  
591 *J Agric Food Chem.* 1997;45(2):304-9.
- 592 108. Kanga SY, Seeramb NP, Narib MG, Bourquin LD. Tart cherry anthocyanins  
593 inhibit tumor development in ApcMin mice and reduce proliferation of human colon  
594 cancer cells. *Cancer Lett.* 2003;194(1):13-9.
- 595 109. Li-Shu W, Stoner GD. Anthocyanins and their role in cancer prevention. *Cancer*  
596 *Lett.* 2008;269(2):281-90.
- 597 110. Liang G, Tang A, Lin X, Li L, Zhang S, Huang Z, Tang H, Li QQ. Green tea  
598 catechins augment the antitumor activity of doxorubicin in an in vivo mouse model  
599 for chemoresistant liver cancer. *Int J Oncol.* 2010; 37(1):111-23.

- 600 111. Kumar S, Pandey AK. Chemistry and Biological Activities of Flavonoids: An  
601 Overview. *Scientific World J.* 2013; 1-16.
- 602 112. Lim H, Darah I, Jain K. Antimicrobial activities of tannins extracted from  
603 *rhizophora apiculata* barks. *J Trop Forest Sci.* 2006; 18(1):59—65.
- 604 113. Katan MB, Grundy SM, Jones P. Efficacy and Safety of Plant Stanols and Sterols  
605 in the Management of Blood Cholesterol Levels. *Mayo Clin Proc.* 2003; 78(8):965–  
606 78.
- 607 114. Takeoka GR, Dao L, Flessa S, Gillespie DM, Jewell WT, Huebner B, Bertow D,  
608 Ebeler SE. Processing Effects on Lycopene Content and Antioxidant Activity of  
609 Tomatoes. *J Agric Food Chem.* 2001; 49(8):3713–17.
- 610 115. Naguib YMA. Antioxidant Activities of Astaxanthin and Related Carotenoids. *J*  
611 *Agric Food Chem.* 2000; 48(4):1150–54.
- 612 116. Olmedilla B, Granado F, Blanco I, Vaquero M, Cajigal C. Lutein in patients with  
613 cataracts and age-related macular degeneration: a long-term supplementation study.  
614 *J Sci Food Agri.* 2001;81(9):904–9.
- 615 117. Kim HY, Yu R, Kim JS, Kim YK, Sung MK. Antiproliferative crude soy saponin  
616 extract modulates the expression of IκBα protein kinase C and cyclooxygenase-2 in  
617 human colon cancer cells. *Cancer Lett.* 2004; 210(1):1–6.
- 618 118. Isolauri E, Rautanen T, Juntunen M, Sillanaukee P, Koivula T. A  
619 Human *Lactobacillus* Strain (*Lactobacillus Casei* sp strain GG) Promotes Recovery  
620 From Acute Diarrhea in Children. *Pediatric.*1991; 88(1):90 -7.
- 621 119. Fuller R, Gibson GR. Probiotics and prebiotics: microflora management for  
622 improved gut health. *Clinic Microbiol Infec.* 1998; 9:477–80.
- 623 120. Ju J, Picinich SC, Yang Z. Cancer-preventive activities of tocopherols and  
624 tocotrienols. *Carcinogenesis.* 2010; 31(4):533–42.
- 625 121. Wang T, Hicks KB, Moreau R. Antioxidant activity of phytosterol oryzanol and  
626 other phytosterol conjugates. *J Ame Oil Chem Soci.* 2002; 79(12):1201-6.
- 627 122. Rozner S, Garti N. The activity and absorption relationship of cholesterol and  
628 phytosterols. *Colloids and Surfaces A: Physicochem Engg Aspec.* 2006; 282-  
629 283:435-56.
- 630 123. Garcia MD, Saenz MT, Gomez MA. Topical anti-inflammatory activity of  
631 phytosterol isolated from *Ezyngium foetidum* on chronic and acute inflammatory  
632 model. *Phytother Res.*1999; 13(1):78-80.
- 633 124. Ranilla LG, Kwon Y, Apostolidis E, Shetty K. Phenolic compounds antioxidant  
634 activity and in vitro inhibitory potential against key enzymes relevant for

635 hyperglycemia and hypertension of commonly used medicinal plants herbs and  
636 spices in Latin America. *Bioresour Tech.* 2010; 101(12):4676–89.

637 125. Morel I, Lescoat G, Cogrel P, Sergent O, Pasdeloup N. Antioxidant and iron-  
638 chelating activities of the flavonoids catechin quercetin and diosmetin on iron-loaded  
639 rat hepatocyte cultures. *Biochem Pharmacol.* 1993; 45(1):13-9.

640 126. Kroes BH, Vanden BAJ, Quarlesvan UHC, Labadie RP. Anti-inflammatory activity  
641 of gallic acid. *Planta Medica.* 1992; 58(6):499-504.

642 127. Birt DF, Hendrich S, Wang W. Dietary agents in cancer prevention: flavonoids  
643 and isoflavonoids. *Pharmacol Thera.* 2001; 90(2–3):157–77.

644 128. Alekel DL, St G, Peterson CT, Hanson KB, Stewart, Toda T. Isoflavone-rich soy  
645 protein isolate attenuates bone loss in the lumbar spine of perimenopausal women.  
646 *Am J Clin Nut.* 2000; 72(3):844-52.

647

648