

Original Research Article

Analysis of Bioactive Compounds in Latex of *Croton bonplandianum* (Baill.) using GC-MS

ABSTRACT

Background: *Croton bonplandianum* Baill. belongs to the family of Euphorbiaceae and commonly known as 'Bantulsi'. Traditionally, the latex of this plant is used for cuts and wounds.

Objective: The present investigation was designed to determine the bioactive constituents of latex of *C. bonplandianum* by GC-MS. In this study, the mass spectrum of the compounds found in the ethanolic extract of latex was matched with the National Institute of Standards and Technology (NIST) library.

Place and Duration of Study: Post Graduate and Research Department of Biochemistry at Government Arts College (Autonomous), Kumbakonam and Centre for Advanced Research in Indian System of Medicine (CARISM), SASTRA University, Tirumalaisamudram, Thanjavur, Tamilnadu, India, during the months between October to December, 2015.

Methodology: 10 ml of fresh latex was mixed with 10 ml of ethanol and kept at shaker for 3 hours. The sample was filtered and concentrated through nitrogen flushing to 1 ml. From this, 2 µl of prepared sample was injected into GC-MS instrument for phytochemical analysis.

Results: The GC-MS analysis of the ethanolic extract of latex of *C. bonplandianum* revealed that the presence of twenty three phytochemicals including the anti-inflammatory, antioxidant and antimicrobial compounds but not all of them are bioactive compounds. The major chemical constituents present in the latex are 2-C-methyl myoinositol (32.4), mequinol (0.74), 4-methyl-phenol (6.86), 1,2,3-benzenetriol (3.54), 3-methyl-quinoline, (0.44), n-hexadecanoic acid [palmitic acid] (5.36) and octadecanoic acid [stearic acid] (1.95).

Conclusion: Further study is needed to isolate, identify, characterize and elucidate the structure of bioactive compounds responsible for therapeutic values of latex of *C. bonplandianum*.

Keywords: *Croton bonplandianum*, latex, GC-MS, ethanolic extract, Bantulsi

1. INTRODUCTION

Medicinal plants have been used by human being since ages in traditional medicine due to their therapeutic potential. The search on medicinal plants has led to the discovery of novel drug candidates used against diverse diseases [1]. According to the World Health Organization [2], more than 80% of the world's population relies on traditional medicine for their primary healthcare needs. Phytochemicals are natural bioactive compounds found in plants such as vegetables, fruits, medicinal plants, flowers, leaves and roots that act among these as a defense system against diseases [3]. Screening of biologically active medicinal compounds has been conducted on well-known species of plants used in traditional medicines [4]. In the recent years GC-MS has been applied to unambiguously identify the structures of different phytoconstituents in plant extracts and biological samples with great success [5,6]. GC-MS is one of the best techniques to identify the constituents of volatile matter such as long and branched chain hydrocarbons, alcohols, acids, esters etc [7]. The phytochemical screening by GC-MS analysis proved that the plants are pharmaceutically important due to presence of different medicinally important phytochemical compounds and secondary plant metabolites. The GC-MS analysis of the plants showed that the presence of important bioactive compounds.

The plant latex is a natural polymer composed of 320 to 35,000 isoprene molecules and it is stored in specialized cells called laticifer [8,9]. The latex is known to be phytochemically diverse that contains complex mixtures of terpenoids, phenolics, proteins and alkaloids [10]. Although, all parts of the plant

have been used for medicinal purposes [11,12]. Comprehensive literature survey on the plant latex revealed that it contains wide diversities of defense chemicals and proteins, which displayed different biological activities [13,14]. Moreover, some compounds with wound healing properties were reportedly found in the plant latex [11]. *C. bonplandianum* belongs to the family of Euphorbiaceae. It is widely distributed throughout the plains of India. It is a small perennial herb, green and laticiferous growing upto 1-2 ft tall [15]. *C. bonplandianum* is also well known for its medicinal value. Seeds of *C. bonplandianum* are used for the treatment of jaundice, acute constipation, abdominal dropsy and internal abscesses [16]. Its ethanolic extract has been found to possess hypotensive, spasmolytic and antimicrobial activities [17,18]. The latex of *C. bonplandianum* is used to treat cuts and wounds traditionally [19,20]. The latex of *C. bonplandianum* (1:5 v/v in 50% acetone) showed antifungal activity [21]. Many pharmacological studies are available using latex of *C. bonplandianum*. But there is no study on the screening of bioactive compounds of latex of *C. bonplandianum*. Therefore, the present study was aimed to investigate the phytoconstituents of ethanolic extract of latex of *C. bonplandianum* by GC-MS.

2. MATERIALS AND METHODS

2.1. Collection of Plant Material

C. bonplandianum latex was collected from the waste lands in and around Mayiladuthurai, Nagapattinam District, Tamilnadu, India, where it was found naturally. The plant was identified by Rev. Dr. S. John Britto, Director, Rabinat Herbarium and Centre for Molecular Systematics, St. Joseph's College, Tiruchirappalli, Tamilnadu, India. The latex of *C. bonplandianum* was obtained as exudates by and plucking of fresh leaves from actively growing plants. The latex was collected into sterile, plastic containers by pressing and squeezing in between fingers, the apex of the leaves to release as much as possible latex into the containers. The collections of latex were made in the mornings [22]. After collection, the latex was centrifuged with 5000 rpm for 15 minutes to remove any solid particles present in it. The latex was decanted off into containers and plugged with cotton and stored at 4°C until the time of use.

2.2. Identification of Phytochemicals by GC-MS

2.2.1. Sample Preparation

10 ml of fresh latex was mixed with 10 ml of ethanol and kept it in the orbital shaker for overnight [23]. The sample was filtered and concentrated through nitrogen flushing to 1 ml. From this, 2 µl of sample was injected into GC-MS instrument for phytochemical analysis.

2.2.2. Equipment

GC-MS analysis was carried out by using a Perkin Elmer Clarus 500 for the screening of phytochemicals of *C. bonplandianum* [24]. The data were obtained on Capillary Column Elite- 5MS (5% phenyl 95% dimethyl poly siloxane). Helium was used as the carrier gas with a flow rate of 1ml/min in the split mode (10:1). An aliquot of 2 µl of ethanol solution of the sample was injected into the column with the injector temperature at 250°C. GC oven temperature started at 110°C and holding for 2 min and it was raised to 200°C at the rate of 10°C/min without holding. Holding was allowed at 280°C for 9 min with program rate of 5°C/min. The injector and detector temperatures were set at 250°C and 280°C, respectively. GC interface and Ion source temperature was maintained at 200°C. The mass spectrum of compounds in the sample was obtained by electron ionization at 70 eV and the detector was operated in scan mode from 40-450 amu (atomic mass units). A scan interval of 0.5 second and fragments from 40 to 450 Da were maintained. The total running time was 36 minutes.

2.2.3. Interpretation of Mass Spectrum (MS)

In the MS Programme, NIST Version 2.0 library database of National Institute of Standard and Technology (NIST) having more than 2, 00,000 patterns were used for identifying the phytochemicals of *C. bonplandianum*. The spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library. The name, molecular weight and structure of the components of the *C. bonplandianum* were ascertained.

3. RESULTS AND DISCUSSION

Nowadays the search on the phytochemicals of medicinal plants with their activity has increased. The combination of the best separation technique (GC) with the best identification technique (MS) made GC-MS and it is an ideal technique for screening of volatile and semi-volatile bioactive compounds [25]. In the present study, twenty three bioactive chemical compounds in the latex of *C. bonplandianum* were identified and their names, retention time (RT), peak area percentage and molecular weight are represented in Table 1. The highest peak area percentage was obtained by 2-C-methyl myoinositol (32.4%) with the retention time of 10.48 and the lowest peak area percentage was obtained by N-ethyl-4-methoxy-1,3,5-Triazin-2-amine (0.07%) with the retention time of 23.31. The most abundant components were found in the latex are 2-C-methyl-myoinositol (42.40%), 13-octadecenal (23.31%), 2-methoxy-4-methylbenzoic acid (10.90%), D-allose (10.63%) and 4-methyl-phenol (6.86%).

In our previous study, 10 ml of latex was mixed with 90 ml of ethanol and prepared the extract as mentioned in the section 2.2.1 and then carried out GC-MS analysis. In that study, the latex extract of *C. bonplandianum* showed the presence of only one compound 2-C-methyl myoinositol (30.8%) [26]. But in the present study, the extract of latex of *C. bonplandianum* showed that the presence of twenty three compounds including 2-C-methyl myoinositol (32.4%). This variation may be due to the dilution of the latex with ethanol because in this study the latex was mixed with equal volume of ethanol, but in the previous study the latex was mixed with 9 fold volume of ethanol for extract preparation. So, the present study confirmed that the dilution of the latex play a main role in the presence of phytochemicals and appearance of peaks in the GC-MS chromatogram.

The biological activities of identified phytochemicals in the latex extract of *C. bonplandianum* were predicted based on Dr. Duke's phytochemical and ethnobotanical databases [27]. Among the identified phytochemicals in the latex of *C. bonplandianum*, a few compounds like 2-C-methyl myoinositol (32.4%), 3-methyl-quinoline (0.44%), 4-amino-6-methyl-piperidin-2-one (5.24%), 1,2,3-benzenetriol (3.54%), 2-methoxy-4-methyl-benzoic acid (10.90%) and n-hexadecanoic acid (5.36%) possess antimicrobial activity. The phytochemical 1,2,3-benzenetriol (pyrogallol) is an important allelochemical belongs to phenolic compound present in the plants and it is used in controlling bacterial diseases [28].

The phytochemicals Mequinol (0.74%), 4-methyl-phenol (6.86%) and 3-methyl-quinoline (0.44%) have anti-inflammatory activity. Mequinol is a common active ingredient in topical drugs used for skin depigmentation and it is a derivative of hydroquinone (4-hydroxyanisole, hydroquinone monomethyl ether). The phenolic compounds are one of the largest and most ubiquitous groups of plant metabolites. A number of studies have focused on the phytochemicals of biological properties such as antiapoptosis, anti-ageing, anticarcinogen, anti-inflammation, anti-atherosclerosis, cardiovascular protection and improvement of the endothelial functions, as well as inhibition of angiogenesis and cell proliferation [29]. Phenolic compounds have been extensively used in disinfections and as bacteriocides [30]. Quinoline compounds have excellent antibacterial activity against *Escherichia coli* and *Staphylococcus aureus* [31].

Based on the Dr. Duke's phytochemical and ethnobotanical databases, the 2-methoxy-4-methyl-phenol (9.55%), D-allose (10.63%), 6-methoxy-1H-indole-3-carbaldehyde (5.98%) and 2-methoxy-4-methyl-benzoic acid (10.90%) have antioxidant activity. The n-hexadecanoic acid (5.36%) and 2-methoxy-1-naphthyl thiocyanate (2.58%) have antitumor activity and 3-methyl-quinoline (0.44%) has antimalarial activity.

The antioxidant properties of phenolic compounds are due their redox properties, ability to chelate metals and quenching of singlet oxygen [32]. Natural phenolic compounds have great potential as natural antimicrobials and food preservatives [33]. D-allose is a novel antitumor monosaccharide, which causes cell growth inhibition on cancer cells by inducing the tumor suppressor gene [34]. The n-hexadecanoic acid has anti-tumor activity against human leukemic cells as well as murine cells [35]. The fatty acids exhibit cytotoxicity against HeLa cells and retard tumor growth [36]. The n-hexadecanoic acid and

octadecanoic acid were showed significant cytotoxicity against oral cancer (KB), breast cancer (MCF-7) and lung cancer (NCI-H187) [37].

The peak area of 1.54% was obtained by benzyl alcohol and it occurs naturally in many plants and it has strong activity against gram positive bacteria and has weak activity against gram negative bacteria, yeasts and moulds [38]. Benzyl alcohol has mild local anesthetic and anti-spasmodic properties [39].

In addition, the results of the GC-MS profile of latex of *C. bonplandianum* may be used as pharmacognostical tool for the identification of bioactive compounds. The presence of various bioactive compounds may be responsible for the therapeutic value of latex of *C. bonplandianum* to treat various ailments by traditional practitioners. However, further study is needed to isolate the individual phytochemical compound from the latex of *C. bonplandianum* and it may be helpful to find a novel drug.




Fig. 1. GC-MS chromatogram of ethanolic extract of latex of *C. bonplandianum*

Table 1. Screening of the phytocompounds of ethanolic extract of latex of *C. bonplandianum* by GC-MS

Sl. No.	RT	Name of the compound	Molecular Formula	MW	Peak %	Nature of compound	**Activity
1.	6.88	1-Methoxy-1,3,5-cycloheptatriene	C ₈ H ₁₀ O	122	1.52	Ether compound	No activity reported
2.	7.51	Benzyl alcohol	C ₇ H ₈ O	108	1.54	Alcoholic compound	Antibacterial
3.	8.62	Mequinol	C ₇ H ₈ O ₂	124	0.74	Phenolic compound	Anti-inflammatory
4.	8.87	4-Methyl-phenol	C ₇ H ₈ O	108	6.86	Phenolic compound	Anti-inflammatory
5.	10.48	2-C-Methyl-myoinositol	C ₇ H ₁₇ O ₆	194	32.4	Sugar derivative	Antimicrobial
6.	11.00	2-Methoxy-4-methyl-phenol	C ₈ H ₁₀ O ₂	138	9.55	Phenolic compound	Antioxidant Cytotoxicity
7.	12.98	4-Ethyl-2-methoxy-phenol	C ₉ H ₁₂ O ₂	152	0.77	Phenolic compound	No activity reported
8.	13.90	2-Methoxy-4-vinylphenol	C ₉ H ₁₀ O ₂	150	0.54	Phenolic compound	No activity reported
9.	14.72	4,4-Dimethyl-1,3-cyclopentanedione	C ₇ H ₁₀ O ₂	126	1.01	Ketone compound	Herbicidal activity
10.	15.33	3-Methyl-quinoline	C ₁₀ H ₉ N	143	0.44	Alkaloid compound	Antimalarial Antibacterial Antifungal Anti-inflammatory
11.	16.98	4-Amino-6-methyl-piperidin-2-one	C ₆ H ₁₂ N ₂ O	128	5.24	Alkaloid compound	Antimicrobial
12.	17.71	1,2,3-Benzenetriol	C ₆ H ₆ O ₃	126	3.54	Phenolic compound	Antimicrobial
13.	18.55	1H-Indol-5-ol	C ₈ H ₇ NO	133	0.35	Phenolic compound	No activity reported
14.	20.34	D-Allose	C ₆ H ₁₂ O ₆	180	10.63	Sugar compound	Antiproliferative, Antioxidant Endocytic

15.	22.61	3-Acetyl-2,5-dimethyl thiophene	C ₈ H ₁₀ OS	154	2.45	Thiophene derivative	No activity reported
16.	23.31	N-Ethyl-4-methoxy-1,3,5-Triazin-2-amine	C ₆ H ₁₀ N ₄ O	154	0.07	Amide compound	No activity reported
17.	26.10	6-Methoxy-1H-Indole-3-carbaldehyde	C ₁₀ H ₉ NO ₂	175	5.98	Indole aldehyde compound	Antioxidant
18.	26.91	2-Methoxy-4-methyl benzoic acid	C ₉ H ₁₀ O ₃	166	10.90	Benzoic acid derivative	Antimicrobial Antioxidant
19.	27.56	5-Methoxy-1H-Indole-3-carboxaldehyde	C ₁₀ H ₉ NO ₂	175	4.20	Indole aldehyde compound	Antioxidant
20.	31.09	n-Hexadecanoic acid (palmitic acid)	C ₁₆ H ₃₂ O ₂	256	5.36	Fatty acid	Antitumor Antibacterial Antifungal
21.	35.43	13-Octadecenal (stearaldehyde)	C ₁₈ H ₃₄ O	266	23.31	Fatty aldehyde compound	No activity reported
22.	35.84	Octadecanoic acid (stearic acid)	C ₁₈ H ₃₆ O ₂	284	1.95	Fatty acid	Antibacterial
23.	38.81	2-Methoxy-1-naphthyl thiocyanate	C ₁₂ H ₉ NOS	215	2.58	Naphthyl derivative	Antitumor Antifungal

4. CONCLUSION

In the present study, the GC-MS analysis of ethanolic extract of latex of *C. bonplandianum* revealed that the presence of twenty three phytochemicals. According to the Dr. Duke's phytochemical and ethnobotanical databases, eight compounds have antimicrobial, three compounds have antiinflammatory, five compounds have antioxidant, two compounds have antitumour and one compound has antimalarial activities. Thus, the GC-MS analysis is the first step towards understanding the nature of active principles in the latex of *C. bonplandianum* and this study may also be helpful to further research related to phytopharmacology.

REFERENCES

1. Vital PG, Rivera WL. Antimicrobial activity and cytotoxicity of *Chromolaena odorata* (L. f.) King and Robinson and *Uncaria perrottetii* (A. Rich) Merr extracts. J Medicinal Plants Res. 2009; 3(7):511-518.
2. World Health Organization (2008). Traditional medicine (Online). <http://www.int/medicines/areas/traditional/en/index.html>.
3. Krishnaiah D, Sarbatly R, Bono A. Phytochemical antioxidants for health and medicine – A move towards nature. Biotechnol Mol Biol Rev. 2007; 1(4): 097-104.
4. Sulekha Joshi, Gupta VP, Utkarsh Sharma. Phytochemical Screening of medicinal plant *Dolichandrone falcata*. Biological Forum – An International Journal. 2016; 8(1): 215-220.
5. Prasain JK, Wang CC, Barnes S. Mass spectrometric methods for the determination of flavonoids in biological samples. Free Radic Biol Med. 2004; 37(9):1324-1350.
6. De Rijke E, Out P, Neissen WMA, Ariese F, Gooijer C, Brinkman UA. The analytical separation and detection methods for flavonoids. J Chromatogr A. 2006; 1112(1-2):31-63.
7. Anjali Ruikar, Rasika Torane, Amruta Tambe, Vedavati Puranik, Nirmala Deshpande. GC-MS study of a steam volatile matter from *Mimusops elengi*. International Journal of Chem Tech Research. 2009; 1(2):158-161.
8. Kang H, Kang MY, Han KH. Identification of natural rubber and characterization of rubber biosynthetic activity in Fig tree. Plant Physiol. 2000; 123(3):1133-1142.
9. Upadhyay RK. Plant latex: A natural source of pharmaceuticals and pesticides. Int J Green Pharm. 2011; 5(3):169–180.
10. Agrawal AA, Konno K. Latex: A model for understanding mechanisms, ecology and evolution of plant defense against herbivory. Annu Rev Ecol Evol Syst. 2009; 40:311–315.
11. Duke JA. Medicinal plant science. London: Macmillian press limited. 1985.
12. Gubitz GM, Mittelbach M, Trabi M. Exploitation of the tropical oil seed plant *Jatropha curcas* L. Bioresour Technol. 1999; 67(1):73–82.
13. Konno K. Plant latex and other exudates as plant defense systems: Roles of various defense chemicals and proteins contained therein. Phytochem. 2011; 72(13):1510–1530.
14. Kitajima S, Kamei K, Taketani S, Yamaguchi M, Kawai F, Komatsu A, Inukai Y. Two chitinase-like proteins abundantly accumulated in latex of mulberry show insecticidal activity. BMC Biochem. 2010; 11:6–11.
15. Sethi A, Sharma RA. Micropropagation of *Croton bonplandinum* Baill. Int Res J Pharm. 2011; 2(10):82-86.
16. Reddy KR. Folk medicines from Chittoor District, Andhra Pradesh, used in the treatment of jaundice. Int J Crude Drug Res. 1988; 26:137-140.
17. Mishra D, Chaturvedi RV, Tripathi SC. The fungitoxic effect of the essential oil of the herb *Nardostachys jatamansi* D.C. Trop Agri. 1995; 72(1):48-52.
18. Verma SP, Kannaiya RS, Upadhyay RR. Antifungal activity of some Euphorbiaceaeous plants. Ind J Plant Path. 1998; 16(1-2):62-63.
19. Chandel KPS, Shukla G, Sharma N. Biodiversity in medicinal and aromatic plants in India, Conservation and utilization, New Delhi, National Bureau of Plant Genetic Resources. 1996: 361.
20. Das AJ, Dutta BK, Sharma GD. Medicinal plants use by different tribes of Cachar District, Assam. Indian J Tradit Know. 2008; 7(3):446–454.

21. Asthana A, Mall HV, Dixit K, Gupta S. Fungitoxic properties of latex of plants with special reference to that of *Croton bonplandianum* Baill. Int J Crude Drug Res. 1989; 27:25-28.
22. Kareem SO, Akpan I, Osho MB. *Calotropis procera* (Sodom apple) – A potential material for enzyme purification. J Biosci Technol. 2003; 87:133-135.
23. Varsha Jadhav, Vaibhav Kalase, Poonam Patil. GC-MS analysis of bioactive compounds in methanolic extract of *Holigarna grahamii* (Wight) Kurz. Int J Herb Med. 2014; 2(4):35-39.
24. Janakiraman N, Johnson M, Sahaya Sathish S. GC-MS analysis of bioactive constituents of *Peristrophe bicalyculata* (Retz.) Nees. (Acanthaceae). Asian Pac J Trop Biomed. 2012; S46-S49.
25. Cowan MM. Plants products as antimicrobial agents. Clin Microbial Rev. 1999; 12(4): 564- 582.
26. Vennila V, Udayakumar R. GC-MS Analysis of leaf, fruits and latex of *Croton bonplandianum* Baill. Int J Biochem Res Rev. 2015; 5(3):187-197.
27. Dr. Duke's Phytochemical and Ethnobotanical Databases, Agricultural Research Service of the US Department of Agriculture. <http://www.ars-grin.gov/duke> 16 June 2014.
28. Ismail Kocacaliskana , Ismet Talanb, Irfan Terzic. Antimicrobial activity of catechol and pyrogallol as allelochemicals. Naturforsch. 2006; 3(61c):639-642.
29. Han X, Shen T, Lou H. Dietary polyphenols and their biological significance. Int J Mol Sci. 2007; 8:950-988.
30. Okwu DE. Evaluation of the chemical composition of indigenous species and flavouring agents. Global J Pure Appl Sci. 2001; 8:455-459.
31. Pintilie Lucia. Quinolones: Synthesis and Antibacterial Activity. Antimicrobial Agents. 2012; 255-272.
32. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. Free Radic Biol Med.1996; 20:933-956.
33. Hayriye Cetin-Karaca. Evaluation of natural antimicrobial phenolic compounds against foodborne pathogens. University of Kentucky Master's Theses. Paper 652. http://uknowledge.uky.edu/gradschool_theses/652.
34. Yamaguchi F, Kamitori K, Sanada K, Horii M, Dong Y, Sui L, Tokuda M. Rare sugar D-allose enhances anti-tumor effect of 5-fluorouracil on the human hepatocellular carcinoma cell line HuH-7. J Biosci Bioeng. 2008; 106(3):248-52.
35. Harada H, Yamashita U, Kurihara H, Fukushi E, Kawabata J. Antitumor activity of palmitic acid found as a selective cytotoxic substance in a marine red alga. Anticancer Res. 2002; 22:2587-2590.
36. Abdel Aziz A, Rady HM, Amer MA, Kiwan HS. Effect of some honey bee extracts on the proliferation, proteolytic and gelatinolytic activities of the hepatocellular carcinoma Hepg2 cell line. Aust J Basic Appl Sci. 2009; 3:2754-2769.
37. Keawsa-Ard S, Liawruangrath B, Liawruangrath S, Teerawutgulrag A, Pyne SG. Chemical constituents and antioxidant and biological activities of the essential oil from leaves of *Solanum spirale*. Nat Prod Commun. 2012; 7:955–958.
38. Bezy alcohol. Committee for veterinary medical products. The European Agency for the Evaluation of Medicinal Products Veterinary Medicines Evaluation Unit. 1997.
39. Srinivasan GV, Sharanappa P, Leela NK, Sadashivam CT, Vijayan KK. Chemical composition and antimicrobial activity of the essential oil *Lea indica* Merr. flowers. Nat Prod Radiance. 2009; 8(5):488-493.