

Original Research Article

Evaluation of Phytochemical and *in-vitro* Antimicrobial Effects of Tomato (*Solanum lycopersicum* Linn.) on Oral Thrush and Human Cariogenic Pathogens

ABSTRACT

Aims: Study on the evaluation of phytochemical and antimicrobial effects of tomato *Solanum lycopersicum* (L) on oral thrush and human cariogenic pathogens was designed to determine the antimicrobial effect of tomato extracts, and to identify the common phytochemical constituents of tomato that may be inhibitory to oral pathogens.

Place and Duration of Study: Fresh tomato fruit samples were collected from Samaru market, Zaria in the month of November and December 2015, and analyzed at Microbiology Laboratory, Nigerian Institute of Leather and Science Technology Zaria, Kaduna.

Methodology: Cold maceration extraction using methanol and water was adopted. All the extracts were subjected to standard phytochemical qualitative screening for the presence or absence of various primary and secondary metabolites. Antimicrobial susceptibility was carried out by agar well diffusion technique. The antimicrobial susceptibility test of tomato (*Solanum lycopersicum*) fruit extracts against selected bacterial (*Streptococcus mutans*, *Bacillus subtilis*) and yeast (*Candida albicans*) pathogens capable of causing oral thrush and dental caries was carried out. Ciprofloxacin and Econazole were used as positive controls. The Minimum Inhibitory Concentrations (MICs) was determined in three concentrations; 100 mg/ml, 50 mg/ml and 25 mg/ml of each extract. Mean zone of inhibition was used to measure the antimicrobial potential of tomato fruit extracts against the test organisms. The microbial isolates were obtained from the Department of Medical Microbiology, Ahmadu Bello University, Zaria.

Results: Alkaloids, flavonoids, glycosides, saponins, tannins, steroids, phlobatannins, terpenoids and tannins are present in tomato fruit extract, but, anthraquinones and phlobatannins are absent in methanolic extract. While, the only phytochemical that was not detected in the aqueous extract is anthraquinones. Highest antibacterial activity was recorded on *Bacillus subtilis* at the concentration of 100mg/ml and 50mg/ml for aqueous and methanolic extracts respectively. *Candida albicans* and *Streptococcus mutans* showed resistance to the various extracts at various concentrations used. The MIC of 100mg/ml was recorded on *Bacillus subtilis*, whereas all other test organisms showed relative resistance to various concentrations of the extracts used.

Conclusion: Tomato fruit contains phytochemicals that showed promising antimicrobial effect on oral thrush and cariogenic pathogens. The tomato extract recorded antimicrobial effect against *Bacillus subtilis* with MIC of 100 mg/ml. The methanolic extract was the most active that could compete favourably with the conventional antibiotic (ciprofloxacin) at higher concentrations, which suggests that possible new drug candidates can be harnessed from the tomato fruit against oral thrush and cariogenic pathogens.

Keywords: Extraction; Lycopene; Minimum inhibitory concentration; Phytochemistry; *Solanum lycopersicum*

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68

1. INTRODUCTION

Tomato *Solanum lycopersicum* (Linn), previously known as *Lycopersicon esculentum* (Mill) is a red berry fruit [1]. It belongs to the plant family solanaceae and genus solanum [2]. Tomato is one of the most important vegetables worldwide because of its high consumption, year round availability and large content of health related components [3] used as juice, soup, ketchup or paste [4]. Although tomatoes are often closely associated with Italian cuisine, they are actually originally native to the Western side of South America in the region occupied by Columbia, Ecuador, Peru, Chile and the western half of Bolivia. Tomato is very rich in vitamins, minerals, essential amino acids, sugars, dietary fibers and lycopene, an antioxidant that reduce the danger related to many cancers and neurodegenerative diseases [5]. Today tomatoes are enjoyed worldwide to the tune of 130 million tons per year to the largest tomato producing country is China (with approximately 34 million tons of production) followed by United States, Turkey, India and China [6]. The plant requires relatively cool weather for optimum yield [7], it typically grows up to 1 to 3 meters (3 – 10 feet) in height and have a weak stem that often sprawls over the ground. It is a perennial in its native habitat, and grown as an annual in temperate climates. The leaves are 10 – 25 cm long, odd pinnate, with five to nine leaflets on petioles, each leaflet up to 8 cm long with a serrated margin [8].

Tomato contains variety of phytonutrients such as carotenoids (including beta-carotene, lutein, and zeaxanthin); flavonoids (including naringenin, chalconaringenin, rutin, kaempferol, and quercetin); hydroxycinnamic acids (including caffeic, ferulic, and coumaric acid); glycosides (including esculeoside A); and fatty acid derivatives (including 9-oxo-octadecadienoic acid). Lycopene, being the major carotenoid in tomato fruit, is the common antioxidant that reduces the danger related to many cancers and neurodegenerative diseases. It is a powerful antioxidant, anti-inflammatory and also has an antimicrobial property [5, 9, 10].

Some studies have shown that tomatoes have cancer prevention effect. The effect of tomato in the eradication of cancer is still unknown, but it was believed that lycopene and the newly discovered bioflavonoids in tomatoes are responsible cancer fighting agents [9]. Tomato is an excellent fruit vegetable for rapid skin cell replacement. Its juice can be used for healing sunburn because of its unique vitamin C. Tomato juice is a good sport drink used to restore one's body from fatigue and sleepiness [11].

The ever increasing demand for safer and cheaper herbal recipes in the developed countries has led to the extraction and development of several drugs and chemotherapeutic agents from plants as well as from traditional herbal remedies [12]. There are considerable economic benefits in the development of indigenous medicines and in the use of medicinal spices for the treatment of various diseases [13].

Majority of chemically synthesized drugs have serious adverse effect to the recipients, which may lead to temporary or permanent disability and incapacitations. Also, oral blisters, oral thrush, dental caries and oral ulcers are very serious infections that can lead to inability to take food or water as a result of dysphagia. The oral thrush disorder is a serious disease that can affect many people at various stages of their lives causing distress and discomfort. Sometimes, the disorder can lead to hospitalization as a result of starvation and dehydration following the victim's inability to ingest food or water [14, 15].

Direct topical application of raw tomato in combination with salt has been used for long in many African tribes to treat oral thrush, blisters, dental caries and ulcers of varying degrees. Therefore this research was designed to determine the antimicrobial effect of tomato, and to identify the common phytochemical constituents of tomato that may be inhibitory to oral pathogens. The research findings can further be used by the dental, pharmaceutical industries, complementary medicine and other medical sectors in tackling the menace of microbial oral disorders.

69

70 2. MATERIAL AND METHODS

71

72 2.1 SAMPLE COLLECTION

73 Fresh physically undamaged fruits of tomato (*Solanum lycopersicum*) were collected from
74 Samaru market Zaria based on the guidelines of convention on Biological Diversity, and
75 taxonomically characterized as Roma VFN, at the herbarium sections of Biological Science
76 Department, Ahmadu Bello University, Zaria and National Horticultural Research Institute,
77 Bagauda Station, Kano with the aid of botanical keys [16], ensuring that the tomato fruit is
78 not among the endangered plant species. The tomato fruits were collected fresh, healthy
79 and free from organic contaminants that may interfere with the substances of interest. The
80 fruits were thoroughly washed through running water, cut and dried under shade for 4 - 16
81 days, after which the dried fruits were pounded into a fine powder using a sterile mortar and
82 pestle [17]. The tomato fruit powder was stored and sealed in labeled sterile reagent bottles
83 for further use. The bioactive components were extracted using the method of Akerele *et al.*
84 [18].
85

86 2.2 PREPARATION OF PLANT EXTRACT

87 Plant extracts were prepared by the method of Alade and Irobi [19] with minor modifications.
88 To study the antibacterial potential of tomato (*Solanum lycopersicum*), non-polar solvent
89 such as methanol, and polar solvent such as water were used. A mass of 30 g of dried
90 powder was weighed on a weighing balance (Mettler 166®, USA) and was extracted by 100
91 ml of each solvent until the aqueous content evaporated completely as adopted by Lin *et al.*
92 [20]. At the end of extraction, the extracts were filtered through Whatman No. 1 filter paper
93 and concentrated over a water bath using a rotary-vacuum evaporator to recover the
94 solvents. The dried extracts were collected and weighed in varying concentrations and kept
95 in airtight container at 4 °C.

96

97 2.3 PRELIMINARY PHYTOCHEMICAL SCREENING

98 The preliminary phytochemical investigation was carried out for aqueous and methanolic
99 extracts of *Solanum lycopersicum* for the detection of various phyto-constituents by using
100 standard procedures to identify the constituents [20, 21].
101

102 2.3.1 Test for the presence of Alkaloids (Wagner's test)

103 Wagner's reagent was prepared by dissolving 2 g of iodine and 6 g of KI in 100 ml of water.
104 The plant extract was prepared by using 500 mg of plant extract, which was treated with few
105 drops of Wagner's reagent. A reddish brown coloured precipitate indicates the presence of
106 alkaloids.

107 2.3.2 Test for the presence of Anthraquinone (Borntrager's test)

108 About 0.5 g of the extract was boiled with 10% HCl for few minutes in a water bathe. It was
109 filtered and allowed to cool. Equal volume of chloroform was added to the filtrate. Few
110 drops of 10% ammonia was added to the mixture and heated. Formation of rose-pink colour
111 indicated the presence of anthraquinones.

112 2.3.3 Test for the presence of Flavonoids

113 The crude powder of dried plant was heated with 10 ml of ethyl acetate over a steam bath
114 for 3 minutes. The mixture was filtered and 4 ml of the filtrate was shaken with 1ml of dilute
115 ammonia solution, after which a yellow colouration was observed.

116 2.3.4 Test for the presence of Phlobatannins

117 An aqueous extract of each plant sample was boiled with 1% aqueous hydrochloric acid
118 (HCl) to observe the deposition of red precipitate.

119 2.3.5 Test for the presence of Glycosides (Fehling's test)

120 The crude plant powder of 0.5 g was dissolved in 5 ml of methanol. Exactly 2 ml of this
121 solution was dispensed into a test tube, and to it 10 ml of HCl was added. The mixture was
122 heated in a boiling water bath for 30 minutes, and to the mixture 5ml of Fehling's solution
123 was added and the mixture was boiled for another 5 minutes to observe a brick red
124 precipitate as an indication for the presence of glycosides.

125 2.3.6 Test for the presence of Saponins (Frothing test)

126 About 0.2 g of the extract was shaken with 5 ml of distilled water and then heated to boil.
127 Frothing (appearance of creamy layer of small bubbles) showed the presence of saponins.

128 2.3.7 Test for the presence of Steroids (Salkowski test)

129 A volume of 1 ml of plant extract was dispensed into a test tube, and to it few drops of
130 concentrated sulphuric acid was added. The presence of red colouration indicated the presence
131 of steroids.

132

133 **2.4 BACTERIAL PATHOGENS**

134 Characterized isolates of overnight cultures of oral thrush and dental caries causing test
135 organisms such as *Candida albicans*, *Streptococcus mutans* and *Bacillus subtilis* were
136 collected from the Dental Section, Medical Microbiology Laboratory, Ahmadu Bello
137 University, Teaching Hospital, Zaria. The isolates were kept in fresh broths to maintain their
138 viability.

139

140 **3.4 ANTIMICROBIAL SUSCEPTIBILITY SCREENING**

141 Screening of the antibacterial activity of the extracts was carried out using agar-well diffusion
142 method. Nutrient agar (N70123, Sigma-Aldrich Co., Germany) for bacterial isolates and
143 Sabouraud's Dextrose agar (RODAC™, Becton-Dickson and Co., USA) for *Candida*
144 *albicans* were prepared according to manufacturer's specifications, sterilized, and poured
145 into Petri dishes and allowed to set. Wells were made in the inoculated media using sterile
146 cork-borer (6 mm diameter, NC-11269, Science Company®, New Jersey, USA) after which a
147 little molten media was used to seal the base of the wells. The characterized and 0.5 scale
148 McFarland standardized test organisms were inoculated on the media by the spread plate
149 method using a sterile rod spreader to obtain uniform microbial growth. A volume of 0.1 ml
150 each of the prepared extracts (100 mg/ml, 50 mg/ml and 25 mg/ml) were transferred into the
151 corresponding wells with a sterile micropipette and were appropriately labelled, while 0.1 ml

of 20% DMSO (DMS, Gaylord Chemical Co., Louisiana, USA) (free of extract) was dispensed into a separate well to serve as the negative control. Ciprofloxacin (30 µg) was used as the positive control for bacterial isolates, while Econazole (30 µg) was used as positive control for *Candida albicans*. The plates were incubated for 48 hours at 37°C. Antibacterial and antifungal activities were evaluated by measuring the diameters of zones of growth inhibition in millimeters (Plate 1) after 48 hours incubation. Oxoid [22] standard susceptibility range was used to classify zones of inhibition as either sensitive (> 10 mm) or resistant (≤ 10 mm). Each experiment was conducted twice, and the mean of two results was taken for both the test and control as adopted by Irobi *et al.* [23].

2.5 DETERMINATION OF MINIMUM INHIBITORY CONCENTRATION

The Minimum Inhibitory Concentration (MIC) was determined as the least concentration that showed an inhibitory effect on test organism using the tube method. Two fold serial dilutions were made using nutrient broth (N70123, Sigma-Aldrich Co., Germany). The 5 ml of a solution of the extract (250 mg/ml) was added aseptically to 5 ml of double strength medium and mixed by shaking. Using a fresh pipette, 5 ml of the mixture was transferred to the second test tube which contained 5 ml of the single strength medium. This too was mixed by shaking and from it 5 ml will be taken into third test tube aseptically and mixed by shaking. The 9th tube containing no test compound served as control. Finally, to each tube 0.2 ml inoculums of the test organisms were added aseptically. The test tubes were covered with cotton wool and incubated at 37°C for 24 hours and then observed for turbidity. The lowest concentration that inhibited growth of test organism was noted as the MIC [24].

3. RESULTS AND DISCUSSION

Table 1 showed the phytochemical screening of tomato fruit extract. Phytochemicals such as alkaloids, flavonoids, phlobatannins, glycosides, saponins, tannins, steroids and terpenoids were detected in aqueous extracts analyzed. While, anthraquinone was not detected in either of the extracts analyzed. Whereas, all the phytochemicals tested were present in the methanolic extract except anthraquinones and phlobatannins.

Table 2 showed the antimicrobial effects of tomato aqueous extract against bacterial isolates. Highest antibacterial activity was recorded on *Bacillus subtilis* at the concentration of 100 mg/ml. All other test organisms showed resistance to the various extracts at various concentrations used.

Table 3 showed the antimicrobial effects of tomato methanol extract against bacterial isolates. Highest antibacterial activity was recorded on *Bacillus subtilis* at the concentration of 100 mg/ml and a slight sensitivity at 50 mg/ml. All other test organisms showed resistance to the various extracts at various concentrations used.

Table 4 showed the minimum inhibitory concentration of the tomato fruit crude extract. The MIC of 100 mg/ml was recorded on *Bacillus subtilis*, whereas all other tests organisms showed relatively resistance to the extract, and hence their MICs were not determined.

Table 1. Qualitative phytochemical screening of tomato fruit extracts

Phytochemicals	Aqueous extract	Methanol extract
Alkaloids	+	+
Flavonoids	+	+
Phlabotannins	+	–
Glycosides	+	+
Saponins	+	+
Tannins	+	+
Terpenoids	+	+
Anthraquinones	–	–
Steroids	+	+

*+ = Present; – = Absent

Based on the phytochemical analysis findings of the research study, it was found out that phytochemicals such as alkaloids, flavonoids, glycosides, saponins, tannins, steroids, phlabotannins and terpenoids and tannins were present in both aqueous and methanolic extracts of tomato fruit which have been associated with antimicrobial activities [25, 26]. Anthraquinone is absent in aqueous extract, while anthraquinone and phlabotannins are absent in methanolic extract. This might be attributed to the difference in the method of extraction employed, polarity of solvents used and also pointing to the fact that the use of solvents during extraction of medicinal plant might have some advantages to individual application. Sukhdev *et al.* [27] also reported the effect of temperature on phyto-components due to the fact that some of these components are thermo-labile in nature. The result reveals the presence of alkaloids which are a large group of secondary chemical constituents of plants made largely of ammonia compounds comprising basically of nitrogen bases synthesized from amino acid building blocks with various radicals replacing one or more of the hydrogen atoms in the peptide ring, mostly containing oxygen and they are intensely bitter in solutions.

The results for the antimicrobial screening have shown that at the concentration of 100mg/ml, all the extracts have antibacterial activity on *Bacillus subtilis*. Effectiveness factor of the tomato is due to the presence of Lycopene that possesses antibacterial and antifungal properties [28, 29]. The results of the inhibition of bacterial growth have shown that the extracts are active at high concentration and inactive at very low concentrations. Thus the study suggests that the inhibition of bacterial growth activity of the extracts is dose dependent. The aqueous and methanolic extracts had no inhibitory effect on the *Streptococcus mutans* and *Candida albicans* but showed inhibitory effects on the Gram-positive rod bacterium, *Bacillus subtilis*, with activity increasing with increase in concentration of extracts. This concurs with the findings of Omodamiro and Amechi [30] who reported antibacterial activity of tomato extract on some of pathogenic microbes. But, resistance of *Candida albicans* to both extracts is contrary to the findings of Sung *et al.* [31] and Pavlovic *et al.* [32] who reported that tomatoes proved to be effective against microorganisms such as *Staphylococcus aureus*, *Proteus*, *Bacillus* and antifungal to *Candida albicans* and *Aspergillus niger*.

234 **Table 2: Antibacterial Effect of Aqueous Tomato fruit Extract**

Test organism	Zones of growth inhibition						
	100 mg/ml	50 mg/ml	25 mg/ml	12.5 mg/ml	Ciprofloxacin	Econazole	DMSO
<i>Bacillus subtilis</i>	10 mm	0 mm	7 mm	0 mm	32 mm	–	0 mm
<i>Candida albicans</i>	0 mm	0 mm	0 mm	0 mm	–	36 mm	0 mm
<i>Streptococcus mutans</i>	0 mm	0 mm	0 mm	0 mm	35 mm	–	0 mm

235
236
237 **Table 3: Antibacterial Effect of methanol Tomato fruit Extract**

Test organism	Zones of growth inhibition						
	100 mg/ml	50 mg/ml	25 mg/ml	12.5 mg/ml	Ciprofloxacin	Econazole	DMSO
<i>Bacillus subtilis</i>	25 mm	15 mm	0 mm	0 mm	32 mm	–	0 mm
<i>Candida albicans</i>	0 mm	0 mm	0 mm	0 mm	–	35 mm	0 mm
<i>Streptococcus mutans</i>	0 mm	0 mm	0 mm	0 mm	36 mm	–	0 mm

238 **Table 4: Minimum Inhibitory Concentration (MIC) of Tomato fruit Extracts**

Test organisms	Concentration		
	Aqueous	Methanol	Control
<i>Bacillus subtilis</i>	100 mg/ml	50 mg/ml	None
<i>Candida albicans</i>	None	None	None
<i>Streptococcus mutans</i>	None	None	None

239

240 This may be due to the better solubility of the bioactive agents in the non-polar solvents. This
 241 conforms to the work of El-Mahmood *et al.* [33] who reported that the extractability of phyto-
 242 constituents differs from solvent to solvent depending on their polarities. It also reveals that
 243 the methanolic extract was more active against the Gram-positive bacteria. Since the
 244 traditional herbal remedy preparation use water as the extractant, it is a paradox that the
 245 aqueous extracts were less active in this study. It is possible that the aqueous extracts may
 246 contain antimicrobial constituents insufficient for efficacy in this study, which may explain
 247 why large amounts of the decoctions must be drunk by the patients. Jigna and Chanda [34]
 248 and Kitonde *et al.* [35] found out that, aqueous extracts showed little or no antimicrobial
 249 activity in contrast to those made using organic solvents. Yinegar *et al.* [36] reported that the
 250 success in traditional medicines may be due to administration of the extracts in large
 251 quantities and over a long period of time.

252 The Minimum Inhibitory Concentration (MIC) of the aqueous extract was highest for *Bacillus*
 253 *subtilis* at the concentration of 100 mg/ml with no activity in all other extract concentrations
 254 used. While, the MIC of the methanolic extract was least for *Bacillus subtilis* at the
 255 concentration of 50 mg/ml, suggesting that the methanol extract is more efficacious on
 256 *Bacillus subtilis* than the aqueous extracts. This is contrary to the studies of Omodamiro and
 257 Amechi [30] who reported MIC of 31.25 mg/ml for *Proteus mirabilis* and *Pseudomonas*
 258 *aeruginosa*. The antimicrobial effect of tomato in the treatment of oral microbial diseases can
 259 probably be enhanced when tomato is used in combination with table salt. The synergy
 260 between raw tomato and salt or honey might inhibit diverse types of oral pathogens.
 261 Previous studies by Al-Oqaili *et al.* [37] reported correlation between synergistic combination
 262 of tomato and honey with high antimicrobial therapeutic effects.

263

264 **4. CONCLUSION**

265

266 The secondary metabolites detected in the tomato fruit extracts include alkaloids, flavonoids,
 267 glycosides, saponins, tannins, steroids, phlabotannins, terpenoids and tannins.
 268 Anthraquinone and phlabotannins are absent in methanolic extract. While, the only
 269 phytochemical that was not detected in the aqueous extract is anthraquinones.

270 *Bacillus subtilis* was susceptible to all the extracts at the concentrations of 100mg/ml, but
 271 *Streptococcus mutans* and *Candida albicans* were not susceptible to the aqueous and
 272 methanolic extracts within the concentration range for this study. The methanolic extract was
 273 the most active that could compete favourably with the conventional antibiotic (ciprofloxacin)
 274 at higher concentrations. The activity of the extracts on the test organisms increased with
 275 increase in concentration. The methanol extract exhibited bacteriostatic effects on the
 276 *Bacillus subtilis* at lower concentrations and bactericidal effects at higher concentrations,
 277 with minimum inhibitory concentration of 100mg/ml.

This research may serve as a scientific basis and lend credence to the claim by the traditional medicine practitioners that the tomato fruit is a useful herbal remedy to various microbial oral infections.

ETHICAL APPROVAL

Not applicable

REFERENCES

1. IAPT (2016). International Code of Nomenclature for algae, fungi and plants. International Association for Plant Taxonomy. Chapter V, section 3. Retrieved on 14th September 2016. Available at: www.iapt-taxon.org
2. Sarah, C. D., Sandra K & Iris E. P. (2003). Taxonomy of tomatoes in the Gal'apagos Islands: native and introduced species of *Solanum* section *Lycopersicon* (Solanaceae). *Journal of systematics and biodiversity*, 1 (1): 29–53.
3. Krishna, M. J., Bhaumik, A. and Kumar, S.P. (2013). Phytochemical Analysis and Antimicrobial Studies of Various Extracts of Tomato (*Solanum lycopersicum* L.). *Scholars Academic Journal of Biosciences (SAJB)*, 1(2):34-38.
4. Dilis B and Trichopoulou A. (2010). Antioxidant intakes and Food Sources in Greek Adults. *The Journal of Nutrition*. Bethesda, 140(7); 1274-1279.
5. Srinivasan, R. (Ed.). (2010). *Safer Tomato Production Methods: A field Guide for soil fertility and pest management*. AVRDC- The World Vegetable Center, Shanhua, Taiwan. AVRDC Publication No. 10-740. pp. 2.
6. Parnell, T., Suslow Trevor V., Lindaj, H. (2014). *Tomato Safe Method to store preserved and enjoy* (PDF) and catalogue University, California Division of Agriculture and Natural Resources Retrieved 18th February, 2015.
7. Rodriguez, G.R. (2007). Effect of Rice Bran Mulching on Growth and Yield of Cherry Tomato. *Cienciae Investigacion Agraria*, 34: 181-186.
8. Acquaah, G (2002). *Horticulture: Principles and Practices*. New Jersey. Prentice Hall. ISBN: 0130331252.
9. Etminan, M., Takkouche, B. and Caamaño-Isoma, F. (2004). The Role of Tomato Products and Lycopene in the Prevention of Prostate cancer: A meta-analysis of Observational Studies. *Cancer Epidemiol. Biomarkers Prev.*, 13 (3): 340-345.
10. Silaste ML, Alfthan G, Aro A, Kesaniemi YA, and Horkko S. (2007). Tomato Juice Decreases LDL Cholesterol Levels and LDL Resistance to Oxidation. *Br J. Nutr.*; 98 (6):1251-8. PMID: 17617941.
11. Makinnon and Rio L. G. (2011). Dietary Restriction of Lycopene for a period of one month resulted in significantly increased biomarkers of Oxidative Stress and bone respiration in postman woman. *J. Nutrition Health Ageing*, 15(2): 133 - 8.
12. Falodun, A., Okunrobo, L. O. and Uzoamaka, N. (2006). Phytochemical screening and anti-inflammatory evaluation of methanolic and aqueous extracts of *Euphobia heterophylla* Linn (Euphobiaceae). *African Journal of Biotechnology*, 5(6): 529-531.
13. Azaizeh, H., Fulder, S., Khalil, K. and Said, O. (2003). Ethnomedicinal knowledge of local Arab practitioners in the Middle East Region. *Fitoterapia*, 74: 98-108.
14. Bouquot Brad W. Neville, Douglass D. Dammi Carl M. Allen Jerry E. (2002). *Oral and Maxillofacial Pathology* (2nd edition) Philadelphia: WB - Sauders pp. 189 - 197 ISBN 072160033.
15. Talib N. (2016). Disorders of oral pigmentation: Drugs and Diseases in Dentistry. *Medscape*. Accessed on August 02 2015. Available at www.emedicine.medscape.com/article/1078143-references?src=refgatesrc1/
16. Arber (1972). *Water Plants: A Study of Aquatic Angiosperm*. Welden Wisely Limited Pp. 436.

- 332 17. Onoruvwe O. and Olorunfemi P. O. (1998). Antimicrobial Screening and
333 Pharmacognostical Evaluation of *Dischrostachys cinerea* Root; West Africa Journal
334 of Biol. Science; 7: 91-99.
- 335 18. Akerele, Z. O., Obasuyi, O., Ebomoyi, M.I., Oboh, I.E., and Uwumarongie. O.H.
336 (2008). Antimicrobial activity of the ethanol extract and fractions of the seeds of
337 *Garcinia kola* Heckel (Guttiferae). African Journal of Biotechnology, 7(2): 169-172.
- 338 19. Alade, P.I. & O.N. Irobi, (1993). Antimicrobial Activities of Crude Leaf Extracts of
339 *Acalypha wilkesiana*. J. Ethnopharmacol. 39, 171-174.
- 340 20. Lin J, Opoku AR, Geheeb-Keller M, Hutchings AD, Terblanche SE, Jager AK,
341 Staden (2015). Available at: www.globalresearchonline.net Also on: www.education-science.com/10/comparative-study-on-phytochemical.html?m=1
342
- 343 21. Tiwari, P., Kumar, B., Kaur M, Kaur G & Kaur, H. (2011). Phyto-chemical Screening
344 and Extraction: A Review. Internationale Pharmaceutica Scientia. 1 (1): 98-106.
- 345 22. Oxoid (1985). Oxoid manual of dehydrated culture media, ingredients and other
346 laboratory services. Basingstoke, England.
- 347 23. Irobi, O. N., Moo-Young, M., Anderson, W. A. and Daramola, S. O. (1994).
348 Antimicrobial activity of bark extracts of *Bridelia ferruginea* (Euphorbiaceae).
349 International Journal of Pharmacognosy, 34(2): 87-90.
- 350 24. Cheruiyot, K. R., Olila, D. and Katerega, J. (2009). In vitro antibacterial activity of
351 selected medicinal plants from Longisa region of Bomet district, Kenya. African
352 Health Sciences, 9(S1): S42-S46.
- 353 25. Reynolds, T. and Dweck, A. C. (1999). Aloe vera Leaf Gel: A Revise Update.
354 Journal of Ethnopharmacology, 68: 3-37.
- 355 26. Aliyu, A. B., Musa, A. M., Abdullahi, M. S., Oyewale, A. O. and Gwarzo, U. S.
356 (2008). Activity of plant extracts used in Northern Nigerian traditional medicine
357 against methicillin-resistant *Staphylococcus aureus* (MRSA). Nigerian Journal of
358 Pharmaceutical Sciences, 7(1): 1-8.
- 359 27. Sukhdev S.H., Suman, P. S. K., Gennaro, L and Dev, D. R. (2008). Extraction
360 Technologies for Medicinal and Aromatic Plants International. Centre for Science
361 and High Technology, Trieste ed, pp 11.
- 362 28. Dahan K, Fennal M, Kumar NB. (2008). Lycopene in the prevention of prostate
363 cancer. J. Soc Integr Oncol., 6:29-36.
- 364 29. Rao AV. (2002). Lycopene, tomatoes, and the prevention of coronary heart disease.
365 Exp Biol Med (Maywood); 227:908-913.
- 366 30. Omodamiro O. D. and Amechi U. (2013). The phytochemical content, antioxidant,
367 antimicrobial and anti-inflammatory activities of *Lycopersicon esculentum* (Tomato).
368 Asian Journal of Plant Science and Research, 2013, 3(5):70-81.
- 369 31. Sung W.S., Lee I.S. and Lee D.G. (2007). Damage to the cytoplasmic membrane
370 and cell death caused by lycopene in *Candida albicans*. J Microbiol Biotechnol.,
371 17(1):797-804.
- 372 32. Pavlović R., Mladenović J., Radovanović B., et al. (2013). In Vitro Antimicrobial
373 Activity of Ethanol Tomato Extracts. Conference VIVUS, 24th-25th April 2013.
374 Biotechnical centre Naklo, Strahinj 99, Naklo Slovenia, Biotechnical Centre Naklo
375 Higher Vocational College, Strahinj 99, Naklo, Slovenia.
- 376 33. El-Mahmood, A. M. and Ameh, J. M. (2007). In vitro antibacterial activity of *Parkia*
377 *biglobosa* (Jacq) root bark extract against some microorganisms associated with
378 urinary tract infections. African Journal of Biotechnology, 6(11): 1272-1275.
- 379 34. Jigna, P. and Chanda, S. V. (2007). In vitro antimicrobial activity and phytochemical
380 analysis of some Indian medicinal plants. Turkish Journal of Biology, 3: 53-58
- 381 35. Kitonde, C. H., Fidahusein, D. S., Lukhoba, C. W. and Jumba, M. M. (2014).
382 Antimicrobial activity and phytochemical screening of *Senna didymobotry* used to
383 treat bacterial and fungal infections in Kenya. International Journal of Education and
384 Research, 2(1): 1-12.
- 385 36. Yineger, H., Kelbessa, E., Bekele, T. and Lulekal, E. (2008). Plants used in
386 traditional management of human ailments at Bale Mountains National Park, South-
387 eastern Ethiopia. Journal of medicinal plant research, 2(6): 132-153.

388 37. Al-Oqaili RMS., Basim B, Istabreq M., Salman MA., Abed D. and Asaad A. (2014). In
389 Vitro Antibacterial Activity of *Solanum lycopersicum* Extract against some
390 Pathogenic Bacteria. Food Science and Quality Management, 27: 12-17.

391

392 ACRONYMS

393

394 **DMSO**: Dimethyl sulfoxide

395 **MIC**: Minimum Inhibitory Concentration

396 **cfu/ml**: Colony forming units per milliliter

397

398 APPENDIX

399



400

401

Plate 1: Antimicrobial susceptibility setup of tomato extract