

Original Research Article

Compositions of *Thunbergia grandiflora* leaf and root essential oils

ABSTRACT

FRESH SAMPLES OF LEAVES AND ROOTS OF *THUNBERGIA GRANDIFLORA* WERE COLLECTED FROM FOREST- RESERVED AREA, IBADAN, NIGERIA FOR EXTRACTION OF THEIR VOLATILE OILS. HYDRO-DISTILLATION METHOD WAS USED IN AN ALL-GLASS CLEVANGER APPARATUS DESIGNED TO BRITISH PHARMACOPEIA SPECIFICATIONS. IT GAVE YIELDS OF 0.09 % AND 0.10 % RESPECTIVELY. THE OILS WERE ANALYZED USING GAS CHROMATOGRAPHY [GC], GAS CHROMATOGRAPHY-MASS SPECTROMETRY [GC-MS] IN UNIVERSITY OF BOTSWANA. A TOTAL OF THIRTY-EIGHT COMPOUNDS WERE IDENTIFIED IN THE LEAF AND ROOT ESSENTIAL OILS, TWENTY-SEVEN COMPOUNDS WERE IDENTIFIED IN LEAF OIL, WHICH ACCOUNT FOR 98.13% OF IT WHILE SEVENTEEN IDENTIFIED COMPOUNDS ARE RESPONSIBLE FOR 73.98% OF ROOT OIL. MOST ABUNDANT COMPOUNDS IN LEAF OIL ARE A-3-OCTANOL (45.96%), 3,7 DIMETHYL 1,6-OCTADIEN-3-OL (13.22%) AND 2-METHOXY-3-(2-PROPENYL) PHENOL (8.36%). DOMINANT COMPOUNDS IN ROOT OIL ARE METHYL SALICYLATE (44.80%), 2,2¹,5,5¹-TETRAMETHYL-1,1¹ - BIPHENYL (5.05%) AND I-PROPYL HEXADECANOATE (2.87%). LEAF OIL IS DOMINATED BY FOLLOWING CLASSES OF COMPOUNDS (%); ALCOHOLS (68.58), CARBOXYLIC ACID (4.01), ALDEHYDES (12.90), KETONES (5.18), ESTERS (2.62), AROMATICS (0.66) AND ALSO SULPHURATES (0.45). CLASSES OF COMPOUNDS IN ROOT OIL ARE ESTERS (47.67), ALCOHOLS (2.64), CARBOXYLIC ACIDS (0.55), ALDEHYDES (7.69), SULPHURATES (1.61), AROMATICS (5.05) AND ALKANE (5.02). THE TWO OILS CONTAIN ALCOHOLS AND ESTERS IN GOOD AMOUNTS. OUR RESULTS INDICATE THAT ALCOHOLS, ESTERS AND ALDEHYDES DOMINATE *THUNBERGIA GRANDIFLORA* ESSENTIAL OILS. WE PRESENT IMPORTANT COMPOUNDS IN THE LEAF AND ROOT ESSENTIAL OILS OF *THUNBERGIA GRANDIFLORA*, WHICH HAVE NOT BEEN REPORTED EARLIER IN LITERATURE.

KEYWORDS: *Thunbergia grandiflora*, *Acanthaceae*, essential oil, terpenoids, Gas Chromatography(GC) and Gas Chromatography – Mass Spectrometry(GC/MS)

33

34 1. INTRODUCTION

35 *Thunbergia grandiflora* Roxb.exRottler (*Acanthaceae*) is an evergreen vine. It is a large climbing and
 36 twining shrub, native to China, India, Nepal, Indochina and Burma and many tropical countries of Africa
 37 [1, 2, 3, 4]. It is also found throughout the Bangladesh, especially in forests of Gajipur and Chittagong
 38 [5]. Generally; it is known as black clock vine and blue trumpet vine. Other common names include Bengal
 39 clockvine, Bengal trumpet, blue skyflower, blue thunbergia, clock vine, sky flower and sky vine[3]. In
 40 Bangladesh, it is known as “Kauathuti”, “Nallata” and “Nillata” [4]. This Plant grows to about 20 metres in
 41 height and has a tuberous root system with a deep tap root [2]. The stalked, opposite leaves, which have
 42 rough surface, are quite variable in shape, they may be triangular or ovate and the margins may be
 43 toothed, lobed or entire. Length of the leaf is up to 20 cm and width is up to 6 cm broad, broad-based
 44 narrowing to a pointed tip, usually with deeply scalloped lobes towards the base. The shape and texture
 45 of the leaf as well as being oval and narrowing to pointed tip makes the plant different from other
 46 *thunbergia* species. The trumpet-shaped flowers have a short broad tube, blue to mauve on the outside,
 47 pale yellow inside, which expands to five rounded, pale lavender-blue petals, one larger than the others.
 48 The seed is flat, up to 1 cm long and covered with brown scales, the pods containing the seeds are
 49 ejected several metres upon ripening[2]. The plant is cultivated as houseplant in temperate regions and
 50 has gained the Royal Horticultural Society's Award of Garden Merit [6, 7].

51 *Thunbergia grandiflora* has very rampant growth habit and is great for the quick covering of arbors,
 52 trellises and fences. It is used as container plant, marvelous cascading over wall. The plant climbs and
 53 blankets native vegetation, with the weight of the vine often pulling down mature trees. *T. grandiflora* is
 54 the most widespread pest species, having been used as a garden ornamental for its attractive large
 55 leaves and hanging groups of large, pale lavender flowers [8]. Other species of *Thunbergia* includes *T.*
 56 *laurifolia*, *T. laevis nees*, *T. alata*, *T. erecta*, *T. coccinea*. Traditionally, some of these species have been
 57 used for treating fever, wounds, skin problems, back and joint pains, eye inflammation, piles and rectal
 58 cancer. They have also been used as aphrodisiac and antidote for snake bite [13-15]. *Thunbergia*
 59 *laurifolia* was found to possess antioxidant, hepatoprotective, neuroprotective and hypoglycaemic
 60 properties[16-19]. *Thunbergia laurifolia* have been reported to contain compounds such as grandifloric
 61 acid, 8-epigrandifloric acid, 3'-O- β -glucopyranosyl-stilbericoside, benzyl β -glucopyranoside, (E)-2-
 62 hexenyl- β -glucopyranoside, hexanol- β -glucopyranoside and 6,8-di-C glucopyranosylapigenin[20]. *T.*
 63 *laevisnees* was reported to display significant antioxidant and anti-diabetic properties [21, 22]. Iridoid
 64 glucosides, apigenin glucosides, stilbericoside and thunbergioside were also obtained from *T. laevis nees*
 65 [23-25] as well as essential oil constituents, phytochemical analyses, and antimicrobial studies on leaf
 66 and stem of *thunbergia laevis nees* [26]. *Thunbergia grandiflora* is used for the treatment of blood
 67 dysentery, cataract, conjunctivitis, diabetes, gout, hydrocele, hysteria, malaria, marasmus, ophthalmia,
 68 postpartum pre-eclampsia, rheumatism, spermatorrhoea, stomach complaints, elephantiasis and urinary

bladder stone[9, 10, 11]. They are used as traditional medicines for cardiac diseases and blood purification. Leaves and stem of *T. grandiflora* are used as a poultice in stomach complaints and to treat eye diseases in Chittagong Hill Tracts and it has anthelmintic activity[12].The methanolic extracts of *T. grandiflora* flowers showed presence of alkaloids and phenols [27, 29]. The novel iridoid glycosides; isounedoside and grandifloric acid were isolated from *thunbergia grandiflora* [28]. Research has not been done on the essential oil composition of *T. grandiflora*; we therefore report the leaf and root essential oil compositions of *thunbergia grandiflora*, which is scarce in literature.

2. EXPERIMENTAL DETAILS

2.1. Plant material

Fresh samples of leaves and roots of *Thunbergia grandiflora* were collected from forest reserved area, Ibadan, Nigeria for extraction of their volatile oils on 8th June, 2015. Plant was authenticated at the Herbarium, Department of Botany, University of Ibadan, where voucher samples have been deposited, with voucher number UIH-22440

2.2. Isolation of essential oils

Sample of *Thunbergia grandiflora* was collected in large quantities, which was separated into leaf and root parts giving 235 g of leaf and 865 g of root. They were crushed separately and hydro distilled for 2.5 to 3 hours in an all glass Clevenger-type apparatus designed to British Pharmacopeia specifications. Oils were collected under iced condition with very small quantity of distilled n-hexane, which the analyzing GC corrected. Leaf and root essential oils were procured in 0.09 % and 0.1 % yields respectively.

2.3. Gas Chromatography

Leaf and root essential oils were analyzed using an Agilent 5975C series Gas Chromatograph - Mass Spectrometer (GC-MS) system. Data analysis was done using Productivity Chemstation Version E.02.01.1177 in conjunction with the Automated Mass Spectral Deconvolution and Identification System (AMDIS). The oven program for the column was as follows: The equilibration time was set at 0.25 min; the initial temperature was set at 70 °C, held there for 4 minutes, and then ramped at 6 °C/min to 140 °C for 6 min. It was further ramped at 15 °C/min to 300 °C for 4 minutes. The total run time was 33.60 minutes. 1 µL of sample extract was injected in the splitless mode. The injector temperature was set at 280°C and the column head pressure was 56.756 kPa yielding a total flow of 64mL/min. The transfer line temperature was 280 °C. A HP-5MS 5% Phenyl Methyl Silox column with dimensions of 30 m x 250 µm x 0.25 µm and maximum recommended temperature (MRT) of 325 °C was used throughout. Helium was used as a carrier gas at a flow rate of 1 mL/min.

2.4. Mass Spectrometry

The acquisition mode was full scan with a solvent delay of 4.00 minutes. Acquisition mass range was set at *m/z* 50– 600 and the electron multiplier detector voltage was auto set to 1082 V. The mass

spectrometer source and quadrupole mass analyzer temperatures were set to 230 °C and 150 °C respectively. The filament emission current was auto set to 34.610 µA.

2.5. Identification of components

Identification of the essential oil components were based on their retention indices (determined with reference to a homologous series of n-alkanes), and by comparison of their mass spectral fragmentation patterns with the National Institute for Standards and Technology (NIST) library version 8.0. The AMDIS software version 2.71 build 134.27 was used for deconvolution i.e. to compare the NIST library spectra with component spectra. The minimum match factor was for deconvolution on AMDIS was set to 70 and all identified components had match factors between 70 and 100. Some mass spectral fragmentation patterns with characteristic retention were compared with standard inbuilt data of Adams (1995) [30].

3. RESULTS AND DISCUSSION

Volatile oils from leaf and root parts of *Thunbergia grandiflora* Roxb.exRottler, an endangered medicinal *Acanthaceae*, were obtained in Nigeria by hydro-distillation using the all-glass Clevenger apparatus designed to British Pharmacopeia specifications and gave yields of 0.09 % and 0.1 % respectively [Table 1]. The oils were analyzed using gas chromatography [GC], gas chromatography-mass spectrometry [GC-MS] in University of Botswana [Figures 1 and 2], which afforded the identification of 27 compounds in leaf and 17 compounds in root. Results of the 38 identified compounds are presented in Table 2. While Table 3 has result of the comparison of the ten classes of compounds found in the two *T. grandiflora* essential oils studied.

Table 1. Essential oils procured from leaf and root parts of *Thunbergia grandiflora*

Plant part	Weight of sample (g)	Weight of essential oil procured (g)	%Yield of essential oil procured	Physical examination
leaf	235	0.2	0.09	Leafy odour
root	865	0.9	0.10	Woody smell




Figure 1: Gas chromatogram of the leaf essential oil of *Thunbergia Grandiflora* [see GC conditions under experimental procedures]

Table 2: Chemical composition of volatile oils of leaf and root thunbergia grandiflora

S/N	RI	Compound	% composition in	
			Leaf	Root
1	1055	3 - octanol	45.96	-
2	1071	3,7 dimethyl 1,6 – octadien - 3 - ol	13.22	-
3	1762	Methyl salicylate	-	44.80
4	2152	2 – methoxy - 3- (2 – propenyl) phenol	8.36	-

5	2226	1 – (2,6,6 – trimethyl – 1,3,cyclohexa – 2 – buten – 1 - one	2.61	-
6	2315	2,3 dihydro -1,1-dimethyl 1H indene – 4 – carboxylic acid	0.52	-
7	2361	4 – (2,2- dimethyl – 6- methylene cyclo – 3 – buten – 2 -one	0.67	-
8	2403	α phenylethyl butanoate	0.69	-
9	2526	Nonylcyclopropane	-	2.50
10	2596	4 – (2,6,6, - trimethyl – 1- cyclohexen – 3- buten – 2 - one	1.55	-
11	3091	2 – formyl methyl – 4,6,6 – bicyclo [3.1.1] hept – 3 - ene	0.86	-
12	3343	2,2 ¹ , 5, 5 ¹ , tetramethyl 1,1 ¹ - biphenyl	0.66	5.05
13	3348	tetradecenal	3.82	-
14	3446	n - pentadecanol	1.04	-
15	3454	9, 10 – dihydodiethyl – anthracene – 9,10 – biimine – 11,12 - dicarboxylate	0.66	-
16	3501	hexadecanal	3.58	-
17	3535	6,10,14, - trimethyl – 2- pentadecanone	0.35	0.92
18	3577	1 - hexadecanol	-	2.64
19	3580	1,3,3 – trimethylbicyclo [2.2.1] heptane	0.43	-
20	3589	(Z) 9,17 octadecadienal	0.96	2.84
21	3596	(Z) 7 tetradecenal	-	2.79
22	3597	Cis – 9 - hexadecenal	2.51	-
23	3621	pentadecanal	1.18	2.06
24	3671	n – hexadecanoic acid	3.49	-

25	3697	Ethyl hexadecanoate	0.56	-
26	3702	(Z) – 13- octadecenal	0.85	-
27	3726	i – propyl hexadecanoate	-	2.87
28	3765	2 – (4 ¹ – methoxyphenyl) – 2- (3 ¹ – methyl – 4 ¹ methoxyphenyl) propane	-	0.63
29	3780	(E) 3 - Eicosene	-	1.68
30	3809	phytol	2.00	1.15
31	3849	2 – chloroethylinoleate	0.31	-
32	3855	Methyl 11,14,17, Eicosatrienoate	0.40	-
33	3953	2,7 dimethyl octane	-	0.82
34	3979	Benzo[b] thiophene – 3 – carboxylic acid	-	0.55
35	4026	3,3 – dimethyl hexane	0.44	-
36	4095	hexadecane	-	1.07
37	4161	Decyl -2- ethyl hexyl sulphurate	-	0.63
38	4224	2 ethyl hexyl isohexylsulphurate	0.45	0.98
% identified			98.13	73.98

135 RI means retention index

136




Figure 2: Gas chromatogram of the root essential oil of *Thunbergia Grandiflora* [see GC conditions under experimental procedures]

Table 3. Classes of compounds in *T.grandiflora* leaf and root essential oils.

S/N	Class of compound	% in each of Thunbergia grandiflora essential oils	
		Leaf	Root
1	alkene	0.86	1.68
2	aldehyde	12.90	7.69
3	ketone	5.18	0.92
4	alkane	0.87	5.02

5	aromatics	0.66	5.05
6	esters	2.62	47.67
7	alcohols	68.58	2.64
8	carboxylic acid	4.01	0.55
9	sulphurate	0.45	1.61
10	terpenoid	2.00	1.15

3.1. DISCUSSION

Volatile oils from leaf and root of *Thunbergia grandiflora* were obtained by hydro-distillation. Oils were procured in 0.09% and 0.10% yields respectively, each having characteristic distinctive notes: Leaf oil (0.09% yield) had a leafy odour while root oil (0.10%) had woody smell (Table 1). Compounds identified in each are listed in Table 2. Thirty-eight volatile compounds were identified in the two essential oils. The leaf oil contain 27 compounds, which make-up 98.13 % of it, while root oil have 17 compounds accounting for 73.98 % of it; Dominant compounds (%) in leaf essential oil are: α -3-octanol (45.96), 3,7 dimethyl 1,6-octadien-3-ol (13.22) and 2-methoxy-3-(2-propenyl) phenol (8.36). While dominant compounds (%) in root essential oil are: methyl salicylate (44.80), 2,2¹,5,5¹-tetramethyl-1,1¹ - biphenyl (5.05) and i-propyl hexadecanoate (2.87). Leaf oil is dominated by following classes of compounds (%): alcohols (68.58), carboxylic acid (4.01), aldehydes (12.90), ketones (5.18), esters (2.62), aromatics (0.66) and sulphurate (0.45). Classes of compounds (%) in root oil are esters (47.67), alcohols (2.64), carboxylic acids (0.55), aldehydes (7.69), sulphurate (1.61), aromatics (5.05) and alkane (5.02). Leaf and root oils contain different amounts of ten classes of compounds shown in **table 3**. The root essential oil is a good source of esters while the leaf essential oil is a good source of alcohols. Compounds present in both leaf and root essential oils include (%); 2,2¹, 5, 5¹, tetramethyl 1,1¹ - biphenyl (0.66 in leaf and 5.05 in root), 6,10,14, - trimethyl – 2- pentadecanone (0.35 in leaf and 0.92 in root), (Z) 9,17 octadecadienal (0.96 in leaf and 2.84 in root), pentadecanal (1.18 in leaf and 2.06 in root) and phytol (2.00 in leaf and 1.15 in root)(**table 2**). The presence in significant amount of oxygenated compounds (esters, alcohols, acids, terpenoids) in these oils is an important quality determining factor, they are known to have favouring and pharmacological properties as well as being important precursors in acetate and shikimate biosynthesis of important metabolites [34]. Phytol is an important component of all plants used in cosmetics, shampoos, toilet soaps, and household cleaners as it shows antimicrobial, anticancer, antidiuretic, antioxidant, anti-inflammatory, antitumor, antimicrobial, and chemopreventive properties and used in vaccine formulations [31, 32]. Methyl salicylate is used in high concentrations as analgesic to treat joint and muscular pain [33], 3-Octanol is found in alcoholic beverages, it is a flavouring agent. n-

hexadecanoic acid have been reported to possess anti-inflammatory, antioxidant, hypocholesterolemic and antibacterial activities, medicated oils rich in n-hexadecanoic acid is used for the treatment of rheumatic symptoms in the traditional medical system. [35, 36]. The identified 38 compounds play important role in the vast ethno-medicinal uses and biological activities demonstrated by *T. grandiflora* [9, 27, 28, 29].

4. CONCLUSION

Our results indicate that *Thunbergia grandiflora* essential oils are dominated by alcohols, esters and aldehydes. This accounts for its important use in ethno-medicinal activities. This study revealed important compounds in the leaf and root essential oils of *Thunbergia grandiflora*, which have not been reported earlier in literature.

References

- [1] "*Thunbergia grandiflora*". Australian Plant Name Index (APNI), IBIS database. Centre for Plant Biodiversity Research, Australian Government, Canberra. 2013.
- [2] "*Thunbergia grandiflora*". Weed Identification. Australian Weeds Committee 2013.
- [3] "Taxon: *Thunbergia grandiflora* Roxb.". Germplasm Resources Information Network (GRIN). United States Department of Agriculture, Agricultural Research Service, Beltsville Area. 2013.
- [4] Motaleb, MA, Hossain, MK, Alam, MK, Abdullah-Al Mamun, MM, Sultana, M. commonly used medicinal herbs and shrubs by traditional herbal practitioners. Glimpses from Thanchiupazila of Bandarban. Dhaka: International Union for Conservation of Nature and Natural Resources; 2013
- [5] Medicinal Plants Database of Bangladesh. *Thunbergia grandiflora* Roxb. Chittagong: Medicinal Plants Database of Bangladesh; 2014. [Accessed on 22nd March, 2015] Available from: <http://www.mpbdb.info/plants/thunbergiagrandiflora.php>
- [6] "RHS Plant Selector – *Thunbergia grandiflora*". Retrieved 27 June 2013.
- [7] RHS A-Z encyclopedia of garden plants. United Kingdom: Dorling Kindersley. 2008 p1136.
- [8] Gilman EF. *Thunbergia grandiflora*, institute of food and agricultural sciences, Florida Cooperative Extension Service university of florida, 1999, 1 - 3,
- [9] Uddin, SN. Traditional uses of ethnomedicinal plants of the Chittagong Hill Tracts. Dhaka: Bangladesh National Herbarium; 2006, p994
- [10] Begum, M. *Thunbergia grandiflora* (Roxb. ex Rottler) Roxb. In: Ahmed ZU, Begum ZNT, Hassan MA, Khondker M, Kabir SMH, Ahmed M, et al. editors. Encyclopedia of flora and fauna of Bangladesh, Angiosperms: dicotyledons (Acanthaceae- Asteraceae). Dhaka: Asiatic Society of Bangladesh; 2008, 6; 74-75

- 207 [11] Yusuf, M. *Crotalaria pallid* Ait. In: Ahmed ZU, Hassan MA, Begum ZNT, Khondker M, Kabir SMH,
208 Ahmed M, et al. editors. Encyclopedia of flora and fauna of Bangladesh, Vol 8. Angiosperms:
209 dicotyledons (Fabaceae-Lythraceae). Dhaka: Asiatic Society of Bangladesh; 2009
- 210 [12] Kabir, MSH, Ahmad, S, Mahamoud, MS, Masum, MAA, Kamal, ATMM, Hoque, MA. Evaluation of
211 total condensed tannin content and anthelmintic activities of organic extracts of four Bangladeshi plants
212 on *Tubifextubifex* worm using in vitro method. *Int J Pharm*;2015, 5: 903 – 910,
- 213 [13]. Kar A, Goswami NK, Saharia D. Distribution and traditional uses of *Thunbergia retzius*
214 (Acanthaceae) in Assam, India. *Pleione*; 2008, 7: 325-32.
- 215 [14]. Obey JK, Anthoney ST. In vitro antimicrobial evaluation of ethanolic extract of *thunbergia alata*
216 leaves extract against selected microorganisms. *Int J Bioassays*, 2015, 4(10): 4418-22.
- 217 [15]. Oratai N. Diversity of medicinal plants for fever healing from Khao Phanom Benja National park,
218 Krabi Province. *KKU Sci J.*, 2014; 42(2): 313-26.
- 219 [16]. Aritajat S, Wutteerapol S, Saenphet K. Anti-diabetic effect of *Thunbergia laurifolia* Linn. aqueous
220 extract. *Southeast Asian J Trop Med Public health*, 2004; 35(2): 53-8.
- 221 [17]. Oonsivilai R, Ferruzzi MG, Ningsanond S. Antioxidant activity and cytotoxicity of Rang Chuet
222 (*Thunbergia laurifolia* Lindl.) extracts. *As J Food Ag-Ind.*, 2008; 1(2): 116-28.
- 223 [18]. Tangpng J, Satarug S. Alleviation of lead poisoning in the brain with aqueous leaf extract of the
224 *Thunbergia laurifolia* Linn. *Toxicol Lett*, 2010; 198(1): 83-88.
- 225 [19]. Pramyothin P, Chirdchupunsare H, Rungsipipat A, Chaichantipyuth C. Hepatoprotective activity of
226 *Thunbergia laurifolia* Linn extract in rat treated with ethanol: in vitro and in vivo studies. *J*
227 *Ethnopharmacol*, 2005; 102(3): 408-11.
- 228 [20] Kanchanapoom T, Kasai R, Yamasaki K. Iridoidglucosides from *Thunbergia laurifolia*,
229 *Phytochemistry*, 2002; 60(80): 769-771.
- 230 [21] Charles A, Joesph M, Ramani, VA. Chemical constituents and anti-oxidant activity of *Thunbergia*
231 *fragrans* leaf extracts. *Proceedings from National conference on evolutionary trends in biological and*
232 *pharmaceutical Chemistry*, 2014; p23.
- 233 [22] Thanislas P, Aruna A and Rathnakumar P. The effect of *Thunbergia fragrans* Roxb in experimental
234 diabetes and their effects on key metabolic enzymes involved in carbohydrate metabolism. *Int J Pharm*
235 *Sci Rev and Res*, 2015; 33(1): 68-73.
- 236 [23] Damtoft S, Frederiksen LB, Jensen SR. Alatoside and Thunaloside, Two Iridoidglucosides from
237 *Thunbergia alata*. *Phytochem*, 1994; 35(5): 1259 – 1261.
- 238 [24] Ismail LD, El-Azizi MM, Khalifa TI, Stermitz FR. Iridoid glycoside from *Thunbergiagrandiflora*.
239 *Phytochem*, 1996; 42(4): 1223 – 1225.
- 240 [25] SdrenRosendal J and Bent JN. Iridoids in *Thunbergia* Species *Phytochemistry*, 1989, 28 (11) 3059-
241 3061,

- [26] Olaoluwa OO, Moronkola DO, Kwenga S, Mgbeoji O. Essential oils constituents, phytochemical analyses, and antimicrobial studies on leaf and stem of *thunbergia laevis* nees. World journal of pharmacy and pharmaceutical sciences, 2016; 5(7), 224-237.
- [27] Antonisamy JM, Aparna JS, Jeeva S, Sukumaran S, Anantham B. Preliminary phytochemical studies on the methanolic flower extracts of some selected medicinal plants from India Asian Pacific Journal of Tropical Biomedicine (2012) S79-S82
- [28] Ismail LD, EL-Azli MM, Khalifat TI, Stermitz FR. Iridoid glycosides from *thunbergia grandiflora*. Phytochemistry 1996; 42 (4), 1223-1225.
- [29] Jeeva S, Johnson M, Aparna JS, Irudayaraj V. Preliminary phytochemical and anti-bacterial studies on flowers of selected medicinal plants. Int. J. Med. Arom. Plants 2011, 1(2) 107 - 114
- [30] Adams, RP. Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy. Allured Pub.Corp., Carol Stream, IL. 1995.
- [31] Prabhadevi, V, Sahaya SS., Johnson M, Venkatramani, B, Janakiraman, N; "Phytochemical studies on *Allamanda cathartica* L. using GC-MS," Asian Pacific Journal of Tropical Biomedicine, 2012;2 (2),550–554.
- [32] Botros RM, Galal TM, Farid AB, Mohamed MAA, Chemistry and Immuno modulatory Activity of Frankincense Oil, Z. Naturforsch 2003; 58 - 230.
- [33] "Topical analgesics introduction". Medicine.ox.ac.uk. 2003-05-26. Retrieved 2012-11-07.
- [34] Tzin V, Galili G. The Biosynthetic Pathways for Shikimate and Aromatic Amino Acids in *Arabidopsis thaliana*. *The Arabidopsis 2010, Book 8*:e0132.
- [35] Aparna, V.; Dileep, KV.; Mandal, PK.; Karthe, P.; Sadasivan, C.; Haridas, M. Anti-inflammatory property of n-hexadecanoic acid: Structural evidence and kinetic assessment. Chem. Biol. Drug Des. 2012, 80, 434–439.
- [36]. Kumar, PP.; Kumaravel, S.; Lalitha, C. Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*. Afr. J. Biochem. Res. 2010, 4, 191–195.