

Original Research Article

Evaluation of CNS depressant, antidiarrheal and antipyretic activities of ethanolic leaf extract of *Calophyllum inophyllum* L. on swiss albino mice.

Abstract

Objective

The main objective of the present study was to evaluate the CNS depressant, antidiarrheal and antipyretic activities of ethanolic leaf extract of *Calophyllum inophyllum* L.

Method

In the present study CNS depressant activity of *C. inophyllum* was carried out using hole-cross, hole-board and open-field models in swiss albino mice whereas castor-oil induced diarrhea was used to evaluate antidiarrheal activity. The antipyretic activity was conducted by Brewer's yeast-induced hyperthermia in mice.

Result

The experimental result indicates that the ethanolic extract of *C. inophyllum* reduced CNS depression significantly ($P = 0.05$) in tested animals and the observed effect was dose dependent as compared to standard drug diazepam. The plant extract at the dose of 250 and 500 mg/kg body weight showed significant inhibition 41.66% and 62.19% in castor oil induced diarrheal method. In yeast induced pyrexia, the plant extract displayed dose dependent inhibition as compared to paracetamol.

Conclusion

The investigated study concludes that *C. inophyllum* leaf has significant CNS depressant, antidiarrheal and antipyretic activities which could support to develop its medicinal applications.

Keywords: CNS depressant; antidiarrheal; antipyretic; *Calophyllum inophyllum*.

Introduction

Calophyllum inophyllum (clusiaceae), commonly known as "Indian laurel" or Alexandrian laurel is a large evergreen tree with a broad spreading crown of irregular branches and thick trunk covered with a rough, black and cracked bark [1,2,3]. The tree is usually 2 to 3m in height with a dense canopy of glossy, shiny, tough, elliptical leaves that are rounded at the bases [4, 5].

It has white flowers arranged in axillary cymes with a sweat lime-like fragrance [6]. Fruits are spherical drupe arranged in cluster having a single large seed. Once ripe, their smooth, yellow epidermis discloses a thin layer of pulp, which tastes somewhat of apple. The grey, ligneous and rather soft nut contains a pale yellow kernel, which is odorless when fresh. Once chewed, it coats the mouth and emulsifies saliva, and its insipid taste becomes bitter. It is widespread tree that is indigenous to tropical Asia and geographically distributed in Melanesia and Polynesia [3]. It grows widely in the coral sands and on the sea shore. The species was brought north to Hawaii from the South Pacific islands in early migrations of Polynesian settlers. Several parts of *Calophyllum inophyllum* is traditionally used in folk medicine for treating various health problems. In Java, the tree is said to have diuretic properties [7, 8]. In China, this plant is used in the treatment of eye diseases, wounds, rheumatisms, and inflammations in traditional Chinese folk medicine [9, 10]. This plant is used in Asia and India for orchitis and in the treatment for gonorrhea. The fresh leaves infusion of *Calophyllum inophyllum* are frequently used for treating bacterial infection, fungal infection and dried leaf for different kinds of skin problems, sores, cuts, and wounds [11,12,13,14]. In Cambodia, the leaves are inhaled as a treatment for migraines and vertigo. The bark is used as an antiseptic and disinfectant. In Cameroon the aqueous extracts of the root bark and leaves are used as a cicatrisant and are also used in the treatment of wounds and herpes [15]. The root bark and nut of *Calophyllum inophyllum* possessed the most significant cytotoxicity against the KB cell line and antimicrobial activities [15]. Seed oil from *Calophyllum inophyllum* has been used in folk medicine in the treatment of aching joints and rheumatism [16]. *Calophyllum inophyllum* possessed strong activity against human immunodeficiency virus type 1 [17, 18]. Some of these species are frequently employed in folk medicine to treat several injuries [2]. This plant has very antihyperlipidemic and antioxidant activity [1].

Preliminary phytochemical analysis of *Calophyllum inophyllum* resulted a wide variety of chemical constituents including Coumarinic derivatives: calophyllolide [20] , inophyllolids including inphyllum B ,P , C and E [15, 20, 21] , calophyllic acid containing calophynic acid [22] , brasiliensic acid and inophylloidic acid [23] . Xanthone derivative such as inoxanthone [15], caloxanthone A and B, macluraxanthine, 1,5-dihydroxyxanthone [24], caloxanthone D ,caloxanthone E , caloxanthone N [9], gerontoxanthone C [25] and 2-hydroxyxanthone [26] ,tannins ,alkaloids [27] , chalcones, benzofurans and triterpenes [4, 5, 28, 29, 30] flavonoid compound Amentoflavone [31,32].

2. Material and methods

2.1. Drugs and chemicals

Ethanol was bought from SIGMA chemical Co. St Louis, USA, Castor oil was purchased from WELL's Heath Care, Spain. Normal saline solution (0.9% NaCl) (Beximeo Infusion Ltd.), Diazepam (Incepta Pharmaceuticals Ltd.), loperamide manufactured by Square Pharmaceuticals Ltd., Bangladesh, was bought from local pharmacy. All the chemicals and reagents were analytical grade.

2.2. Plant materials

Fresh leaves of *C. inophyllum* leaves for experiment was collected by author from Botanical garden, Mirpur, Dhaka, Bangladesh. The plant was identified by the taxonomist of Bangladesh National Herbarium, Mirpur, Dhaka, Bangladesh (accession number DACB:39674) where voucher specimen was deposited as documentation.

2.3. Preparation of extract

250mg of plant extract was soaked in 1000 ml of 99% ethanol in clean, sterilized, and flat-bottomed glass container and shaken two times a day for 15 days at room temperature. Afterwards the plant extract was subjected to filtration with sterilized cotton material followed by Whatman No.1 filter paper. After filtration extract was evaporated using rotary evaporator (Bibby RE-200, Sterilin Ltd., UK) at 4 rpm and 65 °C temperature and the remaining gummy concentrate was taken as crude extract or ethanolic extract. The extract was stored in refrigerator at 4 °C until used for treatment [33].

2.4. Experimental animals

Weighted between 25-30g of young Swiss albino mice of either sex were collected from the animal Resource Branch of the International center For Diarrheal Disease and Research, Bangladesh (ICDDR, B). Animals placed under favorable environmental conditions maintained optimum temperature (25±2°C), humidity (45-55 %) and rodent food and water were provided formulated by the ICDDR, B. During experiment all animals received human care according to the guidelines of 'Guide for the Care and Use of Laboratory Animals', 8th edition, prepared by the National Academy of Sciences and published by the National Institute of Health (US).

2.5. CNS depressant activity test

2.5.1. Hole-cross test

The method was adopted as described by Hussain et al (34). A steel partition was fixed in the middle position of a cage having a size of 30 cm×20 cm×14 cm. A hole of 3 cm diameter was made in the partition for easy passage of mice in the cage. The mice were divided into control,

standard control and test group. The test groups received ethanolic extract of *C. inophyllum* at the dose of 500 and 250 mg/kg body weight orally whereas control group received vehicle (1% Tween 80 in water) at 10 mL/kg body weight orally and standard group received diazepam at a dose of 1 mg/kg body weight orally. The number of movement from one compartment to the other through the hole was counted at 0, 30, 60, 90 and 120 min after oral administration of test drugs for a period of 3 min.

2.5.2. Hole board test

The experiment was performed as described by Emran and Rahman (2014)⁽³⁵⁾. A 30 × 20 × 14cm³ size hole cross box was used to carry out the experiment. A steel partition was fixed in the middle of a cage. There was a 3 cm hole in the partition of the cage. The animals were divided into control, standard, and test groups containing five mice each. The test groups received the ethanolic extract of *C. inophyllum* orally at a dose of 250 and 500 mg/kg body weight whereas the normal control group received vehicle (1% Tween-80 in water orally). The number of movement through the hole from one compartment to the other was counted at 0, 30, 60, 90 and 120 min for a period of 3 min. Diazepam (1 mg/kg body weight i.p.) was used as positive control in the hole board test.

2.5.3. Open field test

The experiment was followed by the method described by Gupta et al. (1971)⁽³⁶⁾. The animals were divided into control, standard and test groups containing five mice each. The test groups received the ethanolic extract of *C. inophyllum* at a dose of 250 and 500 mg/kg body weight orally whereas the normal control group received vehicle (1% Tween-80 in water orally). The floor of open field was divided into a series of squares and each square alternately colored black and white. The apparatus of open field had a 40 cm height wall. Each mouse was placed at the center of the field and the number of squares traveled by the animals was counted at 0, 30, 60, 90 and 120 min for 3 min.

2.6. Antidiarrheal activity test

2.6.1. Castor oil-induced diarrhea

The method described by Shoba and Thomas (2001)⁽³⁷⁾ was followed for this study. Animals were divided into four groups of four mice each. Two test groups administered orally with ethanolic extract of *C. inophyllum* orally at a dose of 250 and 500 mg/kg body weight, respectively. Positive control group was administered loperamide 5 mg/kg as standard, while the control group was administered with only distilled water. One hour after administration of extract and drug each mouse was administered of standard dose of 2 ml of castor oil and housed in a separate cage where stools were collected on blotting paper sheet of uniform weight placed for collecting of fecal material. The number of fecal material in terms of both hard and soft pellet was counted up to 5 h at each hour interval. Percent inhibition (PI) was calculated as follows:

$$PI = \frac{\text{Mean defecation (control group-treated group)} \times 100}{\text{Mean defecation of control group}}$$

2.7. Antipyretic activity test

Brewer's yeast induced pyrexia

The antipyretic activity was evaluated by Brewer's yeast induced pyrexia in experimental animal. The animals were injected with 10 ml/kg body weight 20% aqueous suspension of brewer's yeast sub-cutaneously and rectal temperature of animals was recorded using an Ellab thermometer (33.19 ± 0.40 °C). The animals that after 18 h of subcutaneous administration showed an increase of $0.3\text{--}0.5$ °C in rectal temperature were selected for the antipyretic activity. Treated groups received ethanolic extract of *C. inophyllum* (500 mg/kg) orally, paracetamol (150 mg/kg) as reference drug whereas control group received distilled water (10 ml/kg) only. The rectal temperature was recorded at 1 h intervals up to 3 h after administration of drug⁽³⁹⁾.

2.8. Statistical analysis

Data obtained was analyzed by one-way ANOVA test followed by Dunnett's multiple comparisons post-hoc test using SPSS version 16.0 for Windows® and the results obtained were compared with the control group. $P < 0.05$ was considered to be statistically significant.

3. Results

3.1. CNS depressant activity

3.1.1. Hole-cross test

The table-1 demonstrates the result of hole-cross test of *C. inophyllum* leaves. 2nd observation period (30 min) manifests the locomotor lowering activity which was maintained up to the 5th observation period (120 min) for extracts. The extract showed the dose dependent lowering of locomotion activity which was comparable to standard drug diazepam. The extract showed dose dependent activity of 5.00 ± 0.41 and 3.50 ± 0.96 respectively at the dose of 250 and 500 mg/kg body weight whereas the standard drug diazepam showed 4.75 ± 1.38 at the dose of 1 mg/kg.

Table 1: Effect of ethanolic extract of the *C. inophyllum* leaves on hole cross test in mice

Group	Number of Movements (Mean±SEM)				
	0 min	30 min	60 min	90 min	120 min
Control	5.00 ± 1.08	5.25 ± 2.02	3.25 ± 0.63	2.25 ± 0.48	2.00 ± 0.71
Standard	6.00 ± 1.92	9.00 ± 3.42	8.00 ± 2.04	5.50 ± 1.85	4.75 ± 1.38
<i>C. inophyllum</i> (	$0.50 \pm 0.29^*$	1.75 ± 1.11	4.00 ± 0.71	2.50 ± 1.04	$5.00 \pm 0.41^*$

250 mg/kg)					
<i>C. inophyllum</i> (500 mg/kg)	5.00 ± 1.47	3.50 ± 0.65	4.00 ± 0.91	2.25 ± 0.63	3.50 ± 0.96

Values are reported as mean ± S.E.M. for group of five animals (n = 5). Values are analyzed as compared to control using one way ANOVA followed by Dunnett's post Hoc test. Asterisks indicated statistically significant values from control, * indicates P= 0.05, ** indicates P < 0.01 and *** indicates P < 0.001.

3.1.2. Hole-board test

The table-2 demonstrates the result of hole-board test of *C. inophyllum* leaves. After 90 and 120 minutes in hole board test, the extract at the dose 500 mg/kg body weight displayed a significant (p<0.05) number of head dipping behavior and the results were 13.50 ± 2.22 and 15.00 ± 3.29 respectively compared to control group. The Standard drug diazepam at 120 min showed significant (p<0.05) number of head dipping (21.50 ± 3.30) behavior.

Table 2: Effect of ethanolic extract of the *C. inophyllum* leaves on hole board test in mice

Group	Number of Movements (Mean±SEM)				
	0 min	30 min	60 min	90 min	120 min
Control	39.75 ± 6.71	23.75 ± 1.89	17.00 ± 2.08	20.75 ± 3.92	20.00 ± 1.87
Standard	47.50 ± 4.79	23.25 ± 2.84	27.25 ± 3.79	20.00 ± 1.35	21.50 ± 3.30
<i>C. inophyllum</i> (250 mg/kg)	28.75 ± 0.48	14.25 ± 2.29**	15.00 ± 4.02	17.50 ± 4.33	25.50 ± 2.10
<i>C. inophyllum</i> (500 mg/kg)	42.50 ± 1.04	13.00 ± 1.96*	8.00 ± 2.48*	13.50 ± 2.22	15.00 ± 3.29

Values are reported as mean ± S.E.M. for group of five animals (n = 5). Values are analyzed as compared to control using one way ANOVA followed by Dunnett's post Hoc test. Asterisks indicated statistically significant values from control, * indicates P= 0.05, ** indicates P < 0.01 and *** indicates P < 0.001.

3.1.3. Open Field Test

In open field test, the extract displayed a significant suppression of the number of squares traveled by the mice and the data has been represented into table-3. At 120 min the test sample showed remarkable dose dependant CNS depression activity. extract at the dose of 500 mg/kg (28.25 ± 5.76) body weight showed significant (p<0.001) reduction in the number of movement as compared to control group (39.75 ± 14.76) and standard group (144.3 ± 24.35).

191 **Table 3: Effect of ethanolic extract of the *C. inophyllum* leaves on open field test in mice**

Group	Number of Movements (Mean±SEM)				
	0 min	30 min	60 min	90 min	120 min
Control	122.3 ± 12.15	111.3 ± 10.44	97.00 ± 15.15	35.50 ± 9.579	39.75 ± 14.76
Standard	175.8 ± 10.27	169.3 ± 23.23	134.3 ± 18.96	96.25 ± 35.97	144.3 ± 24.35**
<i>C. inophyllum</i> (250 mg/kg)	161.8 ± 6.29	107.3 ± 7.857	82.50 ± 5.95	59.50 ± 5.74	45.00 ± 13.03
<i>C. inophyllum</i> (500 mg/kg)	135.5 ± 25.59	97.25 ± 12.46	73.75 ± 3.15	57.00 ± 8.89	28.25 ± 5.76

192 Values are reported as mean ± S.E.M. for group of five animals (n = 5). Values are analyzed as
 193 compared to control using one way ANOVA followed by Dunnett's post Hoc test. Asterisks
 194 indicated statistically significant values from control, * indicates P= 0.05, ** indicates P < 0.01
 195 and *** indicates P < 0.001.

196 3.2. Antidiarrheal activity

197 3.2.1. Castor oil-induced diarrhea

198 The results at table-4 showed that *C. inophyllum* leaves extract significantly (p<0.001) reduced
 199 the defecation number as compared to control group of diarrhea which was induced by castor oil.
 200 The extract treated group significantly (p<0.01) decreased the number of stools at 1 to 5 hours
 201 when compared to control group. *C. inophyllum*, showed 64.29% inhibition of diarrheal stools at
 202 the dose of 500 mg/kg body weight whereas the standard drug loperamide at the dose of 5mg/kg
 203 showed 71.43% diarrheal inhibitory activity.

204 **Table 4: Effect of ethanolic extract of *C. inophyllum* leaves on the castor oil-induced**
 205 **diarrhea in mice**

Group	Total number of feces	Number of diarrhea feces	Inhibition of diarrhea (%)
Control	11.50 ± 1.29	3.50 ± 0.58	-
Standard (Loperamide 5 mg/kg)	6.50 ± 1.26 ***	1.00 ± 0.82 **	71.43
<i>C. inophyllum</i> (250 mg/kg)	4.50 ± 2.44 **	2.00 ± 1.10 *	41.66

<i>C. inophyllum</i> (500 mg/kg)	3.15 ± 2.12 ***	1.12 ± 0.5 **	62.19
----------------------------------	-----------------	---------------	-------

206 Values are reported as mean ± S.E.M. for group of five animals (n = 5). Values are analyzed as
 207 compared to control using one way ANOVA followed by Dunnett's post Hoc test. Asterisks
 208 indicated statistically significant values from control, * indicates P= 0.05, ** indicates P < 0.01
 209 and *** indicates P < 0.001.

210 3.3. Antipyretic activity

211 *Brewer's yeast-induced pyrexia*

212 The results at table-4 showed that after 18th of Brewer's Yeast injection experimental mice
 213 displayed a increase in rectal temperature. The extract at the dose of 250 and 500 mg/kg body
 214 weight showed marked reduction of rectal temperature in 1h, 2h and 3h after administration of
 215 the extract and displayed peak anti-pyretic effect at the dose of 500 mg/kg in 3rd h (96.23 ±
 216 0.83°C, p<0.001) as compared to control (96.35 ± 0.44°C). Standard drug paracetamol showed
 217 significant anti-pyretic effect and reached the peak in 3rd h (96.35 ± 0.44°C, p<0.001) as
 218 compared to control group.

219 **Table 6: Effect of ethanolic extract of *C. inophyllum* leaves on Brewer's yeast-induced**
 220 **pyrexia in mice.**

Treatment	Dose (mg/kg)	Temperature in °C				
		Initial	Pyretic	1 h	2 h	3 h
Control	10	96.23 ± 0.49	99.00 ± 0.58	98.68 ± 0.85	96.05 ± 0.20	96.35 ± 0.44
Standard	150	97.35± 0.39*	97.30± 0.34*	96.58 ± 0.81**	95.33 ± 0.20	95.35 ± 0.31
<i>Calophyllum inophyllum</i> (250 mg/kg)	250	96.80 ± 0.26	98.55 ± 0.40	95.28 ± 0.81	96.63 ± 0.21	96.43 ± 0.31
<i>Calophyllum inophyllum</i> (500 mg/kg)	500	96.58 ± 0.55	99.70 ± 0.47	94.83 ± 0.79	95.45 ± 0.42	96.23 ± 0.83

221 Values are reported as mean ± S.E.M. for group of five animals (n = 5). Values are analyzed as
 222 compared to control using one way ANOVA followed by Dunnett's post t-test. Asterisks
 223 indicated statistically significant values from control, * indicates P= 0.05, ** indicates P < 0.01
 224 and *** indicates P < 0.001.

225 4. Discussion

226 Diarrhoea is gastrointestinal disorder, characterized by an excess fluid loss in the faeces results
 227 from an imbalance between the absorptive and secretory mechanisms in the intestinal tract (40).
 228 The liberation of ricinoleic acid from castor oil results in diarrheal effect by inducing
 229 permeability changes in mucosal fluid and electrolyte transport (41). The results of the present

study show that the extract of *C. inophyllum* exerted a significant reduction in the frequency of diarrhoea produced by castor oil. Since ricinoleic acid from castor oil stimulates motility and secretion by the release of prostaglandins (42). It can be assumed that the anti-diarrhoeal effect of ethanol and aqueous extracts may be due to the inhibition of prostaglandin biosynthesis. To obtain the effect of *C. inophyllum* on the central nervous system (CNS) activity, a number of methods namely open field, hole cross, hole-board were used (43).

In all the three experimental antipyretic **test calo ino** **gradually** decreased locomotor activity and produced a significant increase in sedative effect during the period of observation. The locomotor activity is a useful measure of the CNS awareness and any decrease of this activity resulting from depression of the central nervous system indicates a sedative effect(44). Sedation are principally mediated in the CNS by the GABAA receptor complex which are the most abundant inhibitory neurotransmitter in the central nervous system (CNS) (45) and it is commonly thought that central nervous system depressant drugs exert their actions through gamma-aminobutyrate (GABA)-mediated mechanisms (46).

All the above experimental analgesic methods reveal the potential central and peripheral antinociceptive activity of *C. inophyllum* and it is supposed to assume that the central activity of the plant via potentiating GABA inhibition. *C. inophyllum* having many bioactive compounds like flavonoids alkaloids, saponins and steroids are found to be ligands for the gamma aminobutyric acid type A (GABAA) receptors in the central nervous system (CNS) that provides further evidence(47).

Brewer's yeast-induced pyrexia was employed in evaluating antipyretic activities of ethanolic extracts of *C. inophyllum*. The present study showed that the ethanolic extract of *C. inophyllum* leaves caused a better antipyretic action in yeast induced elevation of body temperature. It is well known that the non-steroidal anti-inflammatory drugs control fever by the inhibition of prostaglandin synthesis within the hypothalamus that regulates body temperature (48). It may be assumed that the extract revealed their antipyretic action through inhibition of prostaglandin synthesis within the hypothalamus .

Conclusion

From the above study we can make a conclusion that *C. inophyllum* possess significant CNS depressant, anti-diarrheal and antipyretic activities and various phytochemicals present in the extract may attribute these activities. However, to be clear about the above pharmacological activities further research is needed.

References

1. Janki Prasad, Atul Shrivastava, A.K. Khanna, G. Bhatia, S.K. Awasthi, T. Narender: Antidyslipidemic and antioxidant activity of the constituents isolated from the leaves of *Calophyllum inophyllum* . Phytomedicine 19 (2012) 1245– 1249.

- 267 2. Saravanan. R, Dhachinamoorthi. D, Senthilkumar. K and Thamizhvanan. K: Antimicrobial
268 activity of various extracts from various parts of *Calophyllum inophyllum* L. .Journal of Applied
269 Pharmaceutical Science 01 (03); 2011: 102-106.
- 270 3. Mirza Danish Baig, Syed Basheeruddin, Silpa. S, A. Venkateshwara Reddy: Anti-
271 inflammatory Activity of Ethanolic Extracts of Leaf and Stem Bark of *Calophyllum inophyllum*
272 on Albino Wistar Rats. International Journal of Pharmaceutical Sciences and Drug Research
273 2014; 6(2): 174-177.
- 274 4. Ito C., Itoigawa M., Mishina Y., Cechinel Filho V., Enjo F., Tokuda H., Nishino H., and
275 Furukawa H. Chemical constituents of *Calophyllum brasiliense*, Structure of three new
276 coumarins and cancer chemo- preventive activity of 4-substituted coumarins. J. Nat. Prod. 2003;
277 66: 368-371.
- 278 5. Ito C., Itoigawa M., Mishina Y., Cechinel Filho V., Mukainaka T., Tokuda H., Nishino H.,
279 and Furukawa H. Chemical constituents of *Calophyllum brasiliense*: Structure elucidation of
280 seven new xanthenes and their cancer chemopreventive activity. J. Nat. Prod.2002; 65: 267-272.
- 281 6. Naveed Muhammad, Muhammad Saeed1 and Haroon Khan : Antipyretic, analgesic and anti-
282 inflammatory activity of *Viola betonicifolia* whole plant. BMC Complementary and Alternative
283 Medicine 2012, 12:59.
- 284 7. Kirtikar K. R. and Basu B. T., Indian Medicinal Plants, Bishen Singh Mahendra Pal Singh 23-
285 A, Connaught place, Dehradun- 248001, India, 1998, Vol.1, 2nd edition, 270- 274.
- 286 8. The wealth of India, A Dictionary of Indian raw materials and Industrial products,
287 Publications & Information directorate, CSIR, DR. K.S.Krishnan Marg, New Delhi- 110012,
288 1992, Vol-3:Ca-Ci , Revised edition, 68- 73.
- 289 9. H.Y. Chen, Flora of Hainanica, Vol. 2, (Science Press, Beijing, 1965), p. 56.
- 290 10. H.F. Dai and W.L. Mei, Modern Research on Medicinal Plants in Hainan, (China Science
291 and Technology Press, Beijing, 2007), p. 31.
- 292 11. Clatchey M.C., W. The Ethnopharmacopoeia of Rotuma J Ethnopharmacol, 1996, 50(3), 147-
293 156. [Source was an original research paper. Univ florida florida museum natural history
294 herbarium gainesville FL 32611 USA].
- 295 12. Khan N.U.D., Parveen, N., Singh, M.P., Singh, R., Achari, B., Dastidar, P.P.G and Dutta,
296 P.K., Two Isomeric Benzodipyranone Derivatives From *Calophyllum Inophyllum*,
297 Phytochemistry, 1996, 42 (4), 1181-1183.

- 298 13. Holdsworth, D.K., A Phytochemical Survey of Medicinal Plants of The D'entrecasteaux
299 Islands, Sci New Guinea, 1974, 2(2), 164-171.
- 300 14. Holdsworth D. and Wamoi B., Medicinal Plants of the Admiralty Islands, Papua New
301 Guinea. PART I, Int J Crude Drug Res, 1982, 20(4), 169-181.
- 302 15. Marie C. Yimdjio, Anatole G. Azebaze, Augustin E. Nkengfack, A. Michele Meyer ,
303 Bernard Bodo, Zacharias T. Fomum: Antimicrobial and cytotoxic agents from *Calophyllum*
304 *inophyllum*. Phytochemistry 65 (2004) 2789–2795.
- 305 16. Claude Spino, Marco Dodier, and Subramaniam Sotheeswaran: ANTI-HIV COUMARINS
306 FROM CALOPHYLLUM SEED OIL. Bioorganic & Medicinal Chemistry Letters 8 (1998)
307 3475-3478.
- 308 17. Patil, A.D., Freyer, A.J., Eggleston, D.S., Haltiwanger, R.C., Bean, M.F., Taylor, P.B.,
309 Caranfa, M.J., Brenn, A.L., Bartus, H.R., Johnson, R.K., Hertzberg, R.P., Westley, J.W., 1993.
310 The inophyllums, novel inhibitor of HIV-1 reverse transcriptase isolated from the Malaysian tree,
311 *C. inophyllum* L. J. Med. Chem. 36, 4132– 4138.
- 312 18. Kashman, Y., Gustafson, K.R., Fuller, R.W., Cardellina, J.H., McMahon, J.B., Currens, M.J.,
313 Buckheit Jr., R.W., Hughes, S.H., Cragg, G.M., Boyd, M.R., 1992. The Calanolides, a novel
314 HIV – inhibitory class coumarin derivatives from the tropical rain forest tree, *C. lanigerum*. J.
315 Med. Chem. 35, 2735–2743.
- 316 20. Polonsky J, 1957. Structure chimique du calophyllolide, de l'inophyllolide et de l'acide
317 calophyllique, constituants des noix de *Calophyllum inophyllum*. Bull. Soc. Chem. Fr., 1079–
318 1087.
- 319 21. Kawazu K, Ohigashi H, Mitsui T, 1968: The piscicidal constituents of *Calophyllum*
320 *inophyllum*. Tetrahedron Lett. 19, 2383–2385.
- 321 22. Gautier J, Kunesch G, Polonsky J, 1972: Structure of calophynic acid, a Novel constituent of
322 *Calophyllum inophyllum*. Tetrahedron Lett. 27, 2715–2718.
- 323 23. Stout G.H, Krahn M.M, Breck G.D, 1968: Calophyllum products. II. Brasiliensic and
324 inophylloidic acids. Tetrahedron Lett. 29, 3285– 3290.
- 325 24. Iinuma, M, Tosa, H, Tanaka, T, Yonemori, S, 1994: Two xanthones from root bark of *C.*
326 *inophyllum*. Phytochemistry 35, 527–532.
- 327 25. C.H. Chang, C.C. Lin, H. Masao, and N. Tsuneo, Phytochemistry 28, 595 (1989).
- 328 26. C. Gnerre, U. Thull, P. Gaillard, P.A. Carrupt, B. Testa, E. Fernandes, F. Silva, M. Pinto,

- 329 M.M.M. Pinto, J.L. Wolfender, K. Hostettmann, and G. Cruciani, *Helv. Chim. Acta* 84, 552
330 (2001).
- 331 27. Okemo PO. Antimicrobial efficacy of selected medicinal plants used by Kenyan Herbal
332 doctors. Ph.D. thesis, Kenyatta University of Nairobi, (1996) pp. 173-90.
- 333 28. Da Silva K. L., Santos A. R. S., Mattos P. E. O., Yunes R. A., Delle-Monache F., and
334 Cechinel Filho V. Chemical composition and analgesic activity of *Calophyllum brasiliense*
335 leaves. *Therapie*. 2001; 56: 431- 434.
- 336 29. Oger J. M., Morel C., Helesbeux J. J., Litaudon M., Seraphin D., Dartiguelongue C., Larcher
337 G., Richomme P., and Dural O. First 2- hydroxy-3-methylbut-3-enyl substituted xanthenes
338 isolated from plants: structure elucidation, synthesis and antifungal activity. *Nat. Prod. Res.*2003;
339 17: 195-199.
- 340 30. Isaias D. E. B., Niero R., Noldin V. N., Buzzi F. C., Yunes Inst R. A., Delle-Monache F., and
341 Cechinel Filho V. Pharmacological and phytochemical investigations of different parts of
342 *Calophyllum brasiliense* (Clusiaceae). *Die Pharmazie*. 2004; 59 (11): 879-881.
- 343 31. Desai, P.D., Dutia, M. D., Ganguly, A.K., Govindachari, T.R., Joshi, B.S., Kamat, V.N.,
344 Prakash, D., Rane, D.F., Sathe, S.S. and Viswanathan, N., Chemical Study of Some Indian
345 Plants. Part III, *Indian J Chem*, 1967, 5, 523.
- 346 32. Goh, S.H., Jantan, I. and Waterman, P.G., Neoflavonoid and Biflavonoid Constituents of
347 *Calophyllum Inophylloide*, *J Nat Prod*, 1992, 55 (10), 1415-1420.
- 348 33. Baruah DB, Dash RN, Chaudhari MR, Kadam SS. Plasminogen activators: a comparison.
349 *Vascul Pharmacol* 2011; 44: 1-9.
350
- 351 34. Hussain J, Ur Rehman N, Hussain H, Al-Harrasi A, Ali L, Rizvi TS. **Analgesic, anti-**
352 **inflammatory, and CNS depressant activities of new constituents of Nepeta clarkei.**
353 *Fitoterapia*. 2012; 83(3): 593- 598.
- 354 35. Emran TB, Rahman MA. **Sedative, anxiolytic and analgesic effects of Urena sinuata L.**
355 **leaf extract in animal models.** *Int Food Res J*. 2014; 21(5): 2069-2075.
- 356 36. Gupta BD, Dandiya PC, Gupta ML. **A psychopharmacological analysis of behavior in rat.**
357 *Jpn J Pharmacol*. 1971; 21: 293.
358
- 359 37. Shoba FG, Thomas M. **Study of antidiarrheal activity of four medicinal plants in castor**
360 **oil-induced diarrhea.** *J. Ethnopharmacol*. 2001; 76(1): 73-76.
- 361 38. Chatterjee TK. Handbook on laboratory mice and rats. 1st ed. Kolkata, Department of
362 Pharmaceutical Technology, Jadavpur University, 1993; p 157.

- 363 39. Bhowmick R, Sarwar MS, Dewan SMR, Das A, Das B, Uddin MMN, Islam MS. **In vivo**
364 **analgesic, antipyretic, and anti-inflammatory potential in Swiss albino mice and in vitro**
365 **thrombolytic activity of hydroalcoholic extract from Litsea glutinosa leaves.** Biol Res. 2014;
366 47: 1-8.
- 367 (40) K.K. Senthil Kumar. **Pharmacological studies of anti diarrhoeal activity of Casuarina**
368 **equisetifolia (L.) in experimental animals.** AJPST Vol 1|Issue 1| 2011 |8-11.
- 369 (41) N. Venkatesan, Vadivu Thiyagarajan, Sathiya Narayanan, et.all. **Anti-diarrhoeal**
370 **potential of asparagus racemosus wild root extracts in laboratory animals.**
371 **J Pharm Pharmaceut Sci 8(1):39-46, 2005**
- 372 (42) Havagiray R. Chitme, Ramesh Chandra, Sadhna Kaushik. **Studies on anti-diarrhoeal**
373 **activity of calotropis gigantea r.br. in experimental animals.** J Pharm Pharmaceut Sci
374 **7(1):70-75, 2004**
- 375 (43) Nusrat Subhan, Mohammad Ashraful Alam, Firoj Ahmed et all. **Bioactivity of Excoecaria**
376 **agallocha.** Revista Brasileira de Farmacognosia Brazilian Journal of Pharmacognosy 18(4): 521-
377 526, Out./Dez. 2008
- 378 (44) Farjana Sharmen, Adnan Mannan, Md. Mominur Rahman, et all. **Investigation of in vivo**
379 **neuropharmacological effect of Alpinia nigra leaf extract.** Asian Pac J Trop Biomed 2014;
380 4(2): 137-142.
- 381 (45) file:///C:/Users/POLY/Downloads/C132_1923-1930.pdf
- 382 (46) Verma A, Jana GK, Sen S, Chakraborty R, Sachan S, Mishra A. **Pharmacological**
383 **evaluation of Saraca indica Leaves for central nervous system depressant activity in mice.** J
384 Pharm Sci Res. 2010; 2(6): 338-343.
- 385 (47) Verma A, Jana GK, Sen S, Chakraborty R, Sachan S, Mishra A. **Pharmacological**
386 **evaluation of Saraca indica Leaves for central nervous system depressant activity in mice.** J
387 Pharm Sci Res. 2010; 2(6): 338-343.
- 388 (48) Yousef A. Taher, Awatef M. Samud, Fathy E. El-Taher, et all. **Experimental evaluation of**
389 **anti-inflammatory, antinociceptive and antipyretic activities of clove oil in mice.** Libyan
390 Journal of Medicine 2015, 10: 28685.