

# EFFECT OF RAW QUAIL EGG ON INTRAOCULAR PRESSURE AND BLOOD PRESSURE OF HYPERTENSIVE SUBJECTS

## ABSTRACT

Quail egg is a rich source of vitamins, mineral salts, enzymes and amino acids. In addition to these nutritional values, quail egg has been reported to have medicinal properties such as hypotensive property. This study investigated the effect of fresh raw quail egg on intraocular pressure (IOP) and blood pressure (BP) of hypertensive and normal subjects in a Nigerian population. One hundred and fifty newly diagnosed and untreated mild to moderate hypertensive subjects selected from the screening exercise carried out at the University of Benin Optometry Clinic, Edo State, Nigeria were recruited for this study. They comprised three groups of fifty subjects each, which were systemic hypertensives, ocular hypertensives and normotensive control subjects. IOP and BP were measured and recorded before and after oral administration of 0.6ml/kg body weight of raw quail egg to each subject in the three study groups at 30 minutes interval for 90 minutes. Results showed that the maximum mean difference in IOP after oral administration of 0.6ml/kg body weight of fresh raw quail egg, was  $1.85 \pm 0.12\text{mmHg}$  and  $1.90 \pm 0.05\text{mmHg}$  respectively which occurred at 90 minutes in the normotensive right and left eyes, these were statistically significant ( $p < 0.003$ ). The maximum mean difference in BP was  $15.67 \pm 0.08\text{mmHg}$  which occurred at 60 minutes in the normotensive subjects and it was statistically significant ( $p = 0.000$ ). The hypotensive effect of 0.6ml/kg body weight of fresh raw quail egg was similar in the three study groups, the peak effect on IOP occurred at 90 minutes and this was preceded by the peak effect on BP which occurred at 60 minutes. The hypotensive effect on IOP may have resulted from the fall in BP. In conclusion, 0.6ml/kg body weight of raw quail egg has significant hypotensive effect on intraocular pressure and blood pressure compared to water, when administered orally on empty stomach in ocular hypertensives,

24 systemic hypertensives and in the normotensive control groups. Therefore it may be consumed as a  
25 form of supplement in the control of high blood pressure and raised intraocular pressure.

26 Keywords: raw quail egg, intraocular pressure, blood pressure, hypertension.

## 27 1. INTRODUCTION

28 The prevalence and diagnosis of ocular and systemic hypertension are growing at an  
29 alarming rate especially in the Nigerian society [1]. Ocular hypertension (OHT) is defined as  
30 intraocular pressure higher than normal, in the absence of optic nerve damage or visual field loss  
31 [2]. Intraocular pressure (IOP) is controlled primarily by the rate of secretion and the rate of  
32 drainage of the aqueous humor. Whereby these two factors do not commensurate, there will be  
33 build up pressure leading to increased risk of optic nerve damage; a pathological condition called  
34 glaucoma [3]. The number of people with glaucoma worldwide is expected to rise from 64 million  
35 to 76 million in 2020 and 111 million in 2040, with Africa and Asia being affected more heavily  
36 than the rest of the world [3]. Normal IOP is between 10-20 mmHg while its average value is 15.5  
37 mmHg with fluctuations of about 2.75 mmHg. IOP varies with the time of the day, heartbeat,  
38 respiration and blood pressure [2, 4]. It can be influenced by corneal thickness and rigidity [5, 6].

39 Previous researchers have reported that changes in blood pressure result in changes in  
40 intraocular pressure in humans and animals studies [7-12]. The term 'blood pressure' refers to  
41 arterial blood pressure which is the lateral pressure exerted by the contained column of blood on the  
42 wall of arteries. It is expressed in four different terms: systolic blood pressure, diastolic blood  
43 pressure, pulse pressure and mean arterial blood pressure. Normal systolic pressure is 120 mmHg; it  
44 ranges between 110 to 140mmHg, while normal diastolic pressure is 80 mmHg and varies between  
45 60 and 80 mmHg [13]. Systemic hypertension (SHT) can be defined as a sustained rise in blood  
46 pressure, which can lead to other health complications like stroke and heart attack. A person is said

47 to have systemic hypertension when the mean arterial pressure is greater than the upper range of the  
48 accepted normal measure of 110 mmHg or diastolic pressure greater than 90 mmHg and systolic  
49 pressure greater than 135mmHg [14] . Adeloye *et al.*, (2015) [15] estimated about 20.8 million  
50 cases of systemic hypertension in Nigeria among people aged at least 20 years in 2010, with a  
51 prevalence of 28.0% (24.6, 31.9) in both sexes, 30.7% (24.9, 33.7) among men and 25.2% (22.7,  
52 31.9) among women.

53 Most orthodox medicines used in the treatment of ocular and systemic hypertension have  
54 unwanted adverse effects on patients suffering from these conditions. Therefore most patients prefer  
55 the use of alternative therapy such as natural food supplements to orthodox medicines. Quail egg is  
56 a universal natural food supplement with no health implications and safe to use. It is beneficial in  
57 healthy living because it has been proven to alleviate symptoms of diabetes, hypertension, high  
58 serum cholesterol, arteriosclerosis, asthma, kidney, liver, and gallbladder stones. It strengthens the  
59 immune system, promotes healthy memory, increases brain activity and stabilizes the nervous  
60 system [16-18]. Quail egg is a small speckled egg. Though the chicken egg is five times larger than  
61 quail egg, quail egg contains 7.5 times more Iron, 6 times more vitamin B2 and 5 times more  
62 phosphorus than chicken egg. The egg is a rich source of vitamins, mineral salts, enzymes, fatty  
63 acids and amino acids. Quail egg contains on the average: (per 100 g whole liquid egg), 0.12 mg of  
64 Vitamin B1, 0.85 mg of Vitamin B2, 300 IU of Vitamin A, and 0.10 mg of Niacin. 13.1 g of  
65 Protein, 11.2 g of Fat, 76mg of cholesterol, 84mg of omega-6-fatty acid, 4mg of omega-3-fatty  
66 acid, 59 mg of Calcium, 119mg of potassium, 127mg of sodium, 220 mg of Phosphorus, and 3.8  
67 mg of Iron [19].

68 Quail egg works best when consumed fresh and raw. 240 eggs are needed for one course of  
69 therapy in improving the following conditions: Nervous disorders, Anemia, Migraine, Diabetes,  
70 High blood pressure, Arteriosclerosis, Bronchial asthma, and improving the memory. Instruction

71 for use: Take the egg raw, on an empty stomach, in the morning, half an hour before breakfast, in  
72 the following manner: For adult: 1st -3rd day 3 eggs daily; 4th day on 5-6 eggs daily until  
73 completed therapy. For children: aged: 6-10years, 4 eggs daily (90 eggs are needed for one course  
74 of therapy); 3-6years, 3 eggs daily (60 eggs are needed for one course of therapy); 1-2years, 2 eggs  
75 daily (60 eggs are needed for one course of therapy). 3months-1year, 1 egg daily (30 eggs are  
76 needed for one course of therapy). Wait for six months before repeating therapy. Two courses of  
77 therapy are recommended yearly [19].

78 Previous study has shown that fresh raw quail egg reduces blood glucose level when taken  
79 on empty stomach in hyperglycemia [20] and there are speculations that it lowers blood pressure in  
80 systemic hypertension, but there are no documented experimental studies on its effect on human  
81 blood pressure and intraocular pressure. Therefore this study investigated the effect of oral  
82 administration of 0.6ml/kg body weight of fresh raw quail egg on intraocular pressure and blood  
83 pressure of fifty ocular hypertensives (OHT), fifty systemic hypertensives (SHT) and fifty  
84 normotensive (NT) control subjects; males and females selected from a screening exercise  
85 conducted at the Department of Optometry, University of Benin, Benin City, Nigeria using a  
86 purposive sampling technique.

## 87 2. MATERIALS AND METHODS

### 88 Subjects

89 A screening exercise was conducted at the Department of Optometry, University of Benin,  
90 Benin City, the case history of each subject was taken and each subject was made to go through  
91 general examination of blood pressure measurement and ocular examination of the anterior and  
92 posterior segments of the eyes. The inclusion criteria for the systemic hypertensives selected were  
93 those newly diagnosed from the screening exercise who had not been on any treatment. They had

94 systolic BP of 140-160mmHg, diastolic BP of 90-105mmHg, and average IOP less than 21 mmHg  
95 while the normotensives selected had an average systolic BP less than 140 mmHg, diastolic BP less  
96 than 90 mmHg, and average IOP less than 21 mmHg without treatment. All the subjects had no  
97 other systemic or ocular disorders. Their blood pressure was measured three consecutive times with  
98 U-MEC mercurial sphygmomanometer manufactured in China by Medicare Instrument Ltd, and  
99 Littmann stethoscope manufactured in USA by 3M Health Care. A mean of the three readings was  
100 used in the diagnosis of those with systemic hypertension and those who have normal blood  
101 pressure, during the screening exercise. The inclusion criteria for the ocular hypertensives selected  
102 were those newly diagnosed from the screening exercise and had not been on any treatment. They  
103 had average IOP measured at the three time points greater than 21 mmHg, average systolic BP less  
104 than 140 mmHg, diastolic BP less than 90 mmHg. Their IOP was measured with CT 20 non-contact  
105 computerized tonometer manufactured by Topcon in Japan. The mean of three readings was  
106 recorded for each subject. The three readings were taken at 9am, 3pm, and 6pm in order to observe  
107 the diurnal variations in the IOP of each subject before selection. Central corneal thickness (CCT)  
108 was measured for all the subjects with SW-1000P ultrasound pachymeter manufactured in China by  
109 Tianjin Electronic Technology. Ten readings were continuously taken and the average was recorded  
110 as the CCT. Each Subject's IOP was adjusted and corrected for CCT using the Ehler's formula. The  
111 subjects selected for the three groups had visual field screening using Octopus 900 manufactured in  
112 USA by Haag-streit Company and those selected had no visual field defects.

113 One hundred and fifty subjects, males and females were selected from the screening  
114 exercise using a purposive sampling technique. All the subjects selected were non-alcoholics and  
115 non-smokers and those on previous medication for systemic or ocular disorders were excluded from  
116 the study. They were divided into three groups of fifty each. Group 'A' comprised ocular  
117 hypertensives of twenty males and thirty females, aged 35 -52 (mean age  $45 \pm 3.64$ ) years. Group

118 'B' comprised systemic hypertensives of twenty eight males and twenty two females, aged 45 -58  
119 (mean age  $50 \pm 4.82$ ) years, while Group 'C' comprised normal healthy subjects of twenty five  
120 males and twenty five females, used as the control group, aged 40 -55 (mean age  $46 \pm 4.20$ ) years.  
121 Informed consent was obtained from each of the participant after a detailed explanation of the  
122 procedure was given to them. The study was approved by the Ethics committee of the University of  
123 Benin Teaching Hospital, Benin City, Edo State Nigeria and was performed in accordance with the  
124 Declaration of Helsinki of 1996. All experiments were carried out at the University of Benin  
125 Optometry Clinic, Benin City, Edo State. All subjects were instructed to abstain from all  
126 medication, liquid food, water, juice and beverages in the morning before presenting for the  
127 experiment since these may affect readings. All experiments commenced at 9.00am every morning.

## 128 **Procedure**

129 Day 1 of the experiment, each subject was weighed and body weight recorded. Baseline  
130 intraocular pressure was measured three consecutive times with CT 20 non-contact computerized  
131 tonometer. The average of three readings was recorded for each subject. Baseline blood Pressure  
132 was measured three consecutive times with U-MEC mercurial sphygmomanometer and Littmann  
133 stethoscope. The average of the three readings was recorded for each subject. Thereafter a volume  
134 of 0.6ml/kg body weight of Eva water was administered orally on empty stomach, to each subject in  
135 the three study groups according to their body weight; and their IOP and BP were measured at 30  
136 minutes interval for 90 minutes and the results recorded.

137 Day 2 of the experiment, each subject was weighed and body weight recorded. Intraocular  
138 pressure and blood pressure were measured again at baseline. Thereafter a dose of 0.6ml/kg body  
139 weight of fresh raw quail egg was administered orally on empty stomach, to each subject in the

three study groups according to their body weight; IOP and BP were measured again at 30 minutes interval for 90 minutes using the above mentioned instruments and the results recorded.

### Statistical Analysis

All the data in this study were analyzed with IBM SPSS version 20. Analysis of Variance (ANOVA) was used to determine if quail egg and water had significant effect on intraocular pressure and blood pressure. Least Significant Difference (LSD) was used to determine at what time the effect became significant.

### 3. RESULTS

The results are summarized in Tables. Table 1 showed that after oral administration of 0.6ml/kg body weight of Eva water to the three study groups, there were no significant changes ( $p>0.05$ ) in intraocular pressure and blood pressure at 30, 60 and 90 minutes compared to the baseline values, in systemic and ocular hypertensives as well as in normotensive control subjects

**Table 1: effect of 0.6ml/kg body weight of Eva water on mean intraocular pressure (Right and Left Eyes) and mean arterial blood pressure in ocular hypertensives, systemic hypertensives and normotensive subjects.**

	Time interval	N	Mean of pressure (mmHg) $\pm$ S.E.M. (ocular hypertensives)	Mean of pressure (mmHg) $\pm$ S.E.M. (systemic hypertensives)	Mean of pressure (mmHg) $\pm$ S.E.M. (Normotensives)
<b>Right Eye</b>	Base Line	50	27.61 $\pm$ 0.32	16.50 $\pm$ 0.40	14.80 $\pm$ 0.38
	30 minutes		27.59 $\pm$ 0.31	16.45 $\pm$ 0.39	14.77 $\pm$ 0.35
	60 minutes		27.59 $\pm$ 0.31	16.45 $\pm$ 0.39	14.75 $\pm$ 0.34
	90 minutes		27.58 $\pm$ 0.30	16.45 $\pm$ 0.39	14.75 $\pm$ 0.34
<b>Left Eye</b>	Base Line	50	26.12 $\pm$ 0.44	17.52 $\pm$ 0.55	13.80 $\pm$ 0.30
	30 minutes		26.11 $\pm$ 0.43	17.50 $\pm$ 0.52	13.77 $\pm$ 0.28
	60 minutes		26.08 $\pm$ 0.42	17.48 $\pm$ 0.48	13.77 $\pm$ 0.28
	90 minutes		26.08 $\pm$ 0.42	17.48 $\pm$ 0.48	13.77 $\pm$ 0.28
<b>Mean arterial BP</b>	Base Line	50	90.50 $\pm$ 1.15	105.20 $\pm$ 1.36	88.50 $\pm$ 0.70
	30 minutes		90.48 $\pm$ 1.13	105.17 $\pm$ 1.34	88.48 $\pm$ 0.67
	60 minutes		90.45 $\pm$ 1.12	105.17 $\pm$ 1.34	88.46 $\pm$ 0.67

	90 minutes		90.46 ±1.12	105.17 ±1.34	88.48 ±0.65
--	------------	--	-------------	--------------	-------------

157

158 Table 2 showed that the maximum mean difference in IOP was 0.05 ±0.01mmHg and 0.05  
159 ±0.04mmHg respectively which occurred at 90 minutes in the right eye of the systemic  
160 hypertensives and the normotensive subjects and it was statistically insignificant ( $p>0.05$ ). The  
161 maximum mean difference in BP was 0.05 ±0.03mmHg which occurred at 60 minutes in the ocular  
162 hypertensive subjects and it was statistically insignificant ( $p>0.05$ ).

163 **Table 2: Mean difference in pressure between baseline and different time of assessment after**  
164 **oral administration of 0.6ml/kg body weight of Eva water in OHT, SHT and NT subjects.**

	Time interval	N	Mean difference in pressure (mmHg) ± S.E.M. (ocular hypertensives) with their p-values	Mean difference in pressure (mmHg) ± S.E.M. (systemic hypertensives) with their p-values	Mean difference in pressure (mmHg) ± S.E.M. (Normotensives) with their p-values
Right Eye	30 minutes	50	0.02 ±0.01 (p=0.087)	0.05 ±0.01 (p=0.097)	0.03 ±0.03 (p=0.075)
	60 minutes		0.02 ±0.01 (p=0.087)	0.05 ±0.01 (p=0.097)	0.05 ±0.04 (p=0.085)
	90 minutes		0.03 ±0.02 (p=0.078)	0.05 ±0.01 (p=0.097)	0.05 ±0.04 (p=0.085)
Left Eye	30 minutes	50	0.01 ±0.01 (p=0.070)	0.01 ±0.03 (p=0.071)	0.03 ±0.02 (p=0.078)
	60 minutes		0.04 ±0.02 (p=0.083)	0.04 ±0.05 (p=0.087)	0.03 ±0.02 (p=0.078)
	90 minutes		0.04 ±0.02 (p=0.083)	0.04 ±0.05 (p=0.087)	0.03 ±0.02 (p=0.078)
Mean arterial BP	30 minutes	50	0.02 ±0.02 (p=0.68)	0.03 ±0.02 (p=0.080)	0.02 ±0.03 (p=0.078)
	60 minutes		0.05 ±0.03 (p=0.074)	0.03 ±0.02 (p=0.080)	0.04 ±0.03 (p=0.078)
	90 minutes		0.04 ±0.03 (p=0.098)	0.03 ±0.02 (p=0.080)	0.02 ±0.05 (p=0.070)

165

166 But in Tables 3 and 4, with oral administration of 0.6ml/kg body weight of fresh raw quail  
167 egg, ANOVA showed that 0.6ml/kg body weight of fresh raw quail egg had significant effect  
168 ( $p<0.05$ ) on intraocular pressure and blood pressure in systemic and ocular hypertensives as well as  
169 in normotensive control subjects. LSD showed that the peak effect on mean IOP occurred at 90  
170 minutes and this was statistically significant ( $p=0.000$ ) in the three study groups. It was preceded  
171 by the peak effect on mean arterial blood pressure which occurred at 60 minutes of ingesting



172 0.6ml/kg body weight of fresh raw quail egg, this effect was also statistically significant ( $p=0.000$ )

	Time interval	N	Mean of pressure (mmHg) $\pm$ S.E.M. (ocular hypertensives)	Mean of pressure (mmHg) $\pm$ S.E.M. (systemic hypertensives)	Mean of pressure (mmHg) $\pm$ S.E.M. (Normotensives)
<b>Right Eye</b>	Base Line	50	28.61 $\pm$ 0.29	15.41 $\pm$ 0.39	13.68 $\pm$ 0.26
	30 minutes		27.90 $\pm$ 0.27	14.82 $\pm$ 0.38	12.91 $\pm$ 0.25
	60 minutes		27.34 $\pm$ 0.28	14.23 $\pm$ 0.36	12.39 $\pm$ 0.24
	90 minutes		26.88 $\pm$ 0.26	13.59 $\pm$ 0.33	11.83 $\pm$ 0.24
<b>Left Eye</b>	Base Line	50	27.72 $\pm$ 0.33	16.62 $\pm$ 0.43	13.53 $\pm$ 0.23
	30 minutes		26.92 $\pm$ 0.31	15.90 $\pm$ 0.41	12.87 $\pm$ 0.22
	60 minutes		26.39 $\pm$ 0.30	15.38 $\pm$ 0.40	12.30 $\pm$ 0.22
	90 minutes		25.91 $\pm$ 0.31	14.86 $\pm$ 0.38	11.63 $\pm$ 0.20
<b>Mean arterial BP</b>	Base Line	50	92.87 $\pm$ 1.05	107.92 $\pm$ 1.37	89.21 $\pm$ 0.91
	30 minutes		88.06 $\pm$ 1.01	104.65 $\pm$ 1.26	84.21 $\pm$ 0.87

173 in the three study groups. The peak effect on blood pressure occurred 30 minutes before the peak  
174 effect on intraocular pressure. Thereafter the blood pressure started rising gradually.

175

176

177

178

179 **Table 3: Effect of 0.6ml/kg body weight of raw quail egg on mean intraocular pressure (Right**  
180 **and Left Eyes) and mean arterial blood pressure in OHT, SHT and NT subjects**

	60 minutes		77.33 ±0.97	95.57 ±1.21	73.54 ±0.78
	90 minutes		82.60 ±0.92	98.17 ±1.23	79.16 ±0.83

181

182 **Table 4: Mean difference in pressure between baseline and different time of assessment after**  
183 **oral administration of 0.6ml/kg body weight of raw quail egg in OHT, SHT and NT subjects.**

	Time interval	N	Mean difference in pressure (mmHg) ± S.E.M. (ocular hypertensives) with their p-values	Mean difference in pressure (mmHg) ± S.E.M. (systemic hypertensives) with their p-values	Mean difference in pressure (mmHg) ± S.E.M. (Normotensives) with their p-values
<b>Right Eye</b>	30 minutes	50	0.71 ±0.02 (p=0.072)	0.58 ±0.01 (p=0.092)	0.77 ±0.01 (p=0.063)
	60 minutes		1.27 ±0.01 (p=0.001)	1.18 ±0.03 (p=0.008)	1.29 ±0.02 (p=0.000)
	90 minutes		1.73 ±0.13 (p=0.000)	1.82 ±0.06 (p=0.001)	1.85 ±0.12 (p=0.000)
<b>Left Eye</b>	30 minutes	50	0.80 ±0.01 (p=0.070)	0.72 ±0.02 (p=0.071)	0.66 ±0.01 (p=0.068)
	60 minutes		1.33 ±0.01 (p=0.003)	1.24 ±0.03 (p=0.048)	1.23 ±0.01 (p=0.005)
	90 minutes		1.81 ±0.11 (p=0.000)	1.76 ±0.05 (p=0.002)	1.90 ±0.03 (p=0.000)
<b>Mean arterial BP</b>	30 minutes	50	4.81 ±0.04 (p=0.001)	3.27 ±0.11 (p=0.008)	5.00 ±0.04 (p=0.000)
	60 minutes		15.54 ±0.08 (p=0.000)	12.35 ±0.16 (p=0.000)	15.67 ±0.05 (p=0.000)
	90 minutes		10.27 ±0.13 (p=0.000)	9.75 ±0.14 (p=0.000)	10.05 ±0.04 (p=0.000)

184

185 The maximum mean difference in IOP for the right eye in OHT, SHT and NT subjects after  
186 oral administration of 0.6ml/kg body weight of fresh raw quail egg, was 1.73 ±0.13, 1.82 ±0.06 and  
187 1.85 ±0.12mmHg respectively. Likewise the maximum mean difference in IOP for the left eye in  
188 OHT, SHT and NT subjects were 1.81 ±0.11, 1.76 ±0.05 and 1.90 ±0.05mmHg respectively which  
189 occurred at 90 minutes and these were statistically significant (p<0.003). The maximum mean  
190 difference in BP in OHT, SHT and NT subjects were 15.54 ±0.08 mmHg, 12.35 ±0.16 mmHg,  
191 15.67 ±0.05mmHg which occurred at 60 minutes and they were statistically significant (p=0.000).  
192 The maximum mean difference in IOP and BP after oral administration of 0.6ml/kg body weight of  
193 fresh raw quail egg was similar in the three study groups. This showed that 0.6ml/kg body weight  
194 of fresh raw quail egg has similar hypotensive effects on IOP and BP of ocular and systemic  
195 hypertensives as well as the normotensive control subjects.

#### 196 4. DISCUSSION

197 The mechanism of action of the hypotensive effects of fresh raw quail egg on intraocular  
198 pressure and blood pressure has not been fully understood, but numerous possibilities have been  
199 proposed. Many of the proposals have attempted to relate the effect of dietary calcium on vascular  
200 smooth muscle of blood vessels and how it alters vascular tone to reduce blood pressure [21].  
201 Potassium supplements have also been shown to have blood pressure-lowering effect in persons  
202 with low dietary intake. The study of Frank *et al.*, (1998) [22] and George *et al.*, (2013) [10] further  
203 showed the importance of potassium in lowering blood pressure in the general population.  
204 Therefore the high potassium and calcium content in the quail egg may have played a role in  
205 lowering blood pressure. Hoshi *et al.*, (2013) [23] in their study concluded that Omega-3 fatty acids  
206 also lower blood pressure by directly activating large-conductance calcium-dependent potassium  
207 channels. Hence, Omega-3 fatty acids present in quail egg may have facilitated this mechanism to  
208 decrease the blood pressure.

209 Finally, as long as the aqueous humour drains into the episcleral venous system either by  
210 way of the intrascleral plexus or more directly along the aqueous veins, intraocular pressure varies  
211 directly with the venous pressure [24]. Decrease in episcleral venous pressure will therefore result  
212 in an increased outflow of aqueous causing a fall in intraocular pressure. Previous studies have  
213 shown that intraocular pressure is significantly correlated with systemic blood pressure. There are  
214 significant direct correlations between changes in systemic blood pressure and changes in  
215 intraocular pressure in humans and animals studies [7-12]. Therefore the fall in blood pressure may  
216 have consequently resulted in a fall in intraocular pressure.

#### 217 5. CONCLUSION

218 0.6ml/kg body weight of raw quail egg has significant hypotensive effect on intraocular  
219 pressure and blood pressure compared to water, when administered orally on empty stomach in  
220 ocular hypertensives, systemic hypertensives and in the normotensive control groups. The  
221 hypotensive effects in the three study groups are similar therefore it may be consumed as a form of  
222 supplement in the control of high blood pressure and raised intraocular pressure since the quail egg  
223 is innocuous. Toxicological studies including acute and repeated oral administration on rats as well  
224 as in vitro studies demonstrated good tolerability of the product without mutagenic or genotoxic  
225 effects [18]. Nevertheless further investigations may be required to determine the safety dose of the  
226 quail egg as a supplement in the management of ocular and systemic hypertension.

## 227 **CONFLICT OF INTEREST**

228 Authors have declared that there is no conflict of interest associated with this work.

229

230

231

## 232 **REFERENCES**

- 233 1. Iwuagwu OF, Megwas UA, Irem-Azode A. A study on the gender difference in the prevalence of  
234 glaucoma. J Afr Council Optom. 2007; 1: 104-110.  
235
- 236 2. Viera GM, Oliveira HB, de Andrade DT, Bottaro M, Ritch R. Intraocular Pressure Variation  
237 During Weight Lifting. Arch Ophthalmol. 2006; 124(9): 1251–1254.
- 238 3. Ajayi OB, George GO. Long term effect of exercise on intraocular pressure. JNOA. 2004; 37-  
239 39.
- 240 4. Kanski JJ. Glaucoma. Clinical Ophthalmology: A Systemic Approach, 5<sup>th</sup> ed, Butterworth-  
241 Heinemann, Elsevier Science Limited; 2003. p. 194-198.  
242

- 243 5. Grieshaber MC, Schoetzau A, Zawinka C, Flammer J, Orgul S. Effect of Central Corneal  
244 Thickness on Dynamic Contour Tonometry and Goldmann Applanation Tonometry in Primary  
245 Open-angle Glaucoma. *Arch Ophthalmol*. 2007; 125(6): 740–744.  
246
- 247 6. Tanaka GH. Corneal pachymetry: a prerequisite for applanation tonometry? *Arch Ophthalmol*.  
248 1998; 116(4): 544–545.  
249
- 250 7. Klein BE, Klein R, Knudtson MD. Intraocular pressure and systemic blood pressure:  
251 longitudinal perspective: the Beaver Dam Eye Study. *Br J Ophthalmol*. 2005; 89: 284-287.  
252
- 253 8. Vaajanen A, Mervaala E, Oksala O, Vapaatalo H. Is there a relationship between blood pressure  
254 and intraocular pressure? An experimental study in hypertensive rats. *Curr Eye Res*. 2008; 33:  
255 325-332.  
256
- 257 9. Castejon H, Chiquet C, Savy O, Baguet JP, Khayi H, Tamisier R, Bourdon L, Romanet JP.  
258 Effect of acute increase in blood pressure on intraocular pressure in pigs and humans. *Invest*  
259 *Ophthalmol Vis Sci*. 2010; 51:1599-1605.
- 260 10. George GO, Idu FK, Obika LFO. Effects of corn silk aqueous extract on intraocular pressure of  
261 ocular hypertensive human subjects. *S Afr Optom*. 2013; 72(3): 133-143.
- 262 11. George GO, Obika LFO. Effects of aqueous extract of corn silk on intraocular and blood  
263 pressures of normotensive subjects. *West Afr. J. Pharmacol. Drug Res*. 2014; 29: 16-24.
- 264 12. George GO, Idu FK. Corn silk aqueous extracts and intraocular pressure of systemic and non-  
265 systemic hypertensive subjects. *Clin Exp Optom*. 2015; 98: 138–149.
- 266 13. Sembulingam K, Sembulingam P. Intraocular pressure and Glaucoma. *Essentials of Medical*  
267 *Physiology*, 5th ed. Jaypee Brothers Medical Publishers; 2010. p. 581-591.  
268
- 269 14. Guyton AC, Hall JE. Fluid system of the eye - intraocular fluid. *Textbook of Medical*  
270 *Physiology*, 10th ed. Publishers: Elsevier-Saunders; 2000. p. 575-577.
- 271 15. Adeloye D, Basquill C, Aderemi AV, Thompson JY, Obi FA. An estimate of the prevalence of  
272 hypertension in Nigeria: a systematic review and meta-analysis. *J Hypertens*. 2015; 33(2): 230-  
273 242.
- 274 16. Tunsaringkarn T, Wanna T, Wattasit S. Nutrient Benefits of Quail (*Coturnix Coturnix*  
275 *Japonica*) Eggs. *Intl. J. of Sci and Res. Pub*. 2013; 3(5): 1-8.  
276
- 277 17. Lalwani P. Quail Egg Nutrition. 2011 [Accessed 2016 Mar 28] Available from  
278 <http://www.buzzle.com/articles/quail-egg-nutrition.html>.  
279

- 280 18. Bruttman G. "Ovix" Quail egg homogenate: a clinical evaluation. *La Medicina Biologica*.  
281 1995; 2: 25–29.  
282
- 283 19. British medical researchers for quail egg: Quail Bird Eggs and their Nutritional value. [Cited  
284 2013 Mar 19] Available from <http://www.quailbirdfarmnigeria.blogspot.com/2013/03/>.  
285
- 286 20. Aba PE, Igwebuike DC, Onah JA. Effects of Various Concentrations of Quail Egg Solution on  
287 Glycemia and Antioxidant Parameters of Alloxan–induced Diabetic Rats. *J. Adv. Med. Pharm.*  
288 *Sci.* 2015; 5(4): 1-7.
- 289 21. Hatton DC, Yue Q, McCarron DA. Mechanisms of calcium's effects on blood pressure. *Semin*  
290 *Nephrol.* 1995; 15(6): 593-602.
- 291 22. Frank MS, Walter CW, Smith A, Brown LE, Bernard R, Moore TJ. Effect on blood pressure of  
292 potassium, calcium, and magnesium in women with low habitual intake. *Hypertension*. 1998;  
293 31: 131-138.
- 294 23. Hoshi T, Wissuwa B, Tian Y, Tajima N, Xu R, Bauer M. Omega-3 fatty acids lower blood  
295 pressure by directly activating large-conductance  $\text{Ca}^{2+}$ -dependent  $\text{K}^{+}$  channels. *PNAS*. 2013;  
296 110(12): 4816-4821.
- 297 24. Tarkkanen A, Leikola J. Postural variation of IOP as measured by the Mackay-Marg Tonometer.  
298 *Acta Ophthalmol.* 1967; 45(4): 569-575.
- 299
- 300