

IN-VITRO ANTIBACTERIAL ACTIVITY AND PHYTOCHEMICAL SCREENING OF *PSIDIUM GUAJAVA* ON SOME ENTERIC BACTERIAL ISOLATES OF PUBLIC HEALTH IMPORTANCE

ABSTRACT

Aim: The study was conducted to investigate the antibacterial activity of *Psidium guajava* and its major antibacterial constituents (Phytochemicals).

Material and method: The aqueous and ethanol extracts from the leaves and stem bark of the plant was tested using well diffusion method for their antibacterial activity against some members Enterobacteriaceae family isolated from diarrheic stool sample (*Escherichia coli*, *Shigella* spp, *Salmonella typhi* and *Pseudomonas aeruginosa*).

Results: The result shows that the extracts were active against the microorganisms. The ethanolic extract of stem bark showed the highest zones of inhibition against tested organisms compared to aqueous extract. Statistical analysis of the result shows that the extracts demonstrated higher antibacterial activity against the isolates tested with the average zone of inhibition of 15.44mm, 14.78mm, 12.92mm and 11.31mm for *E. coli*, *Shigella* spp *S. typhi* and *P. aeruginosa*, respectively. The Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the extracts ranges between 6.25 – 100 mg/ml. Preliminary phytochemical analyses showed that both stem bark and leaf extracts contain alkaloids, tannins, terpenoid, Anthraquinone, reducing sugar, amino acid, flavonoid, saponins, glycosides and phenols.

Conclusion: The extracts of the plant demonstrated antibacterial activity due to presence of phytochemical constituents hence, the application of the decoction of leaf and stem bark of the plants in ethno medicine is justified.

Key words: *Psidium guajava*; Enteric bacteria; Phytochemical; well diffusion; antibacterial activity; extracts;

1. INTRODUCTION

There has been a lot of attention focused on producing medicines and products that are natural. Several leaves and leaves extracts have been found to have antibacterial activity against microorganisms [1]. There is no plant that does not have medicinal value [2]. The active components are normally extracted from all plant structure, but the concentration of these compounds varies from

30 structure to structure. However, plants parts known to contain the highest concentration of these
31 phytochemical constituents for therapeutic purpose can either be leaves, stem bark barks, root, bulks,
32 corms, rhizomes, wood, flowers, fruits or the seeds [3].

33 The presence of phytochemical constituents in medicinal plants made them useful for healing as well
34 as for curing of human diseases [4]. Phytochemical are naturally occurring compounds in the
35 medicinal plants, [5]. Large populations of the world, especially in developing countries depend on the
36 traditional system bark of medicine to treat variety of diseases [6]. Several hundred genera of plants
37 were utilized traditionally for medicinal purposes. The world health organization [7] reported that 80%
38 of the world population relies chiefly on traditional medicine and a major part of the traditional
39 therapies which involve the use of plant extract and their constituents [8].

40 Guava, *P. guajava*, belongs to the family Myrtaceae which is considered to have originated from
41 tropical South America, Guava tree grown in tropical and sub-tropical area of the world like Asia,
42 Africa and Hawaii etc [9]. *Psidium guajava* common names are Guava (English), Gwaiba (Hausa),
43 Goifa (Yoruba), Ugova (Igbo), Gutiba (Spanish), Goyave (French) and Goejaaba (Dutch) [10]. Many
44 parts of the plant have been used in traditional medicine to manage conditions like malaria,
45 Gastroenteritis, vomiting, diarrhea, dysentery, wound, ulcer, toothache, cough, sore throat and a
46 number of other conditions [11]. The leaves are particularly rich in flavonoids and distinctly in
47 quercetin [12]. The leaf methanolic extract also showed in vitro antibacterial activity against *E. coli*, *S.*
48 *typhi*, *Staphylococcus aureus* and a strong antibacterial action have been reported on Gram positive
49 and gram negative organism [11]. Therefore, the basic phytochemical investigation of its extract for
50 major phytoconstituents is also vital. The phytochemical analyses of Guava plant shows that its
51 extracts contain over twenty compounds [13].

52 The objective of the study was to determine the antibacterial activity of leaves and stem bark of *P.*
53 *guajava* extracts (Both aqueous and ethanolic) against some members of Enterobacteriaceae namely
54 *E.coli*, *Shigella* spp, *P. aeruginosa* and *S. typhi*, to screen for the phytochemical constituents of the
55 extracts and to evaluate the minimum inhibitory concentration (MIC) and minimum bactericidal
56 concentration (MBC) of the extracts against the test isolates.

57 **2. MATERIALS AND METHODS**

58 **2.1 Plant materials**

The plant materials used in this research consisted of the leaves and stem bark of *P. guajava* which were collected from botanical garden of Government Secondary school Gundutse, Kura Local Government Area of Kano State at about 08:30 a.m. Identification and Authentification of the plant materials has been done at Herbarium unit in the department of Biological science Ahmadu Bello University, Zaria with the following voucher numbers 3253. Voucher specimen has been deposited there for future reference. The samples were washed with water to remove dust and rinsed with distilled water. Samples were air dried for two-weeks and pulverized into powder form using sterile mortar and pestle in the laboratory as described by Mukhtar and Tukur [14]. The powdered samples were bagged in a black polythene bag and store in air tight container for further work.

2.2 Test organisms

Clinical isolates of *E. coli*, *Shigella* spp, *P. aeruginosa* and *S. typhi* were obtained from Department of Microbiology of Murtala Muhammad Specialist hospital for further experiment. Identification and characterization of the isolates was conducted there by using three procedures namely Gram staining, cultural characterization using selective or indicative media and biochemical characterization with reference to Cheesbrough [24]. The pure isolates of each of the test organism were inoculated in sterile slants containing Nutrient agar and transported to the department of Microbiology KUST and refrigerated at 4 °C before use.

2.3 Preparation of Plant Extracts

The ethanol and aqueous extract of both plant samples was carried out according to Bengum [13]. Twenty five gram (25g) each of fine powder of the plant materials was dissolved in 250ml of distilled water and ethanol in a conical flask and kept for 7 days in a cabinet and frequently shaken to dissolve the powder properly. After 7 days the solution of the plant materials was filtered using Whatman filter paper. The filtrate was kept in rotary evaporator for complete evaporation of the solvent. After running this procedure, a gummy extraction was obtained which was preserved in refrigerator before use

2.4 Antibacterial Assay of Extracts

The mixture was filtered using Whatman No.1 filter paper and the extracts were evaporated to dryness using rotary evaporator and water bath. One gram solid residues obtained were reconstituted in 5ml of 5% DMSO to form stock concentration of 200mg/ml, stored in the refrigerator at 4 °C until used. The agar well method was used to determine the antibacterial activity of the plant extracts. 0.1ml of the different standardized organisms were introduced separately and thoroughly mixed with

Mueller Hilton Agar in a sterile Petri dish and allowed to set then labeled. A sterile cork borer 6mm was then used to punch holes (i.e. 5 wells) in the inoculated agar and the agar was then removed. Four wells that were formed were filled with different concentrations of the extract which were labeled accordingly; 200mg/ml, 150 mg/ml, 100mg/ml and 50 mg/ml while the 5th well contained the solution used for the research to serve as control, tetracycline (Chi pharmaceutical limited, Lagos Nigeria) 125mg/ml, was used as control in this research. These were then left on the bench for 1hour for adequate diffusion of the extracts and incubated at 37 °C for 24 hours. After incubation, the diameter of the zones of inhibition around each well were measured to the nearest millimeters along straight line and the mean of the readings were then calculated [2].

2.5 Determination of Minimum Inhibitory Concentration (MIC) and Minimum

Bactericidal Concentration (MBC)

Overnight Nutrient broth cultures of *E. coli*, *Shigella spp*, *S. typhi* and *P. aeruginosa* at 37 °C were prepared. The culture was adjusted to obtain turbidity comparable to that of the turbidity of McFarland 0.5 standard, and then further dilute in Nutrient broth. The inoculums thus prepared expected to obtain 10⁵ cfu/ml. The MIC and MBC were determined using procedure described by Kowser and Fatima [15].

2.6 Phytochemical Screening

This was done on different extract to ascertain the presence of bioactive component present in the leaves and stem bark of *P. guajava*. The presence of Alkaloid, saponin, Glycoside, Tannin, flavonoids, steroid, terpenoid, Anthraquinones, Protein and amino acid were determined using procedure described by Sofowora [16].

3. RESULTS AND DISCUSSION

3.1 Antibacterial activity

The antibacterial activity of aqueous and ethanolic leaf extract *P. guajava* was represented in the table below (Table 1) showing the mean diameter of zone of inhibition of extract on the test isolate with *Shigella spp* being the most susceptible isolate at 200 mg/ml concentration (23.3mm) and *P. aeruginosa* the least susceptible (6 mm).

Table 1: Mean diameter zones of inhibition of Aqueous and ethanolic leaf extract *P. guajava*

CONCENTRATION (mg/ml)	MICROORGANISMS/ ZONE OF INHIBITION (mm)			
	<i>E. coli</i>	<i>Shigella spp</i>	<i>P. aeruginosa</i>	<i>S. typhi</i>

	50	09.6	10.6	06.0*	07.3
ALE	100	11.3	11.6	11.0	10.6
	150	11.6	12.3	10.3	18.0
	200	13.6	16.3	12.6	18.3
CONTROL		18.0	18.6	17.3	17.6
	50	11.6	11.3	07.6	08.3
ELE	100	15.3	18.3	14.3	10.6
	150	17.3	20.3	15.6	13.3
	200	20.0	23.3	17.0	18.6

117 ALE = Aqueous Leaf Extract. ELE = Ethanolic Leaf Extract. * = No Inhibition

118 *E. coli* = Escherichia coli. *P. aeruginosa* = *Pseudomonas aeruginosa*. *S. typhi* = *Salmonella typhi*

119 The antibacterial activity of aqueous and ethanolic stem bark extract *P. guajava* was represented in
120 the table below (Table 2) showing the mean diameter of zone of inhibition of extract on the test isolate
121 with *Shigella* spp being the most susceptible isolate at 200 mg/ml concentration (23.6mm) the least
122 susceptible is *P. aeruginosa*, especially for ALE (11.0 at 100 mg/ml and 10.3 at 150, with no linear
123 increase, and 12.6 at 200 mg/ml)

124 **Table 2: Mean diameter zones of inhibition of Aqueous and ethanolic stem bark extract *P.***
125 ***guajava***

CONCENTRATION (mg/ml)		MICROORGANISMS/ ZONE OF INHIBITION (mm)			
		<i>E. coli</i>	<i>Shigella</i> spp	<i>P. aeruginosa</i>	<i>S. typhi</i>
	50	12.3	13.3	11.6	11.3
ASE	100	14.0	14.6	12.3	13.3
	150	18.3	20.0	14.6	17.3
	200	22.3	23.3	18.3	20.3
CONTROL		18.6	18.3	17.3	17.6
	50	14.3	14.6	12.0	13.3
ESE	100	15.3	17.3	13.3	16.0
	150	20.3	21.3	17.3	18.3
	200	23.3	23.6	20.3	21.3

126 ASE = Aqueous Stem bark Extract. ESE = Ethanolic Stem bark Extract. * = No Inhibition

127 *E. coli* = Escherichia coli. *P. aeruginosa* = *Pseudomonas aeruginosa*. *S. typhi* = *Salmonella typhi*

128 3.2 Minimum inhibitory concentration (MIC) and Minimum bactericidal conc. (MBC)

129 Table below (Table 3) shows minimum inhibitory concentration (MIC) of the plant extracts. The result
130 shows that both aqueous and ethanolic leaf extract of the plant can inhibit the growth of the test
131 isolates at concentration of 6.25 - 25mg/ml. The aqueous stem bark extract has higher MIC of

50mg/ml in *E. coli* and *Shigella spp* than in *S. typhi* and *P. aeruginosa* but MIC of 12.5-50mg/ml for ethanolic extract. This shows that ethanolic extract is more effective against the test isolates.

Table 3: Minimum inhibitory concentration (MIC) and Minimum bactericidal concentration (MBC) of the *P.guajava* extracts against test isolates

EXTRACTS	ORGANISMS/ MIC/MBC (mg/ml)			
	<i>E. coli</i>	<i>Shigella spp</i>	<i>P. aeruginosa</i>	<i>S. typhi</i>
ALE	25/25	25/25	50/50	50/50
ELE	12.5/25	12.5/25	25/50	25/25
ASE	25/50	12.5/50	25/ND	12.5/ND
ESE	6.25/25	6.25/25	12.5/50	6.25/25

ALE = Aqueous leaf extract ELE = Ethanolic leaf extract ASE = Aqueous stem bark extract

ESE = Ethanolic stem bark extract ND = Not detected.

E. coli = *Escherichia coli*. *P. aeruginosa* = *Pseudomonas aeruginosa*. *S. typhi* = *Salmonella typhi*

3.3 Phytochemical screening

The Table below (Table 4) showed that the phytochemicals were present in both leaf and stem bark of *P. guajava*. Both leaf and stem bark contain Alkaloid, saponin, phenol, flavonoids, protein and amino acid, Anthraquinones, terpenoid and tannin except Steroid

Table 4: Phytochemical constituents of leaves and stem bark of *P.guajava*

S/N	PHYTOCHEMICAL	LEAF EXTRACT	STEM BARK EXTRACT
1.	Alkaloid	+	+
2.	Saponin	+	+
3.	Phenol	+	+
4.	Flavonoid	+	+
5.	Protein and amino acid	+	+
6.	Tannin	+	+
7.	Reducing sugar	+	+
8.	Anthraquinone	+	+
9.	Steroid	—	—

154	10.	Terpenoid	+	+
155	+ = Presence of phytochemical,		- = Absence of phytochemical.	
156				

157 The antibacterial activity is higher at a concentration of 200mg/ml against *Shigella spp* and
158 *Escherichia coli* as compared to *Pseudomonas aeruginosa* and *Salmonella typhi*. *Psidium guajava*
159 ethanolic extract showed higher antibacterial activity compared to aqueous extract, this is attributed to
160 better solubility of the active component by the ethanol than water. This shows similarities to the
161 findings of Nwanneka *et al.* [17] which investigated the antibacterial activity of *Psidium guajava* leaf
162 extract, the results showed that both aqueous and ethanolic extracts of guava leaf inhibited the
163 growth of the bacteria and fungi tested but the ethanolic extract showed stronger inhibition than the
164 aqueous extract against the organisms. This result was also in conformity with that of Pandey and
165 Shweta [18] where the results of antibacterial activity of *Psidium guajava* leaf and stem bark reveals
166 that ethanolic extract showed stronger anti-bacterial activity than aqueous extract. However, this was
167 contrary to the findings of Emmanuel, [19] who investigate the antibacterial effects of some Ghanian
168 aromatic plants including *Psidium guajava*, he stated that the antibacterial activity of aqueous extract
169 of *Psidium guajava* is higher than that of ethanolic extract. This is due to differences in geographical
170 location of the plant he used compared to the one of the present study. The result of antibacterial
171 effects of *Psidium guajava* leaf extract by Elekwa *et al.* [20] showed that aqueous extract of guava
172 leaf had higher inhibitory effects on some organisms than ethanolic extract. The results of the present
173 research on the antibacterial activity of stem bark and leaf of *Psidium guajava* on the tested isolates
174 revealed that stem bark extract posses higher antibacterial activity than corresponding leaf extract.
175 This hold true with the results of Elekwa *et al.* [20] in which the antibacterial study on the effect of
176 *Psidium guajava* showed that the stem bark extract is more effective than leaf extract. The higher
177 antibacterial activity shown by stem bark extract is as result of having higher quantity of the
178 phytochemicals in stem bark extract compared to leaves extract.

179 From the results of MIC determination (Table 3), the minimum inhibitory concentration showed that a
180 very low concentration of 6.25 -12.5mg/ml the ethanolic stem bark extract of *Psidium guajava* inhibit
181 the growth of all tested isolates.

182 Preliminary Phytochemical Analysis of *P. guajava* In the present study was carried out to identify the
183 active constituents such as alkaloids, flavonoids, sterols, terpenoid, Anthraquinones, protein and
184 amino acid, phenol, carbohydrate and cellulose present in the leaves and stem bark of guava plant.

Preliminary phytochemical analysis of leaf and stem bark extract, (ethanol and water) of *P. guajava* showed the presence of the following chemical constituents; Alkaloid, saponin, phenol, flavonoids, protein and amino acid, Anthraquinones, terpenoid and tannin (Table 4). Earlier work have revealed the presence of alkaloids, flavonoids, glycosides, poly-phenols, reducing compounds, saponins and tannins in the aqueous extract of *P.guajava* leaf [21]. This finding can be attested to the work of Offo [22] who also reported similar finding on the phytochemical of guava leaf extract, which contain alkaloid, saponin, flavonoids, phenol, steroid, tannin, protein and glycoside. Pandey and Shweta, [18] reported the phytochemicals mainly present in *Psidium guajava* were reducing sugar, tannin, saponin, phlobatannin, terpenoid, alkaloid and phenols. The finding is also similar to that of Joseph and Priya, [23] where the preliminary phytochemical analysis of leaf, stem bark and root extracts of *Psidium guajava* showed the presence of carbohydrate, glycoside, saponin, Anthraquinone, Flavonoid, tannin and alkaloids at high concentration.

4. CONCLUSION

The antibacterial activity of the plant showed that the plant part extracts demonstrated antimicrobial effect against the test isolates with higher activity in stem bark extract compared to leaves extract. Higher antibacterial activity was recorded in ethanol extract compared to aqueous extract. The antibacterial activity of the plant parts is due to the present of the Phytochemicals identified in this study. The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of the extracts ranges from 6.25 – 50mg/ml of the extracts. The Phytochemical screening of the plant parts revealed the present of Alkaloid, Tannin, Saponin, and Cardiac glycoside, protein and amino acid, Flavonoid, Terpenoid, Phenols and Anthraquinone. Findings from this work support the use of extracts from *P. guajava* stem bark and leaves as medicinal plant. It is recommended that Government, Non-governmental organizations (NGOs) and philanthropists should encourage researchers in this field so that the spread of antibacterial resistant pathogens can be restrained.

ETHICAL APPROVAL

Ethical approval (issue number HMB/GEN/488/VOL. 1) was obtained from Kano State Hospital Management Board based on the consent of Murtala Muhammad Specialist Hospital ethical committee

REFERENCES

- [1] Kim, S. and Fung, Y. V. Antibacterial effect of crude water soluble Arrow root (*Peurariae radix*) tea extracts on food borne pathogens in a liquid medium. "Letters in Applied Microbiology" 2004; 39 (4): 319 – 325.
- [2] Anibijuwon, I.I. and Udeze, O.A. Antibacterial Activity of *Carica Papaya* (Paw-paw Leaf) on Some Pathogenic Organisms of Clinical Origin from South-Western, Nigeria. *Ethno botanical Leaflets*. 2009. 13: 850-864.
- [3] Kafaru, E. Immense help formative workshop. In: Khan, R., Islam, B., Akram M., Shakil, S., Ahmad, A., Ali, M.S., Sadiqui, M. and Khan, A.U. (2008). Antibacterial Activity of Five Herbal Extracts against Multi Drug Resistant (MDR) strains of Bacteria and Fungus of Clinical Origin. *Molecules* 1994. 13
- [4] Nostro, A., Germano, M. P., D'angelo, V., Mariano, A. and Lanattel M. A. Extraction method and bio-autography for evaluation of medicinal plants antibacterial activity. *Letter in Applied Microbiology* 2000. 30: 379.
- [5] Abdul Wadood, Ghufuran, M., Babar Jamal, S. B., Naeem, M., Khan, A., Ghaffar, R. and Asnad A. Phytochemical Analysis of Medicinal Plants Occurring in Local Area of Mardan. *Biochemistry & Analytical Biochemistry*, 2013. 2: 144. DOI 10.4172/2161-1009.1000144
- [6] McGaw, L. J., Jager, A. K. and Staden, J. V. Antibacterial, anti-helminthes and anti amoebic activity in South Africa medicinal plants. *Journal of Ethnopharmacol.* 2000; 72: 247 – 263
- [7] World Health Organization (WHO). 2004 Use of antibacterials outside human medicine and result and antibacterial resistance in humans. World Health Organization 2002. Archived from the Original on 13 May, 2004
- [8] Ahmed, I. and Beg, A.Z. Antibacterial and phytochemical studies on 45 Indian Medicinal plants against multi-drug resistance human pathogens. *Journal of Ethnopharmacol.* 2003; 74: 113-123
- [9] Biswal, B., Kimberly, Fredrick, M., Dwayne, D. and Anand, Y. Antibacterial activity of leaf extracts of Guava (*P.guajava*) on two Gram positive and two Gram negative bacteria. *International Journal of Microbiology* 2013; 2: 7
- [10] Burkill, H.M. The Useful Plants of Tropical Africa, Vol. 1, 2nd edition, *Royal Botanical Garden*: 1985. Kew pp 389.
- [11] Abdelrahim, S. I., Almagboul, A.Z., Omer, M. E. and Elegami, A. Antibacterial activity of *P.guajava* L. *Fitoterapia* 2002. 73:713-715.
- [12] Abok, A. Spectrophotometric estimate of Anthraquinones content and antibacterial potential of extracts of some *Cassia specie* used in herbal medicine in Ibadan, Nigeria. *Science forum*, 2000. Vol.1 pp53 – 57

248 [13] Begum, M. The phytochemical and pharmacological investigation of *Carica papaya* leaf. An
 249 undergraduate thesis. East west, University 2014

250 [14] Mukhtar, M. D. and Tukur A. In-vitro screening activity of *Pistia stratiutes* extract. *NISED Journal*
 251 1999. 1 (1) 5 – 6

252 [15] Kowser, M. M. & Fatema, N. Determination of MIC and MBC of selected azithromycin capsule
 253 commercially available in Bagladesh. *The Orion Medical journal* 2009. 32 (1): 619 – 620.

254 [16] Sofowora, A. Research on Medicinal Plants and Traditional Medicine in Africa. *Journal of*
 255 *Alternative and Complementary Medicine*, 1996. 2(3):365 – 372.

256 [17] Nwanneka Lauretta Ofodile, Ndubuisi Moses Chikere Nwakanma, Michael Mordi, and Oluwakemi
 257 Ademolu. Genotoxic and antibacterial studies of the leaves of *P.guajava*. *Euro-Asian Journal of Bio-*
 258 *Sciences*. 2013; 7: 60-68

259 [18] Pandey, A. and Shweta. Antifungal properties of *Psidium guajava* leaves and fruits and fruits
 260 against various pathogens. *Journal of Pharmaceutical and Biomedical Science* 2011. 13 (13): 1-6

261 [19] Emmanuel, O. A. Antibacterial activity profile of the constituents of Ghanaian aromatic medicinal
 262 plants. An M. Sc thesis, Kwame Nkuruma University, Ghana.2010.

263 [20] Elekwa, I. Okereke, S. C. and Ekpo, B. O. Preliminary phytochemical and antibacterial
 264 investigations of the stem bark bark and leaves of *P.guajava* L. *Journal of Medicinal Plants Research*.
 265 2009; 3(1): 45 - 48.

266 [21] Uboh, F.E., Okon, I.E. and Ekong, M.B. Effect of aqueous extract of *Psidium guajava* leaves on
 267 liver enzymes, histological integrity and hematological indices in rats. *Gastroenterol*. 2010; 3(1): 32 –
 268 38

269 [22] Offor, C.E. Phytochemical and Proximate Analyses of *P.Guajava* Leaves, *Quest Journals Journal*
 270 *of Research in Pharmaceutical Science*. 2015; (2): 05- 07

271 [23] Joseph, B. and Priya R. M. Review of nutritional, medicinal and pharmacological properties of
 272 Guava (*P.guajava* L.) *International Journal of pharma Bio Science*. 2011; 2 (1): 53 – 69.

273 [24] Cheesbrough, M. *District Laboratory Practice in Tropical Countries*, Cambridge United Press,
 274 U.K. part. 2000; 27: 105.