

Evaluation of Phytochemical and *in vitro* Antimicrobial Effects of *Solanum lycopersicum* Linn. (Tomato) on Oral Thrush and Human Cariogenic Pathogens

ABSTRACT

Aims: Study on the evaluation of phytochemical and antimicrobial effects of tomato *Solanum lycopersicum* (L.) on oral thrush and human cariogenic pathogens was designed to determine the antimicrobial effect of tomato extracts, and to identify the common phytochemical constituents of tomato that may be inhibitory to oral pathogens.

Place and Duration of Study: Fresh tomato fruit samples were collected from Samaru market, Zaria in the month of November and December 2015, and analyzed at Microbiology Laboratory, Nigerian Institute of Leather and Science Technology Zaria, Kaduna.

Methodology: Cold maceration extraction using methanol and water was adopted. All the extracts were subjected to standard phytochemical qualitative screening for the presence or absence of various primary and secondary metabolites. Antimicrobial susceptibility was carried out by agar well diffusion technique. The antimicrobial susceptibility test of *Solanum lycopersicum* (tomato) fruit extracts against selected bacterial (*Streptococcus mutans*, *Bacillus subtilis*) and yeast (*Candida albicans*) pathogens capable of causing oral thrush and dental caries was carried out. Ciprofloxacin and Econazole were used as positive controls. The Minimum Inhibitory Concentrations (MICs) were determined in three concentrations; 100 mg/ml, 50 mg/ml and 25 mg/ml of each extract. Mean zone of inhibition was used to measure the antimicrobial potential of tomato fruit extracts against the test organisms. The microbial isolates were obtained from the Department of Medical Microbiology, Ahmadu Bello University, Zaria.

Results: Alkaloids, flavonoids, glycosides, saponins, tannins, steroids, phlobatannins, terpenoids and tannins are present in tomato fruit extract, but, anthraquinones and phlobatannins are absent in methanolic extract. While, the only phytochemical that was not detected in the aqueous extract is anthraquinones. Highest antibacterial activity was recorded on *Bacillus subtilis* at the concentration of 100 mg/ml and 50 mg/ml for aqueous and methanolic extracts respectively. *Candida albicans* and *Streptococcus mutans* showed resistance to the various extracts at various concentrations used. The MIC of 100 mg/ml was recorded on *Bacillus subtilis*, whereas all other test organisms showed relative resistance to various concentrations of the extracts used.

Conclusion: Tomato fruit contains phytochemicals that showed promising antimicrobial effect on oral thrush and cariogenic pathogens. The tomato extract recorded antimicrobial effect against *Bacillus subtilis* with MIC of 100 mg/ml. The methanolic extract was the most active that could compete favourably with the conventional antibiotic (ciprofloxacin) at higher concentrations, which suggests that possible new drug candidates can be harnessed from the tomato fruit against oral thrush and cariogenic pathogens.

Keywords: extraction; lycopene; minimum inhibitory concentration; phytochemistry; *solanum lycopersicum*

1. INTRODUCTION

Tomato, *Solanum lycopersicum* (Linn), previously known as *Lycopersicon esculentum* (Mill) is a red berry fruit [1]. It belongs to the plant family solanaceae and genus solanum [2]. Tomato is one of the most important vegetables worldwide because of its high consumption, year round availability and large content of health related components [3]. It is used as juice, soup, ketchup or paste [4]. Although tomatoes are often closely associated with Italian cuisine, they are actually originally native to the Western side of South America in the region occupied by Columbia, Ecuador, Peru, Chile and the western half of Bolivia. Tomato is very rich in vitamins, minerals, essential amino acids, sugars, dietary fibers and lycopene, an antioxidant that reduce the danger related to many cancers and neurodegenerative diseases [5]. Today tomatoes are enjoyed worldwide to the tune of 130 million tons per year by the largest tomato producing country, which is China (with approximately 34 million tons of production) followed by United States, Turkey and India [6]. The plant requires relatively cool weather for optimum yield [7], it typically grows up to 1 to 3 meters (3 – 10 feet) in height and have a weak stem that often sprawls over the ground. It is a perennial in its native habitat, and grown as an annual in temperate climates. The leaves are 10 – 25 cm long, odd pinnate, with five to nine leaflets on petioles; each leaflet is up to 8 cm long with a serrated margin [8].

Tomato contains variety of phytonutrients such as carotenoids (including beta-carotene, lutein, and zeaxanthin); flavonoids (including naringenin, chalconaringenin, rutin, kaempferol, and quercetin); hydroxycinnamic acids (including caffeic, ferulic, and coumaric acid); glycosides (including esculeoside A); and fatty acid derivatives (including 9-oxo-octadecadienoic acid). Lycopene, being the major carotenoid in tomato fruit, is the common antioxidant that reduces the danger related to many cancers and neurodegenerative diseases. It is a powerful antioxidant, anti-inflammatory and also has an antimicrobial property [5, 9, 10].

Some studies have shown that tomatoes have cancer prevention effect. The effect of tomato in the eradication of cancer is still unknown, but it was believed that lycopene and the newly discovered bioflavonoids in tomatoes are responsible cancer fighting agents [9]. Tomato is an excellent fruit vegetable for rapid skin cell replacement. Its juice can be used for healing sunburn because of its unique vitamin C. Tomato juice is a good sport drink used to restore one's body from fatigue and sleepiness [11].

The ever increasing demand for safer and cheaper herbal recipes in the developed countries has led to the extraction and development of several drugs and chemotherapeutic agents from plants as well as from traditional herbal remedies [12]. There are considerable economic benefits in the development of indigenous medicines and in the use of medicinal spices for the treatment of various diseases [13].

Majority of chemically synthesized drugs have serious adverse effect to the recipients, which may lead to temporary or permanent disability and incapacitations. Also, oral blisters, oral thrush, dental caries and oral ulcers are very serious infections that can lead to inability to take food or water as a result of dysphagia. The oral thrush disorder is a serious disease that can affect many people at various stages of their lives causing distress and discomfort. Sometimes, the disorder can lead to hospitalization as a result of starvation and dehydration following the victim's inability to ingest food or water [14, 15].

Direct topical application of raw tomato in combination with salt has been used for long in many African tribes to treat oral thrush, blisters, dental caries and ulcers of varying degrees. Therefore this research was designed to determine the antimicrobial effect of tomato, and to identify the common phytochemical constituents of tomato that may be inhibitory to oral pathogens. The research findings can further be used by the dental, pharmaceutical industries, complementary medicine and other medical sectors in tackling the menace of microbial oral disorders.

2. MATERIAL AND METHODS

2.1 SAMPLE COLLECTION

Fresh physically undamaged fruits of *Solanum lycopersicum* (tomato) were collected from Samaru market Zaria based on the guidelines of convention on Biological Diversity, and taxonomically characterized as Roma VFN, at the herbarium sections of Biological Science Department, Ahmadu Bello University, Zaria and National Horticultural Research Institute, Bagauda Station, Kano with the aid of botanical keys [16], ensuring that the tomato fruit is not among the endangered plant species. The tomato fruits were collected fresh, healthy and free from organic contaminants that may interfere with the substances of interest. The fruits were thoroughly washed through running water, cut and dried under shade for 4 - 16 days, after which the dried fruits were pounded into a fine powder using a sterile mortar and pestle [17]. The tomato fruit powder was stored and sealed in labeled sterile reagent bottles for further use. The bioactive components were extracted using the method of Akerele *et al.* [18].

2.2 PREPARATION OF PLANT EXTRACT

Plant extracts were prepared by the method of Alade and Irobi [19] with minor modifications. To study the antibacterial potential of *Solanum lycopersicum* (tomato), non-polar solvent such as methanol, and polar solvent such as water were used. A mass of 30 g of dried powder was weighed on a weighing balance (Mettler 166®, USA) and was extracted by 100 ml of each solvent until the aqueous content evaporated completely as adopted by Lin *et al.* [20]. At the end of extraction, the extracts were filtered through Whatman No. 1 filter paper and concentrated over a water bath using a rotary-vacuum evaporator to recover the solvents. The dried extracts were collected and weighed in varying concentrations and kept in airtight container at 4°C.

2.3 PRELIMINARY PHYTOCHEMICAL SCREENING

The preliminary phytochemical investigation was carried out for aqueous and methanolic extracts of *Solanum lycopersicum* for the detection of various phyto-constituents by using standard procedures to identify the constituents [20, 21].

2.3.1 Test for the presence of Alkaloids (Wagner's test)

Wagner's reagent was prepared by dissolving 2 g of iodine and 6 g of KI in 100 ml of water. The plant extract was prepared by using 500 mg of plant extract, which was treated with few drops of Wagner's reagent. A reddish brown coloured precipitate indicates the presence of alkaloids [22].

2.3.2 Test for the presence of Anthraquinone (Borntrager's test)

About 0.5 g of the extract was boiled with 10% HCl for few minutes in a water bath. It was filtered and allowed to cool. Equal volume of chloroform was added to the filtrate. Few drops of 10% ammonia was added to the mixture and heated. Formation of rose-pink colour indicated the presence of anthraquinones [23].

112 2.3.3 Test for the presence of Flavonoids (Ethyl acetate test)

113 The crude powder of dried plant was heated with 10 ml of ethyl acetate over a steam bath
114 for 3 minutes. The mixture was filtered and 4 ml of the filtrate was shaken with 1ml of dilute
115 ammonia solution, after which a yellow colouration was observed [24].

116 2.3.4 Test for the presence of Phlobatannins (HCl test)

117 An aqueous extract of each plant sample was boiled with 1% aqueous hydrochloric acid
118 (HCl) to observe the deposition of red precipitate [25].

119 2.3.5 Test for the presence of Glycosides (Fehling's test)

120 The crude plant powder of 0.5 g was dissolved in 5 ml of methanol. Exactly 2 ml of this
121 solution was dispensed into a test tube, and to it 10 ml of HCl was added. The mixture was
122 heated in a boiling water bath for 30 minutes, and to the mixture 5ml of Fehling's solution
123 was added and the mixture was boiled for another 5 minutes to observe a brick red
124 precipitate as an indication for the presence of glycosides [26].

125 2.3.6 Test for the presence of Saponins (Frothing test)

126 About 0.2 g of the extract was shaken with 5 ml of distilled water and then heated to boil.
127 Frothing (appearance of creamy layer of small bubbles) showed the presence of saponins
128 [24].

129 2.3.7 Test for the presence of Steroids (Salkowski test)

130 A volume of 1 ml of plant extract was dispensed into a test tube, and to it few drops of
131 concentrated sulphuric was added. The presence of red colouration indicated the presence
132 of steroids [27].

133 2.3.8 Test for the presence of Terpenoids

134
135 A mass of 0.2 g of the plant extract was mixed with 2 ml of chloroform and filtered. Acetic
136 anhydride was added to the filtrate, and 3 ml concentrated sulphuric acid (H₂SO₄) was
137 added carefully to form a layer. A pink coloration of the interface was formed to indicate
138 positive results [27]

139
140 **2.4 BACTERIAL PATHOGENS**

141 Characterized isolates of overnight cultures of oral thrush and dental caries-causing test
142 organisms such as *Candida albicans*, *Streptococcus mutans* and *Bacillus subtilis* were
143 collected from the Dental Section, Medical Microbiology Laboratory, Ahmadu Bello
144 University, Teaching Hospital, Zaria. The isolates were kept in fresh broths to maintain their
145 viability.

146

147 **3.4 ANTIMICROBIAL SUSCEPTIBILITY SCREENING**

148 Screening of the antibacterial activity of the extracts was carried out using agar-well diffusion
149 method. Nutrient agar (N70123, Sigma-Aldrich Co., Germany) for bacterial isolates and
150 Sabouraud's Dextrose agar (RODAC™, Becton-Dickson and Co., USA) for *Candida*

151 *albicans* were prepared according to manufacturer's specifications, sterilized, and poured
152 into Petri dishes and allowed to set. Wells were made in the inoculated media using sterile
153 cork-borer (6 mm diameter, NC-11269, Science Company®, New Jersey, USA) after which a
154 little molten media was used to seal the base of the wells. The characterized and 0.5 scale
155 McFarland standardized test organisms were inoculated on the media by the spread plate
156 method using a sterile rod spreader to obtain uniform microbial growth. A volume of 0.1 ml
157 each of the prepared extracts (100 mg/ml, 50 mg/ml and 25 mg/ml) were transferred into the
158 corresponding wells with a sterile micropipette and were appropriately labelled, while 0.1 ml
159 of 20% Dimethyl sulfoxide (DMSO) (DMS, Gaylord Chemical Co., Louisiana, USA) (free of
160 extract) was dispensed into a separate well to serve as the negative control. Ciprofloxacin
161 (30 µg) was used as the positive control for bacterial isolates, while Econazole (30 µg) was
162 used as positive control for *Candida albicans*. The plates were incubated for 48 hours at
163 37°C. Antibacterial and antifungal activities were evaluated by measuring the diameters of
164 zones of growth inhibition in millimeters (Plate 1) after 48 hours incubation. Oxoid [28]
165 standard susceptibility range was used to classify zones of inhibition as either sensitive (> 10
166 mm) or resistant (≤ 10 mm). Each experiment was conducted twice, and the mean of two
167 results was taken for both the test and control as adopted by Irobi *et al.* [29].

168 2.5 DETERMINATION OF MINIMUM INHIBITORY CONCENTRATION

169 The Minimum Inhibitory Concentration (MIC) was determined as the least concentration that
170 showed an inhibitory effect on test organism using the tube method. Two fold serial dilutions
171 were made using nutrient broth (N70123, Sigma-Aldrich® Co., Germany). The 5 ml of a
172 solution of the extract (250 mg/ml) was added aseptically to 5 ml of double strength medium
173 and mixed by shaking. Using a fresh pipette, 5 ml of the mixture was transferred to the
174 second test tube which contained 5 ml of the single strength medium. This too was mixed
175 by shaking and from it 5 ml will be taken into third test tube aseptically and mixed by
176 shaking. The 9th tube containing no test compound served as control. Finally, to each tube
177 0.2 ml inoculums of the test organisms were added aseptically. The test tubes were covered
178 with cotton wool and incubated at 37°C for 24 hours and then observed for turbidity. The
179 lowest concentration that inhibited growth of test organism was noted as the MIC [30].

181 3. RESULTS AND DISCUSSION

182
183 Table 1 showed the phytochemical screening of tomato fruit extract. Phytochemicals such as
184 alkaloids, flavonoids, phlabotannins, glycosides, saponins, tannins, steroids and terpenoids
185 were detected in aqueous extracts analyzed. While, anthraquinone was not detected in
186 either of the extracts analyzed. Whereas, all the phytochemicals tested were present in the
187 methanolic extract except anthraquinones and phlabotannins.

188 Table 2 showed the antimicrobial effects of tomato aqueous extract against bacterial
189 isolates. Highest antibacterial activity was recorded on *Bacillus subtilis* at the concentration
190 of 100 mg/ml. All other test organisms showed resistance to the various extracts at various
191 concentrations used.

192 Table 3 showed the antimicrobial effects of tomato methanol extract against bacterial
193 isolates. Highest antibacterial activity was recorded on *Bacillus subtilis* at the concentration
194 of 100 mg/ml and a slight sensitivity at 50 mg/ml. All other test organisms showed resistance
195 to the various extracts at various concentrations used.

196 Table 4 showed the minimum inhibitory concentration of the tomato fruit crude extract. The
197 MIC of 100 mg/ml was recorded on *Bacillus subtilis*, whereas all other tests organisms
198 showed relatively resistance to the extract, and hence their MICs were not determined.

Table 1. Qualitative phytochemical screening of tomato fruit extracts

Phytochemicals	Aqueous extract	Methanol extract
Alkaloids	+	+
Flavonoids	+	+
Phlabotannins	+	–
Glycosides	+	+
Saponins	+	+
Tannins	+	+
Terpenoids	+	+
Anthraquinones	–	–
Steroids	+	+

*+ = Present; – = Absent

Based on the phytochemical analysis findings of the research study, it was found out that phytochemicals such as alkaloids, flavonoids, glycosides, saponins, tannins, steroids, phlabotannins and terpenoids and tannins were present in both aqueous and methanolic extracts of tomato fruit which have been associated with antimicrobial activities [31, 32]. Anthraquinone is absent in aqueous extract, while anthraquinone and phlabotannins are absent in methanolic extract. This might be attributed to the difference in the method of extraction employed, polarity of solvents used and also pointing to the fact that the use of solvents during extraction of medicinal plant might have some advantages to individual application. Sukhdev *et al.* [33] also reported the effect of temperature on phyto-components due to the fact that some of these components are thermo-labile in nature. The result reveals the presence of alkaloids which are a large group of secondary chemical constituents of plants made largely of ammonia compounds comprising basically of nitrogen bases synthesized from amino acid building blocks with various radicals replacing one or more of the hydrogen atoms in the peptide ring, mostly containing oxygen and they are intensely bitter in solutions.

The results for the antimicrobial screening have shown that at the concentration of 100 mg/ml, all the extracts have antibacterial activity on *Bacillus subtilis*. Effectiveness factor of the tomato is due to the presence of Lycopene that possesses antibacterial and antifungal properties [34, 35]. The results of the inhibition of bacterial growth have shown that the extracts are active at high concentration and inactive at very low concentrations. Thus the study suggests that the inhibition of bacterial growth activity of the extracts is dose dependent. The aqueous and methanolic extracts had no inhibitory effect on the *Streptococcus mutans* and *Candida albicans* but showed inhibitory effects on the Gram-positive rod bacterium, *Bacillus subtilis*, with activity increasing with increase in concentration of extracts. This concurs with the findings of Omodamiro and Amechi [36] who reported antibacterial activity of tomato extract on some of pathogenic microbes. But, resistance of *Candida albicans* to both extracts is contrary to the findings of Sung *et al.* [37] and Pavlovic *et al.* [38] who reported that tomatoes proved to be effective against microorganisms such as *Staphylococcus aureus*, *Proteus*, *Bacillus* and antifungal to *Candida albicans* and *Aspergillus niger*.

235 **Table 2: Antibacterial Effect of Aqueous Tomato fruit Extract**

Test organism	Zones of growth inhibition						
	100 mg/ml	50 mg/ml	25 mg/ml	12.5 mg/ml	Ciprofloxacin	Econazole	DMSO
<i>Bacillus subtilis</i>	10 mm	0 mm	7 mm	0 mm	32 mm	–	0 mm
<i>Candida albicans</i>	0 mm	0 mm	0 mm	0 mm	–	36 mm	0 mm
<i>Streptococcus mutans</i>	0 mm	0 mm	0 mm	0 mm	35 mm	–	0 mm

236
237
238 **Table 3: Antibacterial Effect of Methanol Tomato fruit Extract**

Test organism	Zones of growth inhibition						
	100 mg/ml	50 mg/ml	25 mg/ml	12.5 mg/ml	Ciprofloxacin	Econazole	DMSO
<i>Bacillus subtilis</i>	25 mm	15 mm	0 mm	0 mm	32 mm	–	0 mm
<i>Candida albicans</i>	0 mm	0 mm	0 mm	0 mm	–	35 mm	0 mm
<i>Streptococcus mutans</i>	0 mm	0 mm	0 mm	0 mm	36 mm	–	0 mm

239 DMSO= Dimethyl sulfoxide



Plate 1: Antimicrobial susceptibility setup of tomato extract

Table 4: Minimum Inhibitory Concentration (MIC) of Tomato fruit Extracts

Test organisms	Concentration		
	Aqueous	Methanol	Control
<i>Bacillus subtilis</i>	100 mg/ml	50 mg/ml	None
<i>Candida albicans</i>	None	None	None
<i>Streptococcus mutans</i>	None	None	None

This may be due to the better solubility of the bioactive agents in the non-polar solvents. This conforms to the work of El-Mahmood *et al.* [39] who reported that the extractability of phyto-constituents differs from solvent to solvent depending on their polarities. It also reveals that the methanolic extract was more active against the Gram-positive bacteria. Since the traditional herbal remedy preparation use water as the extractant, it is a paradox that the aqueous extracts were less active in this study. It is possible that the aqueous extracts may contain antimicrobial constituents insufficient for efficacy in this study, which may explain why large amounts of the decoctions must be drunk by the patients. Jigna and Chanda [40] and Kitonde *et al.* [41] found out that, aqueous extracts showed little or no antimicrobial activity in contrast to those made using organic solvents. Yinegar *et al.* [42] reported that the success in traditional medicines may be due to administration of the extracts in large quantities and over a long period of time.

The Minimum Inhibitory Concentration (MIC) of the aqueous extract was highest for *Bacillus subtilis* at the concentration of 100 mg/ml with no activity in all other extract concentrations used. While, the MIC of the methanolic extract was least for *Bacillus subtilis* at the concentration of 50 mg/ml, suggesting that the methanol extract is more efficacious on *Bacillus subtilis* than the aqueous extracts. This is contrary to the studies of Omodamiro and Amechi [36] who reported MIC of 31.25 mg/ml for *Proteus mirabilis* and *Pseudomonas aeruginosa*. The antimicrobial effect of tomato in the treatment of oral microbial diseases can probably be enhanced when tomato is used in combination with table salt. The synergy between raw tomato and salt or honey might inhibit diverse types of oral pathogens.

Previous studies by Al-Oqaili *et al.* [43] reported correlation between synergistic combination of tomato and honey with high antimicrobial therapeutic effects.

4. CONCLUSION

The secondary metabolites detected in the tomato fruit extracts include alkaloids, flavonoids, glycosides, saponins, tannins, steroids, phlabotannins, terpenoids and tannins. Anthraquinone and phlabotannins are absent in methanolic extract. While, the only phytochemical that was not detected in the aqueous extract is anthraquinones.

Bacillus subtilis was susceptible to all the extracts at the concentrations of 100mg/ml, but *Streptococcus mutans* and *Candida albicans* were not susceptible to the aqueous and methanolic extracts within the concentration range for this study. The methanolic extract was the most active that could compete favourably with the conventional antibiotic (ciprofloxacin) at higher concentrations. The activity of the extracts on the test organisms increased with increase in concentration. The methanol extract exhibited bacteriostatic effects on the *Bacillus subtilis* at lower concentrations and bactericidal effects at higher concentrations, with minimum inhibitory concentration of 100mg/ml.

This research may serve as a scientific basis and lend credence to the claim by the traditional medicine practitioners that the tomato fruit is a useful herbal remedy to various microbial oral infections.

ETHICAL APPROVAL

Not applicable

REFERENCES

1. IAPT (2016). International Code of Nomenclature for algae, fungi and plants. International Association for Plant Taxonomy. Chapter V, section 3. Retrieved on 14th September 2016. Available at: www.iapt-taxon.org
2. Sarah, C. D., Sandra K & Iris E. P. (2003). Taxonomy of tomatoes in the Gal'apagos Islands: native and introduced species of *Solanum* section *Lycopersicon* (Solanaceae). *Journal of Systematics and Biodiversity*, 1 (1): 29–53.
3. Krishna, M. J., Bhaumik, A. and Kumar, S.P. (2013). Phytochemical Analysis and Antimicrobial Studies of Various Extracts of Tomato (*Solanum lycopersicum* L.). *Scholars Academic Journal of Biosciences (SAJB)*, 1(2):34-38.
4. Dilis B and Trichopoulou A. (2010). Antioxidant intakes and Food Sources in Greek Adults. *The Journal of Nutrition*. Bethesda, 140(7); 1274-1279.
5. Srinivasan, R. (Ed.). (2010). *Safer Tomato Production Methods: A field Guide for soil fertility and pest management*. AVRDC- The World Vegetable Center, Shanhua, Taiwan. AVRDC Publication No. 10-740. pp. 2.
6. Parnell, T., Suslow Trevor V., Lindaj, H. (2014). *Tomato Safe Method to store preserved and enjoy* (PDF) and catalogue University, California Division of Agriculture and Natural Resources Retrieved 18th February, 2015.
7. Rodriguez, G.R. (2007). Effect of Rice Bran Mulching on Growth and Yield of Cherry Tomato. *Cienciae Investigacion Agraria*, 34: 181-186.
8. Acquaah, G (2002). *Horticulture: Principles and Practices*. New Jersey. Prentice Hall. ISBN: 0130331252.
9. Etminan, M., Takkouche, B. and Caamaño-Isoma, F. (2004). The Role of Tomato Products and Lycopene in the Prevention of Prostate cancer: A meta-analysis of Observational Studies. *Cancer Epidemiol. Biomarkers Prev.*, 13 (3): 340-345.

10. Silaste ML, Alfthan G, Aro A, Kesaniemi YA, and Horkko S. (2007). Tomato Juice Decreases LDL Cholesterol Levels and LDL Resistance to Oxidation. *Br J. Nutr.*; 98 (6):1251-8. PMID: 17617941.
11. Makinnon and Rio L. G. (2011). Dietary Restriction of Lycopene for a period of one month resulted in significantly increased biomarkers of Oxidative Stress and bone respiration in postman woman. *J. Nutrition Health Ageing*, 15(2): 133 - 8.
12. Falodun, A., Okunrobo, L. O. and Uzoamaka, N. (2006). Phytochemical screening and anti-inflammatory evaluation of methanolic and aqueous extracts of *Euphobia heterophylla* Linn (Euphobiaceae). *African Journal of Biotechnology*, 5(6): 529-531.
13. Azaizeh, H., Fulder, S., Khalil, K. and Said, O. (2003). Ethnomedicinal knowledge of local Arab practitioners in the Middle East Region. *Fitoterapia*, 74: 98-108.
14. Bouquot Brad W. Neville, Douglass D. Dammi Carl M. Allen Jerry E. (2002). *Oral and Maxillofacial Pathology* (2nd edition) Philadelphia: WB - Saunders pp. 189 - 197 ISBN 072160033.
15. Talib N. (2016). Disorders of oral pigmentation: Drugs and Diseases in Dentistry. Medscape. Accessed on August 02 2015. Available at www.emedicine.medscape.com/article/1078143-references?src=refgatesrc1/
16. Arber (1972). *Water Plants: A Study of Aquatic Angiosperm*. Welden Wisely Limited Pp. 436.
17. Onoruvwe O. and Olorunfemi P. O. (1998). Antimicrobial Screening and Pharmacognostical Evaluation of *Dischrostachys cinerea* Root; *West Africa Journal of Biol. Science*; 7: 91-99.
18. Akerele, Z. O., Obasuyi, O., Ebomoyi, M.I., Oboh, I.E., and Uwumarongie. O.H. (2008). Antimicrobial activity of the ethanol extract and fractions of the seeds of *Garcinia kola* Heckel (Guttiferae). *African Journal of Biotechnology*, 7(2): 169-172.
19. Alade, P.I. & O.N. Irobi, (1993). Antimicrobial Activities of Crude Leaf Extracts of *Acalypha wilkesiana*. *J. Ethnopharmacol.* 39, 171-174.
20. Lin J, Opoku AR, Geheeb-Keller M, Hutchings AD, Terblanche SE, Jager AK, Staden (2015). Available at: www.globalresearchonline.net Also on: www.edu-science-com/10/comparative-study-on-phytochemical.html?m=1
21. Tiwari, P., Kumar, B., Kaur M, Kaur G & Kaur, H. (2011). Phyto-chemical Screening and Extraction: A Review. *Internationale Pharmaceutica Scientia*. 1 (1): 98-106.
22. Wagner, H. (1993). *Pharmazeutische Biologic* 5. (ed), AUFI.15 BN 3 – 437 – 20 498 –X.Gustav fisher Vwelog, Stuttgart, Germany.
23. Evans WC. (1997). *Pharmacology*, Harcourt Brace and Company, Asia, Singapore, 226.
24. Harborne JB. (1998). *A Guide to modern techniques of plant Analysis*. USA: Kluwer Academic Publisher, 54-84
25. Trease GE, Evans WC. (1996). *A textbook of Pharmacognosy*. 14th ed. London: Bailliere Tindall Ltd, 1996.
26. Fehling H. (1849). "Die quantitative Bestimmung von Zucker und Stärkmehl mittelst Kupfervitriol". *Annalen der Chemie und Pharmacie*. 72 (1): 106–113. doi:10.1002/jlac.18490720112.
27. George N J, Obot J B, Ikot A N, Akpan A E and Obi-Egbedi N O. (2010). Phytochemical and Antimicrobial properties of leaves of *Alchonea cordifolia*. *E-J Chem.*, 7(3), 1071-1079.
28. Oxoid (1985). *Oxoid manual of dehydrated culture media, ingredients and other laboratory services*. Basingstoke, England.
29. Irobi, O. N., Moo-Young, M., Anderson, W. A. and Daramola, S. O. (1994). Antimicrobial activity of bark extracts of *Bridelia ferruginea* (Euphorbiaceae). *International Journal of Pharmacognosy*, 34(2): 87-90.
30. Cheruiyot, K. R., Olila, D. and Katereggga, J. (2009). In vitro antibacterial activity of selected medicinal plants from Longisa region of Bomet district, Kenya. *African Health Sciences*, 9(S1): S42-S46.
31. Reynolds, T. and Dweck, A. C. (1999). *Aloe vera Leaf Gel: A Revise Update*. *Journal of Ethnopharmacology*, 68: 3-37.

- 375 32. Aliyu, A. B., Musa, A. M., Abdullahi, M. S., Oyewale, A. O. and Gwarzo, U. S.
 376 (2008). Activity of plant extracts used in Northern Nigerian traditional medicine
 377 against methicillin-resistant *Staphylococcus aureus* (MRSA). Nigerian Journal of
 378 Pharmaceutical Sciences, 7(1): 1-8.
- 379 33. Sukhdev S.H., Suman, P. S. K., Gennaro, L and Dev, D. R. (2008). Extraction
 380 Technologies for Medicinal and Aromatic Plants International. Centre for Science
 381 and High Technology, Trieste ed, pp 11.
- 382 34. Dahan K, Fennal M, Kumar NB. (2008). Lycopene in the prevention of prostate
 383 cancer. J. Soc Integr Oncol., 6:29-36.
- 384 35. Rao AV. (2002). Lycopene, tomatoes, and the prevention of coronary heart disease.
 385 Exp Biol Med (Maywood); 227:908-913.
- 386 36. Omodamiro O. D. and Amechi U. (2013). The phytochemical content, antioxidant,
 387 antimicrobial and anti-inflammatory activities of *Lycopersicon esculentum* (Tomato).
 388 Asian Journal of Plant Science and Research, 2013, 3(5):70-81.
- 389 37. Sung W.S., Lee I.S. and Lee D.G. (2007). Damage to the cytoplasmic membrane
 390 and cell death caused by lycopene in *Candida albicans*. J Microbiol Biotechnol.,
 391 17(1):797-804.
- 392 38. Pavlović R., Mladenović J., Radovanović B., et al. (2013). In Vitro Antimicrobial
 393 Activity of Ethanol Tomato Extracts. Conference VIVUS, 24th-25th April 2013.
 394 Biotechnical centre Naklo, Strahinj 99, Naklo Slovenia, Biotechnical Centre Naklo
 395 Higher Vocational College, Strahinj 99, Naklo, Slovenia.
- 396 39. El-Mahmood, A. M. and Ameh, J. M. (2007). In vitro antibacterial activity of *Parkia*
 397 *biglobosa* (Jacq) root bark extract against some microorganisms associated with
 398 urinary tract infections. African Journal of Biotechnology, 6(11): 1272-1275.
- 399 40. Jigna, P. and Chanda, S. V. (2007). In vitro antimicrobial activity and phytochemical
 400 analysis of some Indian medicinal plants. Turkish Journal of Biology, 3: 53-58
- 401 41. Kitonde, C. H., Fidahusein, D. S., Lukhoba, C. W. and Jumba, M. M. (2014).
 402 Antimicrobial activity and phytochemical screening of *Senna didymobotry* used to
 403 treat bacterial and fungal infections in Kenya. International Journal of Education and
 404 Research, 2(1): 1-12.
- 405 42. Yineger, H., Kelbessa, E., Bekele, T. and Lulekal, E. (2008). Plants used in
 406 traditional management of human ailments at Bale Mountains National Park, South-
 407 eastern Ethiopia. Journal of medicinal plant research, 2(6): 132-153.
- 408 43. Al-Oqaili RMS., Basim B, Istabreq M., Salman MA., Abed D. and Asaad A. (2014). In
 409 Vitro Antibacterial Activity of *Solanum lycopersicum* Extract against some
 410 Pathogenic Bacteria. Food Science and Quality Management, 27: 12-17.