

Original Research Article

In Vitro Antimycobacterial activity of Ficus sycomorus on susceptible strain of Mycobacterium tuberculosis

ABSTRACT

Aims: To evaluate the Anti-mycobacterial activity Of *Ficus sycomorus* by *In Vitro* screening against susceptible strain of *Mycobacterium tuberculosis*.

Study design: Hospital/University based cross sectional study

Place and Duration of Study: National Tuberculosis and Leprosy training center Zaria, Department of Biochemistry, Ahmadu Bello university Zaria. March 2015 to February 2017.

Methodology: The anti-mycobacterial activity of *Ficus sycomorus* (stem bark, root bark, leaves and fruits) was studied *in vitro* using standard Nitrate Reductase Assay techniques against susceptible strain of *Mycobacterium tuberculosis*. Phytochemical analysis of the n-hexane fruit extract was done using standard test reported by Mir *et al.*, 2013, Gracelin *et al.* 2013, and Harborne, 1973. Partial fractionation of the n-hexane fruit extract was done using Thin layer and column chromatography assay.

Results: n-hexane fruit extract showed activity against the tested *Mycobacterium tuberculosis* (*M-Tb*) strain. This activity was observed at 100-400 µg/mL against the susceptible strain. The crude n-hexane, leaves, root bark and stem bark extracts revealed no activity against the susceptible (*M-Tb*) strain. The methanol and aqueous extracts of fruit, leaves, stem bark and root bark also revealed no activity against the tested susceptible *M-Tb* strain. The n-hexane fruit extract (the most active extract) was then partially purified using thin layer chromatography and Colum chromatography where four fractions A-D were obtained, of which B-D were more active than A but B has the lowest minimum inhibitory concentration against the tested *M-Tb* strain. The minimum inhibitory concentration of 6.25µg/mL was obtained against the susceptible strain. The n-hexane fruit extract also revealed phytochemicals of which saponins and Alkaloid have the highest percentage content, (16.67±0.04 %) and (6.00±0.02 %) respectively

Conclusion: Therefore, these findings indicate that if *Ficus sycomorus* is properly explored and standardized it could be used as herbal drug against *Mycobacterium tuberculosis*.

Keywords: [Nitrate Reductase assay, *Ficus sycomorus*, *Mycobacterium tuberculosis*, susceptible strain}

1. INTRODUCTION

Tuberculosis (TB) is a chronic infectious airborne disease caused by the tubercle bacillus *M. tuberculosis* (1). TB infections are characterized by the growth of rod-shaped bundles of the *M. tuberculosis* bacteria which in susceptible animals, including humans produce microscopic "tubercles" consisting of chronic granulomas, some with caseous necrosis. Lung tissue is frequently infected, but other parts of the body can be involved (2). In 2009, 9.4 million new TB cases, including the HIV infected population, were recorded. Although one-third of the world's population is infected by *M. tuberculosis*, only approximately 5-10% of the infected population who are HIV uninfected will develop TB at some stage in their life. HIV and TB co-infected populations are more like to develop TB (3). Factors that contribute to the development of tuberculosis disease are complex and not completely understood but suppression of cell-mediated immunity plays a key role (4).

Tuberculosis is mostly asymptomatic and is aggravated when the immunity is compromise which arises due to conditions like malnutrition, diabetes, malignancy, and HIV/AIDS. A major problem for the control of TB is the requirement of drug regimens for six to nine months. These lengthy regimens lead to non-compliance with therapy, relapse and development of drug resistance. In order to shorten the duration of therapy, novel drugs that are active against *Mycobacterium tuberculosis*, which act through mechanisms different from those employed by the existing frontline and secondary anti-TB drugs are urgently needed. The use of herbs and other alternative therapies for the treatment of Tuberculosis is on the increase.

Phytochemicals are chemical compounds formed during the plants normal metabolic processes, as such the ethanobotanical advantage conferred by these plant based products have surpassed the chemical counter parts owing to their lesser side effect and more potent therapeutic effect. Natural products continue to play the most significant role in the drug discovery and development process. (5) Hence it is a demanding need to study the various pharmacologically valuable aspects of these medicinal plants and one of which is *Ficus sycomorus* in treating human diseases. The anti *Mycobacterium* activity can be detected by the conventional method such as Nitrate Reductase Assay by incorporating the various dilutions of the test antibiotic compound into Lowenstein Jensen (LJ) Medium and inoculating a known amount of test organisms.

Ficus sycomorus known as Baure in Hausa, is a large, semi-deciduous spreading savannah tree, up to 21 (max. 46) m, occasionally buttressed which is found almost round the subtropics of Africa (6). All parts of the plant are used in medicine. The dried fruits are taken orally by adult human beings in Venda (Cyprus) for the management of tuberculosis (7), It has also been reported for the treatment of snake bites, jaundice, chest pains, dysentery, cool, coughs and throat infections (8). Roots extracts are also recommended for cough related cases including tuberculosis, cold, and other related cases (9). Stem-bark, extracts are reported to contain pharmacologically active substances such as gallic tannins, saponins, reducing sugars, alkaloids and flavone aglycones, and was relatively safe in rats with LD₅₀ of 720 mg/kg, causing no hematological, hepatic and renal toxicities (10; 11). Therefore the current study is carried to study the anti mycobacterial activity of the n-hexene extract of *Ficus sycomorus*.

2. MATERIAL AND METHODS

2.1 Collection and processing of plant Material

Fresh leaves, Root back, Stem back and Fruit of *Ficus sycomorus* were obtained from Saye village of Zaria Local Government Area, Kaduna State, Nigeria. The sample was authenticated at the herbarium of the Department of Biological Sciences, Ahmadu Bello University Zaria. The dried leaves, root back, stem back and fruit were pulverized to powder and 100 g of each were extracted in 500ml of Distilled water, methanol, and n-hexane respectively by maceration within 48 hrs. The solution of extract was gently evaporated to

dryness in a water bath at 40°C. The extract was stored in a refrigerator at 4°C until when required.

2.2 Microbial Strain for anti *Mycobacterium tuberculosis* activity

ATCC 27294 (American culture type collection) standard strains of *Mycobacterium tuberculosis* susceptible to both isoniazid and Rifampicin were obtained from National Tuberculosis and Leprosy Training center saye Zaria.

2.3 Susceptibility testing of extracts against *Mycobacterium tuberculosis* Nitrate reductase assay method

NRA was performed as described by Golyshevskaya *et al.* (1996)¹² and Angeby *et al.* (2002)¹³ for anti-mycobacterial activity. The critical concentrations of 0.2µg/mL for INH, 40µg/mL for rifampicin and different extracts concentrations (400, 200, 100, 50, 25, 12.5, 6.25µg /mL) of each test extract sample of *Ficus sycomorus* were used. The Lowenstein Jensen (LJ) media and potassium nitrate (KNO₃ 1mg/mL) were added to the media, and growth of *M. tuberculosis* strains (in the form of pink color) was observed as growth. For each sample test of a concentration, three control bottles were prepared for inoculation of *M. tuberculosis* strain and all the bottles were inspissated.

For each concentration of the extract, 0.2 mL of inoculum suspension was inoculated into the bottles containing LJ medium with potassium nitrate. Similarly, 0.2 mL of inoculum suspension was also inoculated into the std. anti-tubercular drugs, while 0.2 mL of the 1:10 inoculum was inoculated into drug free media which served as growth controls. Falcon tubes in triplicate were utilized for each test sample extracts, anti-tubercular drugs and control bottles. After inoculation all the bottles were incubated at 37°C.

2.4 In-Vitro assay to show Antimycobacterial activity

The method used was the macro-tube dilution method described by Adeniyi *et al.* (1996)¹⁴. The fractions were serially diluted from the solutions of 10000µg/mL to obtained varying concentrations (400, 200, 100, 50, 25, 12.5 and 6.25µg/ml). The concentrations were incorporated into LJ media base and 6ml each was dispensed into falcon tubes, then inspissated at 85°C for 45 mins and inoculated with 0.2ml each with the standardized mycobacterium strain into the various test tubes containing varying concentrations. Another set of three test tubes containing only LJ media base were used as positive controls. Also test tubes containing standard anti-tuberculosis drugs and test organisms were used as negative controls. All the tested tubes and control tubes were then incubated at 37°C for a maximum of 21 days.

Griss reagent (HCl, 2% sulphanilamide and 1% n-1-naphthyl-ethylenediamine dihydrochloride, in the ratio 1:2:2) were used as an indicator of growth, that was added to one drug-free control bottle after 7, 14 or 21 days of incubation. When the color of control bottle changed to pink, then bottles with drugs were then tested with this reagent to indicate activity.

3.0 RESULTS AND DISCUSSION

The result of the anti mycobacterial activity by Nitrate Reductase assay is presented in the following tables against ATCC 27294 which is susceptible standard strain of *Mycobacterium tuberculosis*.

1.1 Anti-mycobacterial activity of aqueous, methanol and n-hexane extract of *Ficus sycomorus* against susceptible strain of Standard *Mycobacterium tuberculosis*

CONC (µg/mL)	AQUEOUS				METHANOL				N-HEXANE			
	LEAVES	STEM	ROOT	FRUIT	LEAVES	STEM	ROOT	FRUIT	LEAVES	STEM	ROOT	FRUIT
400	—	—	—	—	—	—	—	—	—	—	—	+
200	—	—	—	—	—	—	—	—	—	—	—	+
100	—	—	—	—	—	—	—	—	—	—	—	+
50	—	—	—	—	—	—	—	—	—	—	—	—
25	—	—	—	—	—	—	—	—	—	—	—	—
12.5	—	—	—	—	—	—	—	—	—	—	—	—
6.25	—	—	—	—	—	—	—	—	—	—	—	—

NEGATIVE CONTROL

RIF 40 +

INH 0.2 +

Control without Drug (positive control) = Pass

Key: + = Active (Growth absent), — = Not active (Growth present)

Table 1.1 showed the results of the anti-mycobacterial screening at different concentrations (400, 200, 100, 50, 25, 12.5 and 6.25µg/mL) of the aqueous, methanol and n-hexane crude extracts against the susceptible standard *Mycobacterium tuberculosis* strain. Crude n-hexane Fruit extract was found to be the active at a concentration of 100µg/mL. No activity was observed in the n-hexane Leave, stem bark and root bark extracts respectively when compared to the control because there was conversion of nitrate to nitrite which was an indicator by color change to pink, likewise the aqueous and methanol extracts were found to have no activity against the susceptible mycobacterium strain. Two potent first line antibiotics against tuberculosis (rifampicin 40µg/mL and isoniazid 0.2µg/mL) were used as standard drugs against the tested organisms and a free antibiotic drug control.

135 **Table 1.2: Quantitative Phytochemical Analysis of n-hexane *Ficus sycomorus* Fruit**
 136 **extract**

PHYTOCHEMICAL	% QUANTITATIVE VALUE
TANNINS	4.16±0.03
FLAVONOIDS	4.00±0.04
SAPONINS	16.67±0.04
PHENOLS	0.18±0.03
ALKALOIDS	6.00±0.02

137
 138 Table 1.2 showed the quantitative phytochemical analysis of n-hexane *Ficus sycomorus* fruit
 139 extracts. Tannins, Flavonoids Saponins, Phenols, and Alkaloid, were all present. The highest
 140 percentage was obtained in saponins (16.67±0.04) while the lowest percentages were
 141 obtained in phenols (0.18±0.10).

142 The n-hexane fruit extract was subjected to a partial purification using thin layer
 143 chromatography and column chromatography where four fraction A- D was obtained and B-
 144 D was found to be most active, as such it was screened to determined the minimum
 145 inhibitory concentration (MIC).

146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172

173 **Table 1.3: Minimum Inhibitory Concentration (MIC) of column chromatographic**
 174 **fractions of n-hexane Fruit extract against the susceptible Standard *Mycobacterium***
 175 ***tuberculosis* strain**

CONC (µg/mL)	B	C	D
200	+	+	+
100	+	+	+
50	+	+	—
25	+	+	—
12.5	+	—	—
6.25	+	—	—
NEGATIVE CONTROL			
RIF 40	+		
INH 0.2	+		

176 **Control without Drug (Positive control)**

177 **Key: + = Active (Growth absent), — = Not active (Growth present)**

178

179 Table 1.3 showed Minimum Inhibitory Concentration (MIC) of column chromatographic
 180 fractions of *Ficus sycomorus* n-Hexane fruit extract. The minimum inhibitory concentration
 181 (MIC) was determined as the lowest concentration of test samples that resulted in a no color
 182 change on the LJ slant, which signifies complete inhibition of growth in the LJ media. All the
 183 fractions exhibited low MIC against the tested strain. Fraction B exhibited the lowest MIC at
 184 concentration of 6.25 µg/ml (Plate B), while fraction C exhibited MIC at concentration of 25
 185 µg/ml (Plate C), and lastly, Fraction D exhibited MIC at 100 µg/mL (Plate D).



186

187

188

189 **MIC of Chromatographic fraction B, C and D against the standard susceptible**

190 ***Mycobacterium tuberculosis* strain**

191 **D-CONC (Drug control), M-CONC (Media control)**

192

193 The result of the investigation were very encouraging that all the *Mycobacteria* exposed to
 194 the B fraction of *Ficus sycomorus* fruit extract at a lowest concentration of 6.25µg/ml had
 195 *Mycobacterial* activity (Table 1.3 & plate B) in relation to the susceptibility of the standard
 196 drugs against *Mycobacteria* strain. Medium control bottles showed growth of *Mycobacterium*

while test bottle which received the same load of inoculation had complete inhibition of Mycobacteria, these could be due to the non polar extracting properties of n-hexane. Reports from literature show that long-chain unsaturated fatty acids such as oleic and linoleic acid extracted from n-hexane are selective inhibitors of the enoyl-acyl carrier protein reductase, which is a potent inhibitor of cell wall fatty acid synthesis (Zhing *et al.*, 2005)¹⁵. Also n-hexane extract of *Adhatoda vasica* has been reported by Ignacimuthu and Shangumam, (2010)¹⁶ to have antimycobacterial activity. Thus, this study was undertaken to validate the traditional use of *Ficus sycomorus* against *M. tuberculosis*. As such these result is in concordance with the result obtained from Arnold & Galamain, (2002)¹⁷ in the use of *Ficus sycomorus* dried fruit for the management of tuberculosis among the people of Venda (Cyprus) and (Mousa *et al.*, 1994)¹⁷ in the use of *Ficus sycomorus* fruit as Antibacterial, and antifungal activities. On the contrary, extracts from the three solvents of Root bark and Stem bark showed no activity against *M. tuberculosis* which is against the result reported by Morayi, (2013)¹⁸ for the management of tuberculosis.

Natural products are a proven agent for the development of new (19) active drugs and they have receive considerable attention as potential anti-tuberculosis drugs. *Ficus sycomorus* are endown with remarkable phytochemicals which have been found to possess a wide range of activities, which may help in protection against chronic diseases (20). Jha *et al.*, (2012)²¹ reported six quinazoline alkaloid from *Justicia adhatoda* having significant antimycobacterial activity, and in silicon analysis confirmed that these alkaloids inhibit β -ketoacyl-acyl-carrier protein synthase III (FabH), an enzyme involved in the initial step of fatty acid biosynthesis, leading to poor cell wall development and survival of bacilli (22). Sharma, *et al.*, (2008)²³, reported that a flavanoid epigallocatechin gallate/epigallocatechin-3-gallate directly inhibits fatty acid synthase systems I and II which is the fourth step of the fatty acid elongation cycle is carried out by an enoyl-acyl carrier protein reductase (InhA in *M. tuberculosis*) which catalyses an NADH-dependent reduction of the trans-2-enoyl fatty acyl chain to the saturated fatty acyl chain, by interacting with the residues near the NADH binding site. Spaonin also help to fight infection and microbial inversion (24) due the fact that they are glucosides, which includes steroid saponins and triterpenoid saponins (25) which also can form complex with proteins to reduce protein digestibility (26)

A number of studies have explored a wide range of natural products with strong activity against *M. tuberculosis* of which include, *Ricinus communis* Lin (27), *Alliu sativum*, *Zingiber officinale* (28), *Adhatoda vasica* (29) and (30), *Acalypha indica* and *Allium cepa* (31) are among the few reported to have anti-mycobacterium activity.

4. CONCLUSION

The anti mycobacterial activity showed that n-hexane *Ficus sycomorus* fruit extract has the potential to cure tuberculosis and is a promise for future therapeutic interventions, which may be probably be due to the phytochemical constituents present in the plant and could be a function of either the individual or the additive effects of the phytochemical constituents. All these findings justify the claim made in the indigenous system of medicine *Ficus sycomorus* use for the treatment of tuberculosis.

Further detailed in-vivo screening and bio activity studies need be carried out using crude solvent extracts as well as further purified constituents to comprehend their role in anti-tuberculosis activity and also pure purification to strain the active compound that confers this activity so as to develop suitable short time regiment drugs that can effectively kill Mycobacteria with lesser toxicity.

COMPETING INTERESTS

We the authors declared that no competing interests exist.

REFERENCES

- 1 Cole, S. T., R. B. Rosch, J. Parkhill, T. Garnier, C. Churcher, D. Harris, S. V. Gordon, K. Eiglmeier, S. Gas, C. E. Barry, III, F. Tekaiia, K. Badcock, D. Basham, D. Brown, T. Chillingworth, R. Connor, R. Davies, K. Devlin, T. Feltwell, S. Gentles, N. Hamlin, S. Holroyd, T. Hornsby, K. Jagels, A. Krogh, J. McLean, S. Moule, L. Murphy, K. Oliver, J. Osborne, M. A. Quali, M. A. Rajandream, J. Rogers, S. Rutter, K. Seeger, J. Skelton, R. Squares, S. Squares, J. E. Sulston, J. Taylor, S. Whitehead, and B. G. Barrell. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature International Weekly Journal of Science*. 1998; **393**:537-544
- 2 Knechel, N. A. Tuberculosis: Pathophysiology, Clinical features, and Diagnosis. *Journal of the American Association of Critical Care Nurses*. 2011; Doi: 10.4037/ccn2009968.
- 3 World Health Organization (2010) global tuberculosis control report. summary," *Central European Journal of Public Health*. 2010; **18** (4) :237-239.
- 4 Caws, M., Thwaites, G., and Dunstan, S. The influence of host and bacterial genotype on the development of disseminated disease with *Mycobacterium tuberculosis*. *PLoS Pathogens*. 2008; **4** (3): 100-134.
- 5 Newman, D. J and Cragg, G. M. Natural products as sources of new drugs over the last 25 years. *Journal of Natural Products*. 2007; **70**: 461-477.
- 6 Orwa, C., Mutua, A., Kindt, R., Jamnadass, R., and Simons, A. Agroforestry Database a tree reference and selection guide version 4.0 2009; (<http://www.worldagroforestry.org/af/treedb>)
- 7 Arnold, H. J. and Gulumian, M. Pharmacopoeia of traditional medicine in Venda Cyprus; 2002.
- 8 Sofowara, A. Medicinal Plants and Traditional Medicine in Africa, *Spectrum Books Ltd., Ibadan, Nigeria*. 1993; pp. 289-300.
- 9 Maroyi, A. Ethnobotanical study of medicinal plants used by people in Nhema communal area, Zimbabwe. *Journal of Ethno pharmacology*. 2011; **136**:347-354.
- 10 Sandabe, U.K. Pharmacological and toxicological studies of aqueous extract of *Ficus sycomorus* in laboratory animals. Maiduguri. *University of Maiduguri. Phd Thesis*; 2002.
- 11 Sandabe, U.K., Onyeyili, P.A., and Chibuzo, G.A. Phytochemical screening and effect of aqueous extract of *Ficus sycomorus* L (moraceae) stem bark on muscular activity in laboratory animals. *Journal of Ethnopharmacology*. 2006; **103**:481-483.
- 12 Golyshevskaya, V.I., Korneev, A. A., Chernousova, L.N., Selina, L.G., Kazarova, T.A. and Grishina, T.D. New microbiological techniques in diagnosis of tuberculosis. *Problemy Tuberkuleza*. 1996; **6**:22-25.
- 13 Angeby, K. A., Klintz, L. and Hoffner, S. E. Rapid and inexpensive drug susceptibility testing of *Mycobacterium tuberculosis* with a nitrate reductase assay. *Journal of Clinical Microbiology*. 2002; **40**(2):553-555.
- 14 Adeniyi, B. A, Odelola H. A, Oso, B. A Antimicrobial potentials of *Diospyros mespiliformis* (Ebenaceae). *African Journal of Medicinal Science*. 1996; **25**: 221-224
- 15 Zhang, Y., Yew, and W. W. Mechanisms of drug resistant in *Mycobacterium tuberculosis*. *International Journal of Tuberculosis Lung Diseases*. 2009; **13**(11) :1320-1330

- 301 16 Ignacimuthu, S and Shanmugam, N. Antimycobacterial activity of two natural
302 alkaloids, vasicine acetate and 2-acetyl benzylamine, straind from Indian shrub
303 Adhatoda vasica Ness leaves. *Journal of Biological science*. 2010; 35(4): 565–570
- 304 17 Arnold H. J. and Gulumian, M. Pharmacopoeia of traditional medicine in Venda
305 Cyprus; 2002.
- 306 18 Mousa, O., Vuorela, P., Kiviranta, J., Abdel, Wahab, S., Hiltunen, R., and Vuorela, H.
307 Bioactivity of certain Egyptian Ficus species. *Journal of Ethnopharmacology*.1994;
308 41:71–76.
- 309 19 Maroyi, A . Ethnobotanical study of medicinal plants used by people in Nhema
310 communal area, Zimbabwe. *Journal of Ethno pharmacology*. 2011; 136:347–354.
- 311 20 Newman, D.J, Cragg GM. Natural Products as Sources of New Drugs over the last 25
312 years. *J. Nat. Prod*. 2007; 70(3): 461-477.
- 313 21 Mir, M. A., Sawhney, S. S. and Jassal, M. M. S. Qualitative and quantitative analysis
314 of phytochemicals of Taraxacum officinale Wudpecker *Journal of Pharmacy and*
315 *Pharmacology*. 2013; 2 (1), 1 – 5
- 316 22 Jha, D. K., Panda, L., Lavanya, P., Ramaiah, S., and Anbarasu, A. Detection and con-
317 firmation of alkaloids in leaves of Justicia adhatoda and bioinformatics approach to
318 elicit its anti-tuberculosis activity. *Applied Biochemical Biotechnology*. 2012; 168: 980–
319 990.
- 320 23 Sharma, S.K., Kumar, G., Kapoor, M., and Surolia, A. Combined effect of epigallo-
321 catechin gallate and triclosan on enoyl-ACP reductase of Mycobacterium tuberculosis
322 *Biochemical Biophysics Res Commun*. 2008; 368: 12– 17.
- 323 24 Sodipo, O. A. Akiriyi, Y. A and Ogunbanosu Y. U. Studies on certain characteristics
324 of extracts or barks of pansinystalia macruceras (K. Schem) piere, exbeila, *Global*
325 *Journal for Pure Applied Science*. 2000; 6: 83-78.
- 326 25 Dei, H.K (2011). Poultry science, department of animal science, University for
327 developmental studies, Tamale, Chana at June 2011; 90 (6): 1239-44.
- 328 26 Shimoyamada, M., Ikedo, S., Dotsubu, R., and Watanaba, K. Effects of soya beans
329 saponins on chymotrptic hydrophyses. 1998; chem.. vol. 46, P. 4793-4797.
- 330 27 Ladda., P .L., Magdum., C. S. Evaluation of anti-tubercular activity of Ricinus
331 communis Linn. By proportion, NRA and bact / alert methods *International Journal of*
332 *Pharmacy and Pharmaceutical Sciences*. 2012; 4, 3, 474-478
- 333 28 Mann., A, Ibrahim., K, Oyewole., A.O., Amupitan., J.O, Okogun., J.I. Antimycobacterial
334 activity of some medicinal plants in Niger State, Nigeria. *African J Infectious diseases*.
335 2009b; 3:44–48.
- 336 29 Ignacimuthu, S and Shanmugam, N. Antimycobacterial activity of two natural alkaloids,
337 vasicine acetate and 2-acetyl benzylamine, straind from Indian shrub Adhatoda vasica
338 Ness leaves. *Journal of Biological science*. 2010; 35(4): 565–570
- 339 30 Gupta, K.C, and Chopra, I.C. Anti-tubercular action of Adhatoda vasica (N.O.
340 acanthacea). *Indian Journal of Medical Research*. 1954; 42: 355-358.
- 341 31 Gupta, R, Thakur, B, Singh, P, Singh, H.B, Sharma, V.D, Katoch, V.M, and Chauhan,
342 S.V.S. Anti-tuberculosis activity of selected medicinal plants against multidrug
343 resistance mycobacterium tuberculosis strain. *Indian journal of medical research*.
344 2010; 131: 809-813