

Effect of *Strophanthus hispidus* Dc. (Apocynaceae)

Roots Extract on Normal and Hyperglycemic Rats

ABSTRACT

Objective: To investigate the hypoglycemic activity of both ethanol extract *Strophanthus hispidus* root (EESHR) and chloroform extract *Strophanthus hispidus* root (CESHR) respectively.

Methods: Both ethanol extract *Strophanthus hispidus* root and chloroform extract *Strophanthus hispidus* root were evaluated respectively for hypoglycemic activity using Fasting blood glucose (FBG) method. Chloroform extract *Strophanthus hispidus* root was further fractionated using standard column chromatography techniques and the fractions were evaluated for hypoglycemic using standard oral glucose tolerance test method.

Results: In the experiments, the hypoglycemic activity of the chloroform extracts *Strophanthus hispidus* root was better than that of ethanol extract *Strophanthus hispidus* root at 500mg/kg respectively for both extracts which were significant ($p=0.05$) and compared to glibenclamide .

CONCLUSION: The results showed that chloroform extract *Strophanthus hispidus* root fraction (3-4) at 200mg/kg possesses significant hypoglycemic activity better than that of the standard drug-glibenclamide. This supports its use as a potent anti-diabetic drug in alternative herbal medicine.

Keywords: *Strophanthus hispidus*, root extract, hypoglycemic, glibenclamide

23

24

25 1. INTRODUCTION

26 The oral glucose tolerance tests (OGTT) indicates the body's ability to utilize a type of sugar
27 named glucose that is the body's main source of energy. Oral glucose tolerance tests, a test of
28 immerse value and sentiment, in favour of using fasting plasma glucose concentration alone
29 as a practical attempt to simplify the diagnosis of diabetes [1]. Hyperglycemia is an important
30 factor in development and progression of the complication of diabetes mellitus [2].

31 The plant SPH, popularly known as brown *strophanthus*, and hairy *strophanthus* in West
32 Africa, including Nigeria, is indigenous to Africa [3]. A deciduous shrub of 5 m tall and up to
33 100 cm wide, having its stem bark dark grey in colour, with few lenticels, has been reported
34 to have diverse medicinal uses; for example, in the Savannah Zone of West Africa, the latex
35 and seeds of *Strophanthus hispidus* are used as arrow poison, while decoctions of root, stem
36 bark or leaf are used externally to treat skin diseases, leprosy, ulcers, malaria, dysentery,
37 gonorrhea and diabetes [3]. The leaves have been reported to possessed hypoglycemic
38 properties [4]. Similarly, the anti-inflammatory properties of the aqueous of the root extract
39 have been studied [5]. In Nigeria and Ghana, the root decoction is ingested to treat rheumatic
40 diseases and diabetes, while in Togo; the root bark macerate is employed for treating oedema
41 [3]. The trace elements and major elements present in the plant root have been reported by
42 Akinlami et al. [6]. The seeds and the plant extract of *Strophanthus kombéare* beneficial as a
43 cardiac drug or a diuretic [7]. These plants are mainly used in the treatment of congestive
44 heart failure; however, their toxicity limits their extensive use. Ouabain, a more common
45 name for G-strophanthin a poisonous glycoside obtained from the fruits of *Strophanthus*
46 *gratus*, is used widely to block the sodium pump for in vitro studies. K-strophanthin, isolated

from *Strophanthus kombé*, was also used for the treatment of heart diseases until its adverse side effects were observed. The cytotoxicity and structural characterization of various cardenolide glycosides have been extensively studied [7]. Cowan et al isolated lignans from the stem of *Strophanthus gratus* [8]. Three lignans; pinoresinol, 8-hydroxypinoresinol and olivil have been isolated from the stem [8]. In this study, the blood lowering effects of chloroform and ethanol crude extracts of *S.hispidus* respectively were conducted through oral glucose tolerance test in Wistar albino rats.

2. MATERIALS AND METHODS

2.1 Collection of plant materials

Fresh roots of *Strophanthus hispidus* (plate.1) were purchased from the market, in Ondo southwest, Nigeria and identified at the Department of Crop, Soil and Pest Management, Federal University of Technology, Akure, Nigeria and specimen sample were kept in the Harbarium.



Plate 1: Roots of *Strophanthus hispidus* DC. (Apocynaceae)

65

66 **2.2 Extraction**

67 The identified plants materials were air dried for four weeks and pulverized into fine powder
68 using clean and sterile Mortar and Blender. The powdered samples were sieved and quartered
69 to obtain a representative of 1000g used for this study. Two hundred grams of the powered
70 roots of *S. hispidus* were extracted with 1L of 96% ethanol or chloroform respectively at
71 room temperature (25-30°C) by maceration for 72h. After the extraction, the solvents were
72 removed from the extracts by distillation and drying at 60°C in a water bath, leaving the
73 extracted portion as a residue. The crude Ethanol Extract *S. hispidus* Root (EESHR) and
74 Chloroform Extract *S. hispidus* Root (CESHR) were stored and kept in air tight bottle in a
75 refrigerator until used for bio-assay. The crude EESHR was dissolved in normal saline to
76 obtain the required concentration of 500mg/kg while CESHR was first dissolved in Tween-80
77 and followed by normal saline to also obtain the needed concentration of 500mg/kg that were
78 required for the bioassay.

79 **2.3 Phytochemical screening of crude plant extracts**

80 Qualitative phytochemical screening of ethanol extract *S. hispidus* root (EESHR) and
81 chloroform extract *S. hispidus* root (CESHR) respectively were carried-out by standard
82 procedures [9], [10], and [11].

83

84 **2.4 Test animals**

85 A sum total of 50 male albino rats weighing about 120-220 gm. of were procured from
86 Animal house of Federal University of Technology, Akure, Nigeria. All the animals were
87 kept in standard polypropylene cages and maintained at 27°C ± 2°C under 12 h. dark/light

88 cycle. The animals were fed with standard rat feed and water was given ad libitum. Ethical
89 clearance for handling the animals was strictly adhered to as described by Akhtar et al. [13].

90 **2.5 Experimental design**

91 Initial screening of the ethanol extract *S. hispidus* root (EESHR) and chloroform extract
92 *S.hispidus* root (CESHR) respectively was done to evaluate their hypoglycemic potential with
93 dosage of 500mg/kg for each of the extracts. These were administered orally by gavage in
94 normal rats by conducting fasting blood glucose (FBG) study. Oral glucose tolerance test
95 (OGTT) studies were done on separated components of chloroform extract *S.hispidus* root
96 (CESHR), which was subjected to Column chromatography. The chloroform extract *S.*
97 *hispidus* root (CESHR) was identified as the most effective hypoglycemic than ethanol
98 extract *S. hispidus* root (EESHR) by initial screening.

99 **2.6 Assessment of hypoglycemic activity in normal healthy rats**

100 The animals were fasted for 18 h with free access to water prior to the administration of the
101 extracts. After fasting the blood glucose levels was measured using the glucose-oxidase
102 principle and the fasting blood glucose greater than 50 mg/dL was included in the study for
103 the normoglycemic group . The normoglycemic rats were randomly assigned into four
104 groups (1-4) of five rats (n = 5) each as follows,

105 Group 1: Normal treated with 500 mg/kg of CESHR

106 Group 2: Normal treated with 500mg/kg of EESHR

107 Group 3: Normal treated with 5 ml/kg of normal saline

108 Group 4: Normal treated with 5mg/kg of gibenclamide

109 Fasting blood glucose (FBG) was taken initially and then blood samples were collected from
110 tail vein at 2, 4, 6 and 8 h after administering the extracts.

111

112

2.7 Column chromatography separation of chloroform extract *S.hispidus* root (CESHR)

Thin Layer Chromatography (TLC) was employed using silica gel 60 F₂₅₄ precoated plates and solvent system: hexane: ethyl acetate (9:1), (8:2), (7:3), (6:4), (5:5), (4:6), (3:7), (2:8). TLC was carried out on the crude extract to estimate the number of components. One drop of glacial acetic acid was added to each solvent system to aid distinct separation of spots into their different components. After the development, the chromatogram was then allowed to dry before it was transferred into the iodine tank. Initial spotting of the crude chloroform extract *S. hispidus* root obtained on TLC plates indicated more than one components, hence the need for proper separation with a column chromatograph. Column chromatograph was used to separate crude chloroform extract *S. hispidus* root. 3 cm x 100 cm column size and a silica gel of mesh size 60-200 were used. Silica gel (180 g) was made into slurry by adding the solid adsorbent to a quantity of packing solvent (hexane). The slurry was swirled until it was homogeneous and relatively free of entrapped air. Prepared slurry was slowly poured into the column with the tap opened such that it gently settled uniformly. The column was constantly tapped to get rid of any trapped air bubbles and also ensure a leveled layer of the adsorbent. 6 g of crude extract was pre-adsorbed on silica gel. Hexane was used to wash the packed column prior to analysis. This pre-adsorbed sample was allowed to dry before loading onto the column. The ratio of the silica gel to that of extract was 30:1. The loaded sample was then covered with pure white sand and a small piece of cotton wool. The top of the column was usually filled with enough solvent/mixture of solvents to avoid cracking. The elution started with 100 % non-polar hexane after which the polarity of the solvent(s) was increased gradually with EtOAc until 100 % in a gradient elution chromatographic technique. The solvent system used in eluting the column was as follows: hexane 100 %, ethyl acetate/hexane 5 %, 10 %, 15 %, 20 %, 25 %, 40 %, 50 %, ethyl acetate 100 %. A total of 40 eluents of 50 ml each were collected. TLC of the eluents collected was also carried out using

the same solvent system employed in the column chromatography. This was done to ascertain the number of components in it and also to pool together similar eluents which contain the same component based on their R_f values. The fractions obtained (50ml) were grouped and coded to afford four main fractions: CESHr (3-4), CESHr (5-9), CESHr (10-12), and CESHr (18-25).

2.8 Oral glucose tolerance test in chloroform extract *S.hispidus* root (CESHR)

This test was performed according to [14]. Rats were divided into five groups (n = 6). They were fasted overnight and accessed to water only. Blood was taken from the lateral veins of the tail and the blood sugar levels were initially monitored with a glucometer (One touch Horizon). Group I served as control groups were treated with vehicle (0.5% Tween 80 solution), group II treated with 5mg/kg of gibenclamide as a standard drug. Chloroform Fractions CESHr (3-4) serves as group III, CESHr (5-9) served as group IV, CESHr (10-12) served as group V and CESHr (18-25) served as group VI each were treated with a dose of 200 mg/kg and After 30 min, the animals were treated with 5% (wt/v) glucose orally. Blood glucose levels were monitored from lateral tail veins at 30, 90 and 150 min intervals after post glucose challenge. Data were expressed as mean $\pm$ standard error of mean. Statistical comparison were performed by one-way ANOVA followed by Dunnett's multiple comparison test, and the values were considered statistically significant when $p = .05$.

3. RESULT AND DISCUSSION

3.1 Qualitative Phytochemical Screening

Tables 1. presents the result of Qualitative phytochemical screening of ethanol extract *S. hispidus* root (EESHr) and chloroform extract *S.hispidus* root (EESHr) respectively. Both extracts contained flavonoids, alkaloids, saponins. Cardiac glycosides, tannins and anthraquinones. The results are in agreement with similar research work carried out by Agbaje and Fageyinbo [15]. Phytochemicals compounds present as secondary metabolites

play vital roles in the bioactivity exhibited by medicinal plants used ethno botanically [16]. Some plant compounds have pharmacology effects, which forms the basis of chemical synthesis of drugs used in modern medicine [16]. Flavonoids are polyphenols compounds that are structurally classified as flavonols, flavones, isoflavones, catechins, anthocyanidins and chalcones [17]. Numerous flavonoids have been identified; many of which occur in fruits, vegetables and beverages tea, coffee, beer, wine and fruit drinks [18]. Flavonoids (Figure 1) have potential beneficial effects on human health. They have been reported to have anti-viral, anti-platelet, ant-inflammatory, anti-tumors, anti-diabetic and antioxidant activities [18]; [19].

Table 1: Qualitative phytochemical screening of ethanol extract *S. hispidus* root (EESHR) and chloroform extract *S.hispidus* root (CESHR) respectively.


General Test	Specific Test	EESHR	CESHR
Test for Flavonoids	Ferric Chloride Test	+	+
Test for Glycosides	Keller-Kiliani Test	+	+
Test for Tannins	Ferric Chloride Test	+	+
Test for Alkaloids	Dragendoffs test	+	+
Test for Anthraquinones	Hydrogen Chloride Test	+	+
Test for Saponins	Frothing Test	+	+

3.2 Hypoglycemic effect of extracts on normal rats

Figure 1 describes the result of hypoglycemic effect of chloroform extract of *S.hispidus* roots (CESHR) and ethanol extract of *S.hispidus* roots (EESHR) respectively, on fasting blood glucose (FBG) of normal rats. Rats treated with 500mg/kg CESHR showed a maximum fall of 39.2% in Fasting blood glucose (FBG) after 8hrs. of oral administration, while the fall of

26.62% was observed with 500mg/kg of EESHR respectively. The hypoglycemic activity of the chloroform root extract was better than that of ethanol root extract. The result was in agreement with similar report by of Ojiako et al. [20].

Fig.1. Hypoglycemic effect of *S.hispidus* root extracts on BGL of normal rats



Each value shown in mean $\pm$ S.D. (n = 5). Control (distilled water); Chloroform extract *S.hispidus* roots (CESHR-500mg/kg); Ethanol extract *S.hispidus* roots (EESHR-500mg/kg); STD (gibenclamide -5mg/kg).


3.3 Hypoglycemic effect of chloroform extracts fractions on OGTT of normal rats

Figure 2 depicts the hypoglycemic effect of a single oral administration of 200mg/kg each of different chloroform extract fractions of *S.hispidus* root on OGTT of normal rats.

Experimental induction of hyperglycemia by intragastric ingestion of glucose resulted in a 1.5 to 2-fold increase in plasma glucose levels (comparing bar groups of 0 minute with the bar groups of 30 minute, (fig. 2). The CESH fraction (3-4) produced a maximum fall of 42.45% at two hours after glucose administration, whereas fall of 32.74%, 34.66% and

205 28.93% were observed with the CESHHR fraction (5-9), CESHHR fraction (10-12) and CESHHR
206 fraction (18-25) respectively. A standard antihyperglycemic drug, glibenclamide, when
207 administered to glucose loaded rats, reduced blood glucose level by 49.00%. The
208 results demonstrated that the CESHHR fraction (3-4) possesses antihyperglycemic potential
209 more than the others fractions. The CESHHR fraction (3-4) was comparable to that of the
210 standard drug, glibenclamide, in its effectiveness in lowering blood glucose. This suggests
211 that the fraction contains antihyperglycemic constituents.

212 **Fig.2. Hypoglycemic effect of Chloroform extract fractions *S.hispidus* roots (CESHR) on**
213 **BGL of normal rats during OGTT**



214
 215 .Each value shown in mean $\pm$ S.D. (n = 6). Control ((0.5% Tween 80 solution); Chloroform
 216 extract fractions *S.hispidus* roots (CESHR-200mg/kg); STD (glibenclamide-5mg/kg).

217

218 However, it can be suggested that the mode of action of action of fractions CESHR (3-4) is
 219 probably mediated by an enhanced secretion of insulin, like glibenclamide a biguanides. In
 220 the light of the above pharmacology results, further works is on-going in our laboratory to
 221 possible isolate and charaterize the possible hypoglycemic compounds from the most active
 222 fraction using Infra-red spectroscopic, ^1H , ^{13}C , 2D-NMR spectroscopic and Electron Spray

223 Mass spectroscopy. Finally, the present study has given idea of the hypoglycemic compounds
224 present in the reported plant fraction.

225 **4. CONCLUSION**

226 The result showed that CESH fraction (3-4) at 200mg/kg possesses significant
227 hypoglycemic activity better than that of the standard drug-glibenclamide. This supports its
228 use as a potent anti-diabetic drug in alternative herbal medicine.

229 **CONSENT**

230 It is not applicable

231 **ETHICAL APPROVAL**

232 The authors declare that this work was not against public interest. Animal experiments were
233 conducted in accordance with NIH guidelines for care and use of Laboratory animals (Pub.
234 No. 85-23, Revised).

235 **REFERENCES**

- 236 1. Famobuwa OE, Akinlami OO, Agbowuro AA. Oral glucose tolerance test in normal
237 control and glucose induced hyperglycemic rats with *Tetracarpidium conophorum*
238 (Nigerian Walnuts). Journal of Natural Products and Biomedical Research.
239 2015;1(2),48-50.
- 240 2. Luzi I. Pancreas transplantation and diabetic complications. The New England Journal
241 of Medicine. 1998; 339, 115-117.
- 242 3. Hutchinson J and Dalziel JM. Flora of West Tropical Africa. II. Millbank, London
243 crown agents for oversea Government and Administration 4, 1963; 221-221.

4. Ojiako, OA. and Igwe, CU. A time-trend hypoglycemic study of ethanol and chloroform extracts of *Strophanthus hispidus*. Journal of Herbs, Spices and Medicinal Plants. 2009; 15(1), 1-8.
5. Agbaje, EO and Fageyinbo, MS. Evaluating Anti-Inflammatory activity of aqueous root extract of *Strophanthus hispidus* DC. (Apocynaceae) International Journal of Applied Research in Natural Products. 2012; 4 (4), 7-14.
6. Akinlami, O., Lajide, L., Oloyede, H and Ojo, B. Trace Elements and Majors Minerals Evaluation in *Fluerya aestuans* Linn. (Urticaceae). International Journal Pharma Science. 2014; 3(5): 333-337.
7. Burkil, HM. The useful plants of west tropical Africa (2nd ed.). Vol. 1: Royal Botanical Garden. UK, KEW 1985:641-3
8. Cowan, S., Stewart, M., Abbiw, DK., Latif, Z., Sarker, SD., and Nash.RJ. Lignans from *Strophanthus gratus*. *Phytochemical communication*. Fitoterapia. 2001; 72 .80 – 82.
9. Harborne, JB. Methods of plant analysis. In: *Phytochemical Methods*. Chapman and Hall, London; 1980, 67-78.
10. Sofowora, LA. African Medicinal plants proceeding of conference, university of Ife, Nigeria. 1982; 70-73.
11. Edeoga, HO., Okwu, DE, and Mbaebie, BO. 2005. Phytochemical constituents of some Nigerian medicinal plants. *Africa journal of Biotechnology*. 2005; 4(7), 685-688.
12. Nayak, BS and Pinto, LM. *Catharanthus roseus* flower extract has wound-healing activity on Sptague Dawley rats. *BMC Complementary and Alternative Medicine* 2006; 6, 41.

- 268 13. Akhtar, MA, Rashid, M, Ibne Waheed, MI, Islam, MR, Shaheen, SM, Islam, MA and
269 Ahmed, M. Comparison of large-term antihyperglycemic and hypolipidemic effects
270 between *Coccinia cordifolia* (Linn) and *Catharanthus roseus* (Linn) in alloxan-
271 induced diabetic rats. Research of Medical Sciences Medical Science. 2007; 2(1), 29-
272 34.
- 273 14. Shirwaikar, A and Rajendran, K. Effect of aqueous bark extract of *Garuga pinnata*
274 Roxb. in streptozotocin-nicotinamide induced type-II diabetes mellitus. Journal of
275 Ethnopharmacology.2006; 112, 1-6
- 276 15. Agbaje, EO and Fageyinbo, MS. Evaluating Anti-Inflammatory activity of aqueous
277 root extract of *Strophanthus hispidus* DC. (Apocynaceae). International Journal of
278 Applied Research in Natural Products. 2012; 4 (4), 7-14.
- 279 16. Kar, A and Choufhary, BK. Important mineral content of a few Ayurvedic herbs with
280 a discussion on medicinal aspects. Indian Drugs.1994; 31(3): 127-130
- 281 17. Lee, KV, KIM, YJ, Lee, HJ and Lee, CY. Cocoa has more Phenolic Phytochemical
282 and higher antioxidant capacity than teas and red wine. Journal of Agriculture and.
283 Food Chemistry. 2003; 51(25):7292-5
- 284 18. Wikipedia., Flavonoids.en.wikipedia.org/wiki/Flavonoids, 2012b.
- 285 19. Rice-Evans, C.A. Flavonoids and isoflavonones absorption, metabolism and
286 bioactivity. Free Radical and Biological Medicine. 2004; 36:827-828.
- 287 20. Ojiako, OA and Igwe, CU. A Time-Trend Hypoglycemic Study of Ethanol and
288 Chloroform Extracts of *Strophanthus hispidus*. Journal of Herbs, Spices & Medicinal
289 Plants. 2009; 4, 45-49.