

BODY WEIGHT CHANGES AND SERUM GROWTH HORMONE COMPARATIVE ASSESMENT IN FEMALE LACTATING WISTAR RATS (Rattus Norvegicus) TREATED WITH METOCLOPRAMIDE AND SOME ATYPICAL ANTIPSYCHOTICS

ABSTRACT

Aims: The design of this study was to comparatively evaluate body weight changes and the serum growth hormone level in female lactating Wistar rats treated with Normal saline, Metoclopramide Olanzapine and Risperidone.

Study design: Twenty (20) Female wistar rats weighing between 140-180g after parturition, randomly divided into four (4) groups of five (5) animals each (n=5) were used for the study

Methodology: The study was an experimental study comprising of four groups. Group I was the control that received 2ml/kg normal saline; Group II: received Metoclopramide (5 mg/kg b.w); Group III: received Olanzapine (5 mg/kg b.w) while group IV received Risperidone (5 mg/kg b.w). All administration was done via oral gavage using an oral cannula, for the period of fourteen (14) days at 06:00 h daily. At the end of the experiment, blood samples were obtained using a 5 ml syringe via cardiac puncture and the sera used for growth hormone. Animals were weighed using a digital weighing balance at 06:00h and their weight recorded in grams for days 0, 3, 6, 9, 12 and 14.

Results: There was a significant increase in mean serum growth hormone concentration ($P < .05$) in the Metoclopramide treated group; (19.35 ± 1.14 ng/ml) compared to normal control; (11.53 ± 1.64 ng/ml), Olanzapine treated; (11.28 ± 0.35 ng/ml) and Risperidone treated; (11.30 ± 0.70 ng/ml). However, there was no statistically significant difference in the Olanzapine and Risperidone treated groups compared to the Normal control ($P > 0.05$). There was no statistically significant difference ($P > .05$) observed in the body weight changes although at day 14, Metoclopramide showed the highest increase (180 ± 1.30 g) compared to the Normal control (172 ± 1.83), Olanzapine (173 ± 2.30) and Risperidone (175 ± 3.80).

Conclusion: It is concluded that metoclopramide at a dose of 5 mg/kg significantly increases serum growth hormone level (ng/ml) compared to Olanzapine and Risperidone in female lactating wistar rats, with a non-significant increase in body weight of the dams.

Keywords: Lactation, Olanzapine, Risperidone and Dopamine.

1.0 INTRODUCTION

Growth hormone, also known as somatotropin is a peptide hormone which stimulates growth, cell reproduction and regeneration in humans and other animals. It is a single chain polypeptide with 191 amino acid synthesized, stored and secreted by the somatotrophic cells

12 within the adenohypophysis. Somatostatin also secreted from the hypothalamus, inhibits the
13 release of Growth hormone [1]. Growth hormone is secreted from the anterior pituitary gland
14 in a pulsatile fashion and acts on target tissues to promote growth. Secretion from the
15 pituitary is the major source of Growth hormone; however several other tissues synthesize
16 GH, including brain, placenta, mammary and pineal gland [2]. GH is involved in maintaining
17 mammary cell number during lactation of mice. In ruminants, the action of GH on the
18 mammary gland is thought to be mediated mainly by the insulin-like growth factor (IGF)
19 signaling axis. Although the action of GH is mediated mainly through the IGF axis, there is
20 evidence that GH may act independently of IGF-I to stimulate milk production [3, 4]. In
21 addition, expression of GH receptor has been detected in mammary tissue. The GH receptor
22 belongs to a super family of transmembrane receptors, of which PRL receptor is a member
23 [5].

24 Metoclopramide drug was originally commercialized in Europe as an antipsychotic and later
25 in the US as a gastro-kinetic agent that increases gastrointestinal motility. Its first reported
26 use as a galactagogue was in 1975 [6]. It has also been evaluated in many clinical trials [7].
27 It's reported side effects in humans include anxiety, insomnia, severe depression and
28 seizure. Infants that consumed milk from treated mother had intestinal discomfort [8]. 10 mg
29 administered by oral route three times daily for 10 days increases milk production [9].
30 Dopamine D₂ receptor blockade in the brain is a general pharmacodynamic property of all
31 antipsychotics which are mainly of two (2) groups known as 'typical' and 'atypical'
32 antipsychotics. Although with typical antipsychotics the level of D₂ receptor blockade is
33 directly related to the antipsychotic effect, with atypical agents the situation is more
34 complicated [8].

35 Olanzapine (OLZ) is a thienobenzodiazepine derivate structurally similar to clozapine that is
36 effective in treating schizophrenia and acute manic episodes, and in preventing the
37 recurrence of bipolar disorders [10]. Treatment with OLZ is associated with a higher risk of
38 weight gain and, more extensively, metabolic syndrome than other typical and atypical

antipsychotics [11, 12]. It also has a low propensity to cause extrapyramidal effects or sustained increases in prolactin levels [13].

Risperidone is a benzisoxazole derivative, and has a very strong affinity for serotonin (5-HT_{2A} and 5-HT₇) and dopamine D₂ receptors (its affinity for D₃ and D₄ receptors is three times lower) [13]. It is rapidly absorbed after oral administration, with peak plasma concentrations being reached in about one hour; its oral bioavailability is about 70-85 %. It mainly undergoes 9-hydroxylation in the liver that yields the active 9-hydroxy risperidone metabolite (9-OH-RSP), a step that is mainly catalysed by CYP2D6 and, to a lesser extent, CYP3A4; alicyclic dehydroxylation and oxidative N-dealkylation are minor metabolic pathways [13].

Therefore this study was aimed at body weight changes and serum growth hormone in female lactating wistar rats treated with metoclopramide and some atypical antipsychotic drugs.

1.1 EXPERIMENTAL SITE

This study was conducted at the Department of Human Physiology, Faculty of Medical Sciences, College of Health Sciences, Ahmadu Bello University. Zaria is located between latitudes 11° and 3' N, and between 7° and 42' E, at an altitude of 670 m above the sea level and 664 km away from the sea, in the Northern Guinea Savanna zone. [14]. The research was carried out according to the guidelines of Ahmadu Bello University animal use and care policy.

58


1.2 CHEMICALS AND DRUGS

Olanzapine and Risperidone; Stallion Laboratories Pvt. Ltd.: Ahmedabad India. Manufacturing Licence No.: G/898, NAFDAC Reg. No.: A4-1677 and Metoclopramide hydrochloride tablets 10 mg; Jiangsu Pengyao Pharmaceutical Inc. China, Manufacturing Licence No.: B131205, NAFDAC Reg. No.: 04-6476 and normal saline, used for this study

64 were purchased from the Pharmacy of the Ahmadu Bello University Teaching Hospital
65 (ABUTH) Shika; 8 km away from Samaru on Latitude 112° 12' N and Longitude 07° 37' E,
66 Zaria Kaduna state, Nigeria

67 1.3 EXPERIMENTAL DESIGN

68 Twenty female lactating Wistar rats were purchased from the animal house of Human
69 physiology department, faculty of medicine, Ahmadu Bello University. Mating protocol was
70 carried out during which the 20 females were divided randomly into four (4) groups of five
71 rats each (n=5). Each group of females was then mixed with a corresponding male group in
72 a ratio of 5:5 (male to female) and allowed to cohabit for mating. At parturition, the 20
73 females were randomly divided into four (4) groups of five (5) animals each (n=5) and
74 treated as follows:



75

76 1.4 EVALUATION OF SERUM GROWTH HORMONE CONCENTRATION IN FEMALE 77 LACTATING WISTAR RATS

78 The animals were anaesthetized on day 14 at the termination of the experiment by
79 intraperitoneal injection of ketamine and diazepam at 75 and 25 (mg/kg) respectively. The
80 thorax of the each anaesthetized animal was cut open and with the aid of a 5ml syringe, the
81 pulsating heart of the rat pierced at the left ventricle and blood obtained from which sera was
82 obtained for the determination of growth hormone using the Rat-Growth hormone ELISA kit
83 according to the manufacturer's manual.

84

85

86 1.5 STATISTICAL ANALYSIS


87 Data obtained from the study were expressed as mean $\pm$ SEM. Statistical analysis was
 88 carried out using version 20 of SPSS with the aid of one way analysis of variance (ANOVA)
 89 and Tukey's post-hoc test. Values with ($P < .05$) were considered significant

90 2.0 RESULT

91 2.1 Effect of Metoclopramide, Olanzapine and Risperidone on serum growth hormone 92 levels in female lactating wistar rats

93 The serum growth hormone level (ng/ml) in female lactating wistar rats treated with
 94 metoclopramide was significantly increased ($P < .05$) compared to the Normal control;
 95 19.35 ± 1.14 ng/ml vs 11.53 ± 1.164 ng/ml respectively. Although changes were observed in
 96 the other treated groups, it was however not statistically significant compared to the Normal
 97 control.

98



99

100

101 Figure I: Serum Growth hormone levels in female lactating wistar rats treated with Normal
 102 saline, Metoclopramide, Olanzapine and Risperidone for fourteen (14) days. D= day.
 103 Different superscript (a,b) indicates statistically significant changes ($P < .05$), N. saline=
 104 Normal Saline, GH= Growth hormone. Data was analysed using ANOVA after which Tukey
 105 post hoc test was carried out on SPSS.

2.2 Effect of Metoclopramide, Olanzapine and Risperidone on Body weight changes in female lactating wistar rats treated for fourteen (14) days;

Metoclopramide treated groups showed the highest gain in weight at day 14 of the experiment; $180 \pm 1.30\text{g}$ compared to Normal control ($172 \pm 1.83\text{g}$), Olanzapine ($173 \pm 2.30\text{g}$) and Risperidone ($175 \pm 3.80\text{g}$). Although changes were observed, it was however not statistically significant ($P > .05$)




Figure II: Showing body weight changes in female lactating wistar rats treated with Normal saline, Metoclopramide, Olanzapine and Risperidone for fourteen (14) days. D= day. Data was analysed using SPSS while Repeated Measure ANOVA was employed to check the effect of time.

3.0 DISCUSSION

The levels of serum growth hormone from this present study showed significant increase in the Metoclopramide treated group compared to the control ($P < .05$). The result obtained from the metoclopramide treated could have been due to a pro-activity of this substance on the growth hormone releasing hormone in the hypothalamus resulting in increased serum growth hormone secretions from the cells in the pituitary gland. It has been shown that dopaminergic drugs like metoclopramide can elicit GH secretion through an increased hypothalamic growth hormone releasing hormone [15]. This result has also been corroborated by the increase in weight observed in the metoclopramide treated group which

is consequence of the activity of elevated serum level of growth hormone. The action of GH on the mammary gland is thought to be mediated mainly by the insulin-like growth factor (IGF) signaling axis [16]. Metoclopramide is known to elevate serum prolactin during which is responsible for milk synthesis. This action of the said drug could have also indirectly caused the complementary release of growth hormone in response to increased milk synthesis for the purpose of galactopoiesis and mammogenesis. Improved lactation performance can be achieved under conditions that enhance mammary cell proliferation (or decrease apoptosis), biochemical and structural differentiation of mammary epithelium, and synthesis and secretion of milk components. Moreover, any factors involved in the regulation of these processes can directly impact mammary function and milk yield [17, 18].

The increase in body weight observed in metoclopramide treatment could also have been from a possible direct action of the drug on the hepatocytes resulting in the secretion of insulin like growth factor-1, which is known to cause the growth of tissues in the body. The increase in body weight could also have been due to a possible upregulation of GH receptors in the various tissues, hence elevating their susceptibility and response to secreted GH. Although Olanzapine and Risperidone both act on dopaminergic cells like metoclopramide, the mild effect observed in the GH level of these treated groups could have been from their less affinity to D₂ receptors within the adenohypophysis resulting in less effect on the serum GH level compared to the control and metoclopramide. The result obtained on the body weight in both olanzapine and risperidone treated groups also corroborate with the findings on the serum growth hormone level.

3.1 Conclusion

Metoclopramide elevates serum GH level as well as the body weight of dams during lactation in female Wistar rats more than Olanzapine and Risperidone.

Ethical Approval:

The research was carried out according to the guidelines of Ahmadu Bello University animal use and care policy.

This research is an excerpt from a work we conducted a while back as part of a dissertation during which a proposal was presented to the department and approved taking into consideration the University's ethical standards on the use of animals.

As per international standard or university standard ethical approval has been collected and preserved by the authors.

3.2 REFERENCES

1. Tuggle CK, Trenkle A. Control of growth hormone synthesis: domestic Animal. *Endocrinology*. 1996;13(1): 1-33.
2. Butler AA, Le Roith D. Control of growth by the somatotropic axis: Growth hormone and the insulin-like growth factors have related and independent roles. *Annual Review of Physiology*. 2001; 63: 141-164
3. Hadsell DL, Parlow AF, Torres D, George J, Olea W. Enhancement of maternal lactation performance during prolonged lactation in the mouse by mouse GH and long-R3-IGF-I is linked to changes in mammary signaling and gene expression. *Journal of Endocrinology*. 2008; 198: 61-70
4. Wall E, McFadden T. Regulation of Mammary Development as It Relates to Changes in Milk Production Efficiency. 2012; <http://dx.doi.org/10.5772/50777>
5. Postel-Vinay MC, Kelly PA. Growth hormone receptor signalling. *Baillieres Clinical Endocrinology and Metabolism*. 1996; 10: 323-336
6. Guzman V, Toscano G, Canales ES, Zarate A. Improvement of defective lactation by using oral Metoclopramide. *Acta Obstetrica et Gynecologica Scandinavica*. 1979; 58(1):53–55.
7. Anderson PO, Valdés V. A critical review of pharmaceutical galactagogues. *Breastfeeding Medicine* 2007; 2(4):229–242
8. Zuppa A, Sindico A, Orchi C. Safety and efficacy of galactagogues: substances that induce, maintain, and increase breast milk production. *Journal of Pharmacy and Pharmaceutical Sciences*. 2010; 13(2): 113-187.
9. Ingram J, Taylor H, Churchill C, Pike A, Greenwood R. Metoclopramide or domperidone for increasing maternal breast milk output: a randomized controlled trial. *Archives of Disease in Childhood: Fetal and Neonatal Edition*. 2012; 97(4): F241–F245
10. McCormack PL, Wiseman LR. Olanzapine: a review of its use in the management of bipolar disorder. *Drugs*. 2004; 64: 2709.

- 185 11. Mirena V, Boyko SD, Peychinskam I, Veleva P, Dimitrova PR. Metoclopramide – induced
186 extrapyramidal signs and symptoms – brief review of literature and case report. *Journal of*
187 *IMAB*, 2014; 20(6): 339-342
- 188 12. Lambert M, Haro JM, Novick D, Edgell ET, Kennedy L, Ratcliffe M. Olanzapine vs. other
189 antipsychotics in actual out-patient settings: six months tolerability results from the European
190 Schizophrenia Out-patient Health Outcomes study. *Acta Psychiatrica Scandinavica*. 2005; 111:
191 232-43
- 192 13. Leysen JE, Gommeren W, Eens A, Stoof JC, and Jansen PA. Profile of Risperidone, a
193 new antipsychotic *Journal pharmacology*. 1998; 247: 661-70.
- 194 14. Marthins MI.. *Zaria and its regions*. Department of Geography Ahmadu Bello University,
195 Zaria, pp-41. 2006
- 196 15. Guistina A, Bussi AR, Doga M, Licini M, Schettino M, Wehrenberg WB. Effect of
197 metoclopramide on the growth hormone response to galanin in normal man. *Hormones*
198 *metabolism and Research*. 1993; 25(19): 489-492.
- 199 16. Mortel M, Mehta SD. Systematic review of the efficacy of herbal galactogogues *Journal*
200 *of Human Lactation*. 2013; 29(2): 154–162
- 201 17. Akers RM. Major advances associated with hormone and growth factor regulation of
202 mammary growth and lactation in dairy cows. *Journal of Dairy Science*. 2006; 89: 1222-
203 1234.
- 204 18. Lieberman JA, Tollefson G, Tohen M, Green AI, Gur RE, Kahn R. Comparative efficacy
205 and safety of atypical and conventional antipsychotic drugs in first episode psychosis: a
206 randomized, double-blind trial of Olanzapine versus haloperidol. *American Journal*
207 *Psychiatry*. 2003; 160: 1396-404.